

NEAT (North East Atlantic Taxa):

South Scandinavian marine & maritime Chelicerata & Uniramia

Check-List

compiled at TMBL (Tjärnö Marine Biological Laboratory) by:

Hans G. Hansson 1990-04-11 / small revisions until Nov. 1994, when it for the first time was published on Internet.
Published as a pdf file February 1996 and again August 1998.

Citation suggested: Hansson, H.G. (Comp.). 1998. NEAT (North East Atlantic Taxa): South Scandinavian marine & maritime Chelicerata & Uniramia. Check-List. Internet pdf Ed., Aug. 1998. [<http://www.tmbi.gu.se>].

Denotations: (TM) = Genotype @ = Associated to * = General note

N.B.: This is one of several preliminary check-lists, covering S. Scandinavian marine animal (and partly marine protist) taxa. Some financial support from (or via) NKMB (Nordiskt Kollegium för Marin Biologi), during the last years of the existence of this organisation (until 1993), is thankfully acknowledged. The primary purpose of these checklists is to facilitate for everyone, trying to identify organisms from the area, to know which species that earlier have been encountered there, or in neighbouring areas. A secondary purpose is to facilitate for non-experts to find as correct names as possible for organisms, including names of authors and years of description. So far these checklists are very preliminary. Due to restricted access to literature there are (some known, and probably many unknown) omissions in the lists. Certainly also several errors may be found, especially regarding taxa like Plathelminthes and Nemata, where the experience of the compiler is very rudimentary. The credibility of some other lists, containing taxa like e.g. Porifera, where, at least in certain subtaxa, taxonomic confusion seems to prevail, is certainly also suboptimal so far.

The compiler must admit his almost definite practical inexpérience of Chelicerata and Uniramia, so I guess that this list is very much more incomplete than the lists dealing with marine animals s.str. The borderline between animals living more or less permanently in the marine littoral and temporary guests in this environment is also more or less artificial, so some of the names in this list should probably not have been included.

Species found in South Scandinavia, as well as from neighbouring areas, chiefly the British Isles, have been considered, as some of them may show to have a slightly more northern distribution, than what is known today. However, species with a typical Lusitanian distribution, with their northern distribution limit around France or Southern British Isles, have as a rule been omitted here, albeit a few species with probable northern limits around the British Isles are listed here until distribution patterns are better known. Arctic Taxa, known only from areas north of N Norway are probably not as well represented in this list as they should be. The compiler would be very grateful for every correction of presumptive lapses and omissions an initiated reader could make.

N.B. This document is intended to be completely unavailable regarding nomenclature within the meaning of ICZN.

ARTHROPODA Latreille,1829

* Taxa within the remaining sub-phylum, Crustacea, are treated in a separate list.

CHELICERATA Heymons,1901

MEROSTOMATA Woodward,1866

XIPHOSURIDA R. & E. Richter,1924
= **LIMULIDA** "Dolksvansar"

Xiphosuridae
= **Limulidae**

Xiphosurinae Pocock,1902

Limulus O.F. Müller,1785
= **Xiphosura** Gronovius,1764 (Is this name rejected for any reason?)

L. polyphemus (Linnaeus,1758)
* American N. Atlantic species, found now and then in Europe
S. Baltic, Skagerrak, North Sea, W. N. Atlantic

ARACHNIDA Cuvier,1812 "Spindeldjur"

* Only "marine families" included

ARANAEAE Clerck,1758 "Spindlar"

* There are no strictly marine spiders in Europe; those in the following list may be found intertidally.

ENTELEGYNAE

Araneoidea

Linyphiidae Blackwall,1859
= **Erigonidae** Gerhardt,1923
= **Micryphantidae** Bertkau,1872 "Täckvävare"

Cneophalocotes Simon,1884

C. curtis Simon,1884?
England

Erigone Audouin, in Savigny,1826

E. arctica (White,1852) **maritima** Kulczinski,1902
Böhnsan (e.g. Saltö)-Uppland, Norway, Finnland, Denmark, Germany, Belgium, Br. Isles, Arctic shores etc.

E. longipalpis (Sundevall,1830)
Skåne, Norway, Finland, Denmark, Germany, Br. Isles, Arcachon, etc.

Halorates Hull,1911

H. reprobus (O. Pickard-Cambridge,1879)
* Littoral & in wrack.
Skåne, Norway, Belgium, England, France

Acanthophyma? 19??

* Probably not a recognized genus today?

A. gowerensis (Lockett,19??)
Ireland, Britain (Gower peninsula)

Oedothorax Bertkau,1883

O. fuscus (Blackwall,1834)
Norway, Belgium, Britain

O. apicatus (Blackwall,1850)
*Littoral
Finnland, Norway, Belgium

O. retosus (Westring,1851)
* Littoral & in wrack.
Uppland, Åland, Finnland, Norway, Belgium

Baryphyma Simon,1884
=? **Praestigia** Millidge,1954

B. duffeyi (Millidge,1954)
= **Praestigia duffeyi** Millidge,1954
SE England, Belgium

B. maritimum (Crocker & Parker,1970)
Belgium

Satilatlas Keyserling,1886
=? **Perimones** Jackson,1932

P. britteni (Jackson,1913)
Britain, Belgium

Trichoncus Simon,1884

T. hackmanni Millidge,1955
S England

Lycosoidea Sundevall,1833 "Jaktspindlar"

Lycosidae Sundevall,1833 "Vargspindlar"

Arctosa Koch,1847

A. perita (Latreille,1799)
Europe; e.g. Sweden, Norway, Germany, ?Denmark

A. cinerea (J.C. Fabricius,1777) "Strandjaktspindeln"
= **Trochosa cinerea** (J.C. Fabricius,1777)
Sweden, Norway, Denmark

Pardosa C.L. Koch,1847
= **Lycosa** Latreille,1804 (p.p.)

P. arenicola (F.O. Pickard-Cambridge,18??) **fucicola** (Dahl,1???)
= **Lycosa arenicola** F.O. Pickard-Cambridge,18?? **fucicola** Dahl,1???

* Exclusively littoral.

Uppland, Norway, England, Germany

P. pullata (Clerck,1758)
Sweden, Norway, England, Germany, S. Finland

P. purbeckensis F.O. Pickard-Cambridge,1895
Br. Isles, Belgium

P. ? fluvialis (Blackwall,1???)
= **Lycosa fluvialis** : Auctt.
Småland-Uppland, Finnland

Trochosa Koch,1847

T. ruricola (De Geer,1778)

* Littoral.
Sörmland-Uppland, Norway, Britain, Germany

Dictynoidea F.O. Pickard-Cambridge,1871

Dictynidae F.O. Pickard-Cambridge,1871

Dictyna Sundevall,1833

D. major Menge,1869

N England, Scotland, Belgium

Clubionoidea (Simon,1895) Petrunkevitch,1955

Gnaphosidae Pocock,1898 "Plattbukspindlar"

Haplodrassus Chamberlin,1922

H. minor (F.O. Pickard-Cambridge,1879)

Norway, S England, Germany

Thomisoidea Sundevall,1833

Thomisidae Sundevall,1833 "Krabbspindlar"

Misumena Latreille,1804

M. vatia (Clerck,1758)

= **Araneus vatius** Martyn,1793 [ex Clerck,1757]

Sweden, Norway, Arcachon

Salicidae Blackwall,1841 "Hoppspindlar"

Salticidae Blackwall,1841 "Egentliga hoppspindlar"

Euophrays Koch, in Panzer,1834

E. browningi Millidge & Lockett,1955

SE England

Salicus Latreille,1804

S. mutabilis H. Lucas,1846

Arcachon

S. scenicus (Clerck,1758)

= **Aranea scenica** Linnaeus,1758 ex Clerck

Norway, Belgium, Arcachon

Heliophanus C.L. Koch, in Panzer,1835

H. auratus C.L. Koch, in Panzer,1835

Belgium, Arcachon

H. tribulosus Simon,1868

Belgium, Arcachon

PSEUDOSCORPIONIDA Latreille,1825 "Klokrypare"

* With the exception of gen. **Chthonius**, only species restricted to sea shores have been considered

CHTHONIINA Beier,1932

Chthonioidae Beier,1932

Chthoniidae Daday,1889

Kewochthonius Chamberlin,1929

- K. halberti** (Kew,1916)
England, Ireland, France
- Chthonius** C.L. Koch,1843
* Found also in certain inland localities.
- C. tetrachelatus** (Preyssler,1790)
Bohuslän-Östergötland, Öland, Gotland, Åland, C. Europe, Br. Isles, Denmark, Norway
- C. kewi** Gabbott,1966
Br. Isles
- NEOBISIIINA** Beier,1932
- Neobisiodea** Chamberlin,1931
- Neobisidae** Chamberlin,1931
- Neobisium** Chamberlin,1931
- N. (Neobisium) maritimum** (Leach,1812)
W. & SW. Br. Isles, Ireland, W. France
- CHELIFERINA**
- Cheliferoidea** Chamberlin,1931
- Cheliferidae** Hagen,1879
- Cheliferinae** Simon,1879
- Dactylochelifer** Beier,1932
- D. latreilli** (Leach,1817)
@ *Amnophila arenaria* & *Elymus pycnanthus*
Br. Isles, E. & S. Sweden (e.g. Uppland), Denmark, Finland, W. & Central Europe; e.g. N. Germany
- ACARI** Leach,1817 "Kvalster"
- ACTINOTRICHIDA**
- ASTIGMATIDA**
- Hemisarcopoidea**
- Hyadesiidae** Halbert,1915
= *Glyphagidae*
- Hyadesia** Mégnin,1889 (TM *Hyadesia uncinifer* Mégnin,1889 - Tierra del Fuego)
= *Lentungula* Michael,1893
- H. fusca** (Lohmann,1894)
= *Lentungula fusca* Lohmann,1894
Baltic, Bohuslän, W Ireland, North Sea shores
- H. algivorans** (Michael,1893)
= *Lentungula algivorans* Michael,1893
S England
- Algophagidae**
- Algophagus** Hughes,1955
- Neohyadesia** ??,19??
- Acaroidea**
- Acaridae** Oudemans,1904 "Skabbkvalster"
- Pontoppidania** Oudemans,1923
- P. littoralis** (Halbert,1920)
= *Tyroglyphus littoralis* Halbert,1920
@ in moist decaying seaweeds
W Ireland
- Rhizoglyphus** Claparède,1869
- R. sp.**
Britain
- Tyrophagus** Oudemans,1924
- T. sp.**
Britain
- Suidasiidae**
- Suidasia**
- PSOROPTINA**
@ with few exceptions parasites of birds or mammals
- Psoroptoidea**
- Sarcopidae**
- Sarcopes** Latreille,1802
- ?S. sp.: Jacobsen,1966
@ skin of *Phoca vitulina*
German North Sea, Netherlands
- PROSTIGMATIDA** Kramer,1887
= **TROMBIDIFORMES**
= **ACTINEDIDA**
- Nautarachnidiae**
* What infraorder & superfamily?
- Nautarachna** Moniez,1888
- Thyasidae**
* What infraorder & superfamily?
- Thyopsis** Piersig,1899
- ENDEOSTIGMATINA** Grandjean,1937
- Nanorchestidae** Grandjean,1937
- Nanorchestes** Topsent & Trouessart,1890
- N. amphibius** Topsent & Trouessart,1890
Ireland, Isle of Whigt
- Bimichaelidae**
= **Pachygnathidae**
= **Alycidae** G. Canestrini,1891
- Pachygnathus** Dugès,1834
- P. sp.**
Britain
- Alicus** Berlese,1890
= *Alycus* Koch,1842, non Rafinesque,1815 (Coleoptera)
* This genus is not found in Hayward & Ryland (Eds.) 1990. Synonymized?
- A. (Leptalicus** Berlese,1910) **oblongus** Halbert,1920
W Ireland
- A. latus** Halbert,1920
W Ireland
- EUPODINA**
- Bdelloidea**
- Bdellidae** Dugès,1834
- Bdella** Latreille,1795
- B. interrupta** G.O. Evans,1952
* In crevices of the rocky littoral.
Br. Isles
- B. capillata** Kramer,1881
Scandinavia, Germany, Britain, Clare Island, Italy
- B. decipiens** Thorell,1871
= *Bdella vulgaris* (Hermann,1???) var. *littoralis* Moniez,1890
Sweden, W Ireland, Arcachon, Siberia
- B. sanguinea** Trouessart,1???
?Germany
- B. littoralis** (Linnaeus,1758)
= *Molgus littoralis* (Linnaeus,1758) (TM of *Molgus* Dujardin,1842)
= *Molgus acutus* (Thorell,1871)
= *M. villosus* (Kramer,1883)
= *M. basteri* (Michael,1896)
* An up to 3.5 mm long, shining red species, on decaying seaweed.
Scotland, Norway, Ireland
- Cyta** von Heyden,1826
- C. latirostris** (Hermann,1???)
W Ireland
- Hoplomolgus** Berlese,1923
- H. sp.**
Britain
- Neomolgus** Oudemans,1938
- N. sp.**
Britain
- Eupodoidea**
- Eupodidae** C.L. Koch,1842
- Eupodes** Koch, in Panzer,1835
- E. halophilus** Halbert,1920
England, W Ireland
- Penthaleidae** Oudemans,1931
- Hylotydeus** Berlese,1891
- H. hydrodromus** (Berlese & Trouessart,1889)
= *Notophallus hydrodromus* Berlese & Trouessart,1889
Br. Isles
- H. hydrodromus** (Berlese & Trouessart,1889) var. *albolineatus* Halbert,1915
W Ireland
- Chromotydaeus** Berlese,1903
* This genus is not found in Hayward & Ryland (Eds.) 1990. Synonymized? - Correct family?
- C. ovatus** (C.L. Koch,1838)
= *Penthalodus ovatus* C.L. Koch,1838
@ among *Semibalanus*, *Fucus spiralis* & *F. vesiculosus*
W Norway, Germany, Ireland, Britain, Italy
- Penthalodiidae** Sig Thor,1933
- Penthalodes** Murray,1877
- P. sp.**
Britain
- Rhagidiidae** Oudemans,1922
- Foveacheles** Zacharda,1???
= *Rhagidia* Thorell,1871
- F. halophila** (Laboulbène,1851)
= *Gamasus halophilus* Laboulbène,1851
= *Normeria halophila* : Moniez,1889
W Ireland, ?French Atlantic coast
- Robustochelus** Zacharda,1???
- R. sp.**
Britain
- Tydeoidea**

- Tydeidae** Kramer,1877
- Lasiotydaeus** Berlese,1908 (TM*L. glycyphaginus* Berlese,1908?)
- L. (Melanotydaeus** Berlese,1910) **brevistylus** Halbert,1920
W Ireland
- Retetydeus** Thor,1931
- R. sp.**
Britain
- Nematalycoidea**
- Nematalycidae**
- Halacaroidea**
- Halacaridae** Murray,1877 "Havskvalster"
- Rhombognatinae** Viets,1927
- Isobactrus** Newell,1947
- I. levis** (Viets,1927)
= *Rhombognathus levis* Viets,1927
Baltic, Bohuslän, North Sea
- I. setosus** (Lohmann,1889)
= *Aletes setosus* Lohmann,1889
Baltic, Halland, North Sea
- I. uniscutatus** (Viets,1939)
= *Rhombognathus uniscutatus* Viets,1939
Baltic, North Sea
- I. hartmanni** Bartsch,1972
* Not included in Green & Macquitty, 1987
Öresund, Helgoland
- I. ungulatus** Bartsch,1975
Roscoff, Falmouth, Isle of Man, Pembrokeshire
- Metarhombognathus** Newell,1947
- M. armatus** (Lohmann,1893)
= *Rhombognathus mollis* Viets
Baltic, North Sea
- M. nudus** (Viets,1928)
= *Rhombognathus conctectus* Viets
Cullercoats, Scotland, Murmansk
- Rhombognathides** Viets,1927
- R. mucronatus** (Viets,1927)
Bornholm, North Sea, Norway
- R. pascens** (Lohmann,1889)
Baltic, Bohuslän, North Sea, Norway
- R. seahami** (Hodge,1860)
= *Pachynathus seahami* Hodge,1860
Baltic, Bohuslän, Norway, North Sea, Kola peninsula, Novaja Zemlja, Iceland, Shetlands, Ireland, E & S Britain, E USA
- R. spinipes** (Viets,1933)
Baltic, North Sea
- R. merrimani** Newell,1947
Bohuslän, Norway, England, Wales, Ireland
- R. exoplus** (Trouessart,1901)
* Not included in Green & Macquitty, 1987
Bretagne, North Sea
- R. trionyx** (Trouessart,1899)
* Not included in Green & Macquitty, 1987
Bretagne
- Rhombognathus** Trouessart,1888
- R. notops** (Gosse,1855)
Baltic, Bohuslän, North Sea
- R. subtilis** Bartsch,1975
Norway, Spitzbergen, Cullercoats, Northumberland
- R. lioy whole** Trouessart,1900
Baltic, Bergen
- R. magnirostris** Trouessart,1889
Kullen, Bergen, French mediterranean
- R. cetratus** Bartsch,1975
Elbe-estuary
- R. cryptorhynchus** Trouessart,1901
* Not included in Green & Macquitty, 1987
Bretagne
- R. plumifer** ?
?
- R. intermedius** Schulz,1933
Kieler Bucht, Helgoland
- R. procerus** ?
?
- Actacarinae** Viets,1939
- Actacarus** Schulz,1936
- A. obductus** Bartsch,1977
* Not included in Green & Macquitty, 1987
Bohuslän, North Carolina
- A. pygmaeus** Schulz,1936
* Not included in Green & Macquitty, 1987
Kieler Bucht, North Sea, Mediterranean, Black Sea
- A. bacescui** Konnerth-Ionescu,1970
* Not included in Green & Macquitty, 1987
N Spain-Mediterranean-Black Sea
- Halacarinae** Viets,1927
- Agauopsis** Viets,1927
- A. brevipalpus** (Trouessart,1889)
= *Agae brevipalpus* Trouessart,1889
England, Wales, Ireland, France
- A. tricuspis** Benard,1962
= *A. vidua* Krantz,1970
England, Bretagne
- A. aculeata** (Trouessart,1896)
* Not included in Green & Macquitty, 1987
Biscay in 1410m among corals
- A. microrhyncha** (Trouessart,1889)
* Not included in Green & Macquitty, 1987
Calais
- A. hirsuta** (Trouessart,1889)
* Not included in Green & Macquitty, 1987
Mediterranean
- Agae** Lohmann,1889
- A. chevreuxi** (Trouessart,1889)
* Not included in Green & Macquitty, 1987
Roscoff
- A. panopae** (Lohmann,1893)
* Not included in Green & Macquitty, 1987
Monaco
- Anomalohalacarus** Newell,1949
- A. anomala** (Trouessart,1894)
* In sublittoral sediments.
Bohuslän, Skagerrak, Engl. channel, Bretagne
- A. intermedius** Bartsch,1976
* Shallow water species.
North Sea, Bretagne, Ireland
- A. marcandrei** Monniot,1967
* In sublittoral sediments.
Bohuslän, Skagerrak, Bretagne, Yorkshire
- A. minutus** Bartsch,1976
* Shallow water species.
Baltic, North Sea, Bretagne, Yorkshire
- A. poizati** Bartsch,1985
* In sublittoral sediments.
Bohuslän, Yorkshire
- A. acnemus** Bartsch,1985
* In sublittoral sediments.
Bohuslän, Yorkshire
- A. arenarius** Bartsch,1978
* Shallow water species. Not included in Green & Macquitty, 1987
North Sea
- A. similis** Bartsch,1976
* Not included in Green & Macquitty, 1987
?
- A. septentrionalis** Bartsch,1991
* In sublittoral sediments.
Bohuslän
- Arhodeoporus** Newell,1947
- A. gracilipes** (Trouessart,1889)
Bohuslän, North Sea, Tromsö, England
- A. minor** Bartsch,1977
Bretagne, Yorkshire, Ireland
- Coloboceras** Trouessart,1889
- C. drachi** Monniot,1962
Ireland
- C. longiusculus** Trouessart,1889
* Not included in Green & Macquitty, 1987
Bohuslän, French atlantic coast
- C. kochleri** Trouessart,1896
* Not included in Green & Macquitty, 1987
Biscay in 1410m among corals
- Copidognathus** Trouessart,1888
- C. brevirostris** Viets,1927
Baltic, Bohuslän, Bergen, North Sea
- C. brifaciuss** Bartsch,1989
Bohuslän
- C. dentatus** Viets,1940
Grimsby, Kent
- C. fabricii** (Lohmann,1889)
= *Halacarus fabricii* Lohmann,1889
Baltic, Halland, Bohuslän, North Sea, Tromsö, Iceland, Faeroes, Ireland, W France, Mediterranean, Black Sea, Azores, Cape Verde
- C. gibbus** (Trouessart,1889)
Helgoland, Engl. Channel
- C. granulatus** (Hodge,1863)
England, French atlantic coast
- C. glyptoderma** Trouessart, ?
Plymouth??

- C. hartwigi* Bartsch,1979
Irish Sea, Bermuda
- C. lamellosus* (Lohmann,1893)
= *C. tabellio* (Trouessart,1894)
Engl. Channel, Yorkshire, Ireland
- C. latisetus* Viets,1940
Bohuslän, North Sea, Bretagne, Yorkshire, Kent, Ireland
- C. loricifer* André,1946
Baltic, Engl. Channel
- C. magnipalpus* (Police,1909)
* Not included in Green & Macquitty, 1987
Bohuslän, Mediterranean
- C. oculatus* (Hodge,1863)
Baltic, Kattegatt, Bohuslän, North Sea, Engl. Channel
- C. poucheti* (Trouessart,1894)
= *Halacarus poucheti* Trouessart,1894
Iceland, Labrador
- C. remipes* (Trouessart,1894)
Engl. Channel, Ireland
- C. reticulatus* (Trouessart,1893)
= *Halacarus reticulatus* Trouessart,1893
Bohuslän (Svartskär N of Kalkgrund - Bartsch, unpubl.), Norway, Spitzbergen, Iceland, Ireland
- C. rhodostigma* (Gosse,1855)
Baltic, Halland, Bohuslän, North Sea, England, Scotland, Ireland
- C. septentrionalis* (Halbert,1915)
= *Halacarus (Copidognathus) lamellosus* Lohmann,1893 var. *septentrionalis* Halbert,1915
Engl. Channel, Clew Bay
- C. steveici* Bartsch,1976
@ Ectoparasite on eggs of *Maia squinado*
Engl. Channel
- C. tectirostris* Bartsch,1979
Bohuslän, Bretagne, Yorkshire, Irish Sea, Mediterranean
- C. gracilipes* (Trouessart,1889)
* Not included in Green & Macquitty, 1987
N. Kattegatt, Bohuslän, Helgoland, Bergen
- C. caudani* (Trouessart,1896)
* Not included in Green & Macquitty, 1987
Biscay
- C. humerosus* (Trouessart,1896)
* Not included in Green & Macquitty, 1987
Engl. Channel
- C. latus* Viets,19??
* Not included in Green & Macquitty, 1987
Öresund, Bergen
- Halacarus* Gosse,1855
- H. bisulcus* Viets,1927
Bohuslän, Helgoland, Bergen, Iceland, Faeroes, Yorkshire, Devon, Ireland
- H. anomalus* Trouessart,1894
* Not included in Green & Macquitty, 1987
Baltic, Helgoland
- H. ctenopus* Gosse,1855 (TM)
North Sea, Shetland, Engl. Channel, Ireland
- H. actenos* Trouessart,1889
Engl. Channel, Wales, Ireland
- H. borealis* Trouessart,1893
NW Iceland
- Thalassarachna* Packard,1871
= *Halacarellus* Schulz,1933
- T. baltica* (Lohmann,1889)
Baltic, Bohuslän, North Sea
- T. basteri* (Johnston,1836)
= *Acarus basteri* Johnston,1836
= *T. verrilli* Packard,1871
= *Halacarus spinifer* Lohmann,1889
Baltic, Bohuslän, North Sea, Iceland, Murmansk, Novaja Zemlya, Greenland, Faeroes, Ireland, E & S England, France, U.S.A
- T. capuzina* (Lohmann,1893)
= *Halacarus (Halacarus) areodus* Halbert,1915
Baltic, Öresund, Bohuslän, North Sea, England, Scotland, Ireland
- T. petiti* (Angelier,1950)
Kieler Bucht, Bretagne, Humber estuary
- T. procura* (Viets,1927)
Baltic, Halland, Helgoland, Bretagne, Ireland
- T. southerni* (Halbert,1915)
= *Halacarus (Halacarus) southerni* Halbert,1915
Bohuslän, Yorkshire, Clew Bay, Irish Sea
- T. striatus* (Lohmann,1889)
= *Halacarus similis* Viets,1927
Kieler Bucht, Bohuslän, Bergen, Bretagne, England, Ireland
- T. subterranea* (Schulz,1933)
Baltic, North Sea, Norway, Bretagne, Essex
- T. bergensis* (Viets,1927)
* Not included in Green & Macquitty, 1987
Kieler Bucht, Bergen
- T. floridearum* (Lohmann,1889)
* Not included in Green & Macquitty, 1987
Baltic
- T. subcrispus* (Bartsch,1978)
* Not included in Green & Macquitty, 1987
?
- T. inermis* (Trouessart,1888)
* Not included in Green & Macquitty, 1987
Baltic (e.g. Greifswald), Halland, North Sea
- T. longipes* (Trouessart,1888)
* Not included in Green & Macquitty, 1987
Baltic, Bohuslän, North Sea
- T. micropectinatus* (Bartsch,1972)
* Not included in Green & Macquitty, 1987
Schleswig-Holstein
- T. hexacanthus* (Viets,1927)
* Not included in Green & Macquitty, 1987
Kieler Bucht, Öresund, Kattegatt, Bohuslän, Bergen
- T. abyssorum* (Trouessart,1896)
* Not included in Green & Macquitty, 1987
Biscay in 400-1410m among corals
- Camactognathus* Newell,1984
* Correct subfamily?
- C. borealis* Bartsch,1991
Bohuslän
- Acarochelopodinae* Bartsch,1977
- Acarochelopodia* Angelier,1954
- A. delamarei* Angelier,1954
* Not included in Green & Macquitty, 1987
Europe
- Simognathinae* Viets,1927
- Simognathus* Trouessart,1889
- S. leiomerus* Trouessart,1894
= *S. andrei* Monniot,1961
Engl. Channel, Ireland
- S. minutus* (Hodge,1863)
= *Pachynathus sculpus* Brady,1875
Kullen (NW Skåne), Helgoland, Bergen, England, Scotland
- Acaromantis* Trouessart & Neumann,1893
- A. squilla* Trouessart & Neumann,1893
* Not included in Green & Macquitty, 1987
French atlantic coast
- A. minutus* Bartsch,1976
* Not included in Green & Macquitty, 1987
?
- Atelopsalis* Trouessart,1896
- A. tricuspidis* Trouessart,1896
* Not included in Green & Macquitty, 1987
Biscay in 1410m among corals
- Lohmannellinae* Viets,1927
- Lohmannella* Trouessart,1901
- L. falcata* (Hodge,1863)
= *Leptognathus falcatus* Hodge,1863
Baltic, Bohuslän, Norway, North Sea, Shetlands, Hebrides, Ireland, Britain, Iceland, W France, Mediterranean, Black Sea, E North America, (possibly cosmopolitan)
- L. kervillei* (Trouessart,1894)
Engl. Channel, Wales
- L. multiseta* Bartsch,1986
* Not included in Green & Macquitty, 1987
Bohuslän, Engl. Channel, Mediterranean
- L. norvegica* Viets,1927
* Not included in Green & Macquitty, 1987
Bergen
- Porolohmannella* Viets,1933
- P. violacea* (Kramer,1879)
* NB! Freshwater species
Widespread in Europe from Belgium to U.S.S.R.
- Scaptognathus* Trouessart,1889
- S. tridens* Trouessart,1889
Engl. Channel, French atlantic coast
- S. trouessarti* Halbert,1915
Dingle Bay (Ireland)
- S. hallezi* Trouessart,1894
* Not included in Green & Macquitty, 1987
Calais
- Limnohalacarinae* Viets,1927
- Limnohalacarus* Walter,1917
- L. wackeri* (Walter,1914)
* NB! Freshwater species
Widespread in Europe
- Parasoldanellonyx* Viets,1929
- P. parviscutatus* (Walter,1917)
* NB! Freshwater species
Belgium, England, Scotland, Switzerland, Italy, Germany, U.S.S.R.
- Soldanellonyx* Walter,1917
- S. chappuisi* Walter,1917
* NB! Freshwater species

England, Scotland, France to Roumania and U.S.S.R.

S. monardi Walter,1919

* NB! Freshwater species

England, Wales, Scotland, France to Roumania and U.S.S.R.

Porohalacarinae Viets,1933

Caspihalacarus Viets,1928

C. hyrcanus Viets,1928

* NB! Brackish water species

Wales, Caspian and Black Seas, Dnieper

Lobohalacarus Viets,1939

L. dolgarae Green,1954

* NB! Freshwater species

England

L. gallicus (Migot,1926)

*NB! Freshwater species

French Pyrenees

L. weberi (Romijn & Viets,1924)

* NB! Freshwater species

England, Scotland, Widespread in European inland waters

Porohalacarus Thor,1923

P. alpinus (Thor,1910)

* NB! Freshwater species

England, Wales, Widespread in European and N. African waters

ANYSTINA

Anystoidea

Anystidae Oudemans,1902

Chaussieria Oudemans,1937

C. sp.

Britain

Teneriffiidae

Heteroteneriffia Hirst,1926

PARASITENGONINA Oudemans,1909

Erythraeoidae

Erythraeidae Oudemans,1902

Balaustium von Heyden,1826

B. sp.

Britain

Erythraeus Latreille,1806

E. sp.

Britain

Trombidioidea

Podothrombiidae

Podothrombium Berlese,1910

P. filipes (Koch,18??)

Norway, Germany, Ireland, Britain

Trombiculidae

Trombidiidae Leach,1814

Rhyncholophus Dugès,1834

* This genus is not found in Hayward & Ryland (Eds.) 1990. Synonymized? - Correct family?

R. rubripes Berlese & Trouessart,1889

= *R. mineatus* var. *rubripes* Berlese & Trouessart,1889

= *Ritteria hirsuta* George,1910

@ Commensal on *Semibalanus balanoides*.

W Ireland, Britain, France, ?Germany

R. norvegicus (Sig. Thor,1900)

= *Ritteria norvegica* Sig. Thor.

Norway, Ireland

R. tardus Halbert,1915

W Ireland

R. passerinii (Berlese,1904)

= *Erythraeus passerinii* Berlese,1904

W Ireland, Italy

R. araneoides (Berlese,1910)

= *Achorolophus araneoides* Berlese,1910

W Ireland, Sicily

Microtrombidium Haller,1882

M. pusillum (Hermann,18??) var. *major* Halbert,1920

W Ireland

Hydryphantoidea

Hydryphantidae

@ larvae on Odonata, Dipera, Hemiptera, Plecoptera & Trichoptera

Hydryphantes Koch,1842

Hydrodromidae

@ larvae on Chaoboridae, Culicidae & Chironomidae

Hydrodroma Koch,1837

Hygrobatoidea

Pontarachnidae Sig Thor,1929

* In the littoral zone of marine waters, but probably not N of the Mediterranean.

Pontarachna Philippi,1840

Litarachna Walter,1925

Pionidae

Piona Koch,1842

ELEUTHERERGONINA Oudemans,1909

RAPHIGNATHAE

Raphignathoidea

Raphignathidae Kramer,1877

Raphignathus Dugès,1834

* This genus is not found in Hayward & Ryland (Eds.) 1990.

R. scutatus Halbert,1920

W Ireland

Stigmaeidae Oudemans,1931

Stigmacus Koch,1835

S. rhodomelas var. *fissuricola* Halbert,1920

W Ireland

Cheylostigmaeus Willmann,19??

C. sp.

Britain

Tetranychoidae

Tetranychidae

Monoceronychus McGregor,1945

Schizotetranychus Trägårdh,1915

Tenuipalpidae

=? *Phytopiptalpidae*

=? *Pseudoleptidae*

Dolichotetranychus Sayed,1938

Pseudoleptus Bruyant,1911

CRYPTOSTIGMATIDA

= *QRIBATIDA*

= *ORIBATEI* Dugès,1834

Zachvatkina ??,19??

* Correct suborder?; Family?

Z. sp. cf. stercorarii (Dubinin,1949)

@ *Stercorarius parasiticus*

British Columbia

Cymbaeremaidae

* Which superfamily?

Cymbaemaeus Berlese,1896

C. sp.

@ *Uria aalge*

British Columbia

CIRCUMDEHISCENTIAEINA

?Superfamily?

Oribata Latreille,1802

* The species in this genus should probably go to other oribateid genera.

O. setosa C.L. Koch,18??

W. Ireland

O. quadricornuta Michael,18??

W Ireland

O. lucasii Nicolet,18??

W Ireland

Ameronothroidea

Ameronothridae Willmann,1931

Ameronothrus Berlese,1896 (TM *Eremaeus lineatus* Thorell,18??)

A. marinus Banks,18??

* On rocks in intertidal zone.

France

A. bilineatus Michael,18??

* Alivorous species restricted to littoral or estuarine situations.

Br. Isles

A. schneideri (Oudemans,1905)

* Coastal

Borkum

A. sp. aff. oblongatus Sitnikova,1975

@ *Uria aalge*

British Columbia

Fortuninia van der Hammen,1960

Hygroribates Jacot,1934

Selenoribates Strenzke,1961

Carabodoidea

Carabodidae Willmann,1931

<i>Carabodes</i> C.L. Koch, in Panzer, 1835 (TM <i>C. coriaceus</i> Koch, in Panzer, 1835)	<i>P.</i> sp. Britain
<i>C.</i> sp. Britain	
<i>Passalozetes</i> Grandjean, 1932	
<u>Ceratozetoidea</u>	
<i>Ceratozetidae</i> Jacot, 1925	
<i>Ceratozetes</i> Berlese, 1908 (TM <i>Oribata gracilis</i> Michael, 1884)	
<i>C.?</i> <i>quadrivertex</i> (Halbert, 1920) = <i>Oribata quadrivertex</i> Halbert, 1920 W Ireland	
<i>Trichoribates</i> Berlese, 1910 (TM <i>Murcia trimaculata</i> C.L. Koch, 1836)	
<i>T.</i> sp. Britain	
?Families?	
<i>Chamobates</i> Hull, 1916 (TM <i>Oribata cuspidata</i> Michael, 1884)	
<i>C.</i> sp. Britain	
<i>Euzetes</i> Berlese, 1908 (TM <i>Oribata globula</i> Nicolet, 1855)	
<i>E.</i> sp. Britain	
<i>Fuscozetes</i> Sellnick, 1928 (TM <i>Oribates fuscipes</i> C.L. Koch, 1844)	
<i>F.</i> sp. Britain	
<i>Minunthozetes</i> Hull, 1916 (TM <i>Oribata fusigera</i> Michael, 1884 = <i>Zetes semirufus</i> C.L. Koch, 1840)	
<i>M.</i> sp. Britain	
<i>Mycobates</i> Hull, 1916	
<i>M. parmeliae</i> (Michael, 1884) (TM) = <i>Oribata parmeliae</i> Michael, 1884 * Under lichens growing on granite rocks Cornwall, W Ireland	
<i>Punctoribates</i> Berlese, 1908 (TM <i>Oribates punctum</i> C.L. Koch-Berl., 1840)	
<i>P.?</i> <i>avenifera</i> Michael, 18?? = <i>Oribata avenifera</i> Michael, 18?? W Ireland	
<i>Sphaerozetes</i> Berlese, 1885 (TM <i>Oribates orbicularis</i> C.L. Koch, 1836)	
<i>S.</i> sp. Britain	
<u>Galumnoidea</u>	
<u>Galumnidae</u> Grandjean, 1936	
<i>Galumna</i> von Heyden, 1826 (TM <i>Notaspis alatus</i> Hermann, 1804)	
<i>G.</i> sp. Britain	
<u>Liacaroidea</u>	
<i>Adoristes</i> Hull, 1916 (TM <i>Oribates ovatus</i> C.L. Koch, 1840)	
<i>A.</i> sp. Britain	
<i>Xenillus</i> Robineau-Desvoidy, 1839 (TM <i>X. clypeator</i> R.-D., 1839 = <i>Notaspis tegeocranus</i> Hermann, 1804)	
<i>X.</i> sp. Britain	
<u>Oppoidea</u>	
<i>Oribella</i> Berlese, 1908 (TM <i>Notaspis pectinata</i> Michael, 1885)	
<i>O.</i> sp. Britain	
<u>Eremaeidae</u>	
<i>Halozetes</i> Berlese, 1917	
<i>Oppia</i> C.L. Koch, 1835 (TM <i>O. nitens</i> C.L. Koch, 1835)	
<i>O. maritima</i> (Willmann, 1929) @ in seaweed Cuxhaven	
<i>O.</i> sp. Britain	
<i>Thalassozetes</i> Schuster, 1963	
<u>Oribatelloidea</u>	
<u>Oribatellidae</u> Jacot, 1925	
<i>Oribatella</i> Banks, 1895 (TM <i>O. quadridentata</i> Banks, 1895?)	
<i>O. maritima</i> v. <i>litoralis</i> Strenzke, 1952 ?	
<i>O.</i> sp. Britain	
?Family?	
<i>Parachipteria</i> van der Hammen, 1952	
	<i>P.</i> sp. Britain
	<u>Oribatuloidea</u>
	<i>Oribatulidae</i> Jacot, 1929
	<i>Haloribatula</i> Schuster, 1957
	<i>Liebstadia</i> Oudemans, 1906 (TM <i>Notaspis similis</i> Michael, 1888)
	<i>L.</i> sp. Britain
	<i>Oribatula</i> Berlese, 1895 (TM <i>Notaspis tibialis</i> Nicolet, 1855)
	<i>O. saxicola</i> Halbert, 1920 W Ireland
	<i>O. similis</i> Michael, 18?? * Is this identical with the genotype of <i>Liebstadia</i> Oudemans? W Ireland
	<i>O. venusta</i> Berlese, 1908 = <i>Notaspis venustus</i> : Halbert, 1915 Norway, Ireland
	<u>Scheloribatidae</u> Grandjean, 1933
	<i>Scheloribates</i> Berlese, 1908 (TM <i>Zetes latipes</i> C.L. Koch, 1844)
	<i>S.</i> sp. Britain
	<u>Passalozetoidea</u>
	<i>Scutovortex</i> Michael, 1879 (TM <i>S. sculptus</i> Michael, 18??)
	<i>S. spoofi</i> Oudemans, 1900 = <i>S. bilineatus</i> : Oudemans, 1901, non Michael, 18?? = <i>Ameronothrus spoofi</i> Auct. * Alivorous species restricted to littoral or estuarine situations. Finland, Br. Isles
	<i>S. bidactylus</i> Coggi, 1900 * in maritime sand dunes Norderney, Ireland, Netherlands, Italy
	<i>S. bilineatus</i> Michael, 18?? W Ireland, England
	<i>S. corrugatus</i> Michael, 18?? =? <i>Eremaeus lineatus</i> Thorell, 18?? W Ireland
	<i>S. perforatus</i> Berlese, 1910 Ireland
	<i>S. maculatus</i> Michael, 18?? @ in <i>Lichina pygmaea</i> W Ireland
	<u>Pelopoidea</u>
	<u>Pelopsidae</u> Ewing, 1917
	<i>Pelops</i> C.L. Koch, 1835, non Gistl, 1834 (Coleoptera) (TM <i>Notaspis hirsutus</i> C.L. Koch, 18??)
	<i>P.</i> sp. Britain
	<u>Eupelopsidae</u> Ewing, 1917
	<i>E.</i> sp. Britain
	<i>Peloptulus</i> Berlese, 1908
	<u>EUPTYCTIMINA</u>
	<u>Phitracaroidea</u>
	<i>Steganacarus</i> Ewing, 1917
	<i>S.</i> sp. Britain
	<u>NOTHRONATINA</u>
	<u>Nanhermannidae</u>
	* Which superfamily?
	<i>Nanhermannia</i> Berlese, 1913
	<u>Nothroidea</u>
	<u>Camisiidae</u> Sellnick, 1928
	<i>Camisia</i> von Heyden, 1826 (TM <i>Notaspis segnis</i> Hermann, 1804)
	<i>C.</i> sp. Britain
	<i>Platynothrus</i> Berlese, 1913 (TM <i>Nothrus peltifer</i> C.L. Koch, 1840)
	<i>P.</i> sp. Britain
	<u>Nothridae</u>
	<i>Nothrus</i> Koch, in Panzer, 1835 * This genus is not found in Hayward & Ryland (Eds.) 1990. Synonymized? - Correct family?
	<i>N. invenustus</i> Michael, 18?? @ in <i>Lichina pygmaea</i> W Ireland
	<u>Hermannioidea</u>
	<u>Hermannniidae</u> Sellnick, 1928

- | | |
|--|---|
| <i>Hermannia</i> Nicolet,1855 (TM <i>Hermannia crassipes</i> Nicolet,1855 = <i>Nothrus gibbus</i> Koch,1840) | Britain |
| <i>H. scabra</i> (L. Koch,18??)
= <i>H. nodosa</i> Michael,18??
Finland, Sweden, Germany, Br. Isles, W Ireland, Novaja Zemlja, Siberia | <i>Pergamasus</i> Berlese,1904 |
| <i>H. reticulata</i> Thorell,18??
Netherlands, Britain, W Ireland, E Greenland, Spitsbergen, Novaja Zemlja | <i>P.</i> sp.
Britain |
| <u>ANACTINOTRICHIDA</u>
= <u>PARASITIFORMES</u> Reuter,1909
= <u>GAMASIIFORMES</u>
= <u>ANACTINOTRICHIDA</u> | <i>Trachygamasus</i> Berlese,1906 |
| <u>MESOSTIGMATIDA</u> Canestrini,1891
= <u>GAMASIDA</u> | <i>T.</i> sp.
Britain |
| <u>UROPODINA</u> Kramer,1881 | <i>Vulgarogamasus</i> Tichomirov,19?? |
| <u>Uropodidae</u> Berlese,1917 "Sköldkvalster" | <i>V.</i> sp.
Britain |
| <u>Dendrouropoda</u> Willmann,1959 | <u>DERMANYSSINA</u> |
| <i>Dentibaiulus</i> ??,19?? | <u>Dermanyssoidae</u> |
| <i>Phaulodinychus</i> Berlese,1905
= <i>Haluropoda</i> Halbert,1915 | <u>Halacaridae</u> |
| <i>P. minor</i> (Halbert,1915)
= <i>Haluropoda minor</i> Halbert,1915
= <i>Trachuropoda minor</i> Halbert,1920
* In decaying wrack.
W Ireland | <i>Halarachne</i> Allman,1847 |
| <i>P. orchesitiidarum</i> (Barrois,1887)
= <i>Uropoda orchesitiidarum</i> Barrois,1887
* In decaying wrack.
W Ireland | <i>H. halichoerii</i> Allman,1847
@ nostrils of <i>Halichoerus gryphus</i>
Baltic, Ireland |
| <i>P. repletus</i> Berlese,1905
= <i>Haluropoda interrupta</i> Halbert,1915 (TM of <i>Haluropoda</i> Halbert,1915 - Ireland) | <i>H. miroungae</i> Ferris,1925
@ <i>Phoca vitulina</i>
US Pacific |
| Ireland | <i>Orthohalarachne</i> Newell,1947
@ walruses & eared seals |
| <i>Phaulocyliba</i> Berlese,1905
* This genus is not found in Hayward & Ryland (Eds.) 1990. Synonymized? | <i>O. diminuta</i> (Doetschman,1944)
@ <i>Callorhinus ursinus</i> & <i>Zalophus californianus</i>
Bering's Sound-C California |
| <i>P. littoralis</i> (Trouessart,1902)
= <i>P. berlesei</i> Halbert,1915
Ireland, Britain | <i>O. attenuata</i> (Banks,1910)
@ <i>Callorhinus ursinus</i> & <i>Eumetopias jubata</i>
Bering's Sound-Oregon |
| <i>Dinychus</i> Kramer,1886 | <u>Laelapidae</u> Berlese,1892 |
| <i>D.</i> sp.: Halbert,1920
W Ireland | <i>Arctoseius</i> Thor,1930
* Described from Spitsbergen |
| <i>Uropoda</i> Latreille,1806 | <i>Aspidilaelaps</i> Trägårdh,1949 |
| <i>U.</i> sp.
Britain | <i>Gammaridacarus</i> Canaris,1962
@ <i>Orchestoidea</i> |
| <i>Urosternella</i> Berlese,1918 | <i>Laelaps</i> Koch,1836
* This genus is not found in Hayward & Ryland (Eds.) 1990. Synonymized? - Correct family? |
| <u>Thinozerconidae</u> Halbert,1915 | <i>L. dentatus</i> Halbert,1920
W Ireland |
| <i>Thinozercon</i> Halbert,1915 | <i>Hypoaspis</i> Canestrini,1884 |
| <i>T. michaeli</i> Halbert,1915
Ireland | <i>H.</i> sp.
Britain |
| <u>Polyaspidae</u> Berlese,1917 | <i>Ololaelaps</i> Berlese,1905 |
| <i>Trachytes</i> Michael,1894 | <i>O.</i> sp.
Britain |
| <i>T.</i> sp.
Britain | <i>Pseudoparasitus</i> Oudemans,1902 |
| <u>PARASITINA</u> | <i>P.</i> sp.
Britain |
| <u>Parasitidae</u> Oudemans,1901 "Parasitkvalster" | <u>Rhinonyssidae</u> |
| <i>Eugamasus</i> Berlese,1892
* This genus is not found in Hayward & Ryland (Eds.) 1990. Synonymized? It was used as a subgenus together with <i>Gamasus</i> by Halbert,1915. | <i>Rhinonyssus</i> Trouessart,1884 |
| <i>E. immanis</i> (Berlese,1903)
* Collected from tidal drift; feeding on small Oligochaetes.
Norway, Millport, Northumberland, Ireland | <i>R.</i> sp.
@ <i>Uria aalge</i>
British Columbia |
| <i>E. trouessarti</i> (Berlese,1892)
= <i>G. thalassinus</i> Berlese & Trouessart,1889
= <i>G. fucorum</i> var. β Berlese & Trouessart,1889
Norway, Northumberland, Ireland, France | <u>Rhodacaroidea</u> |
| <i>Parasitus</i> Latreille,1795 | <i>Rhodacaridae</i> Oudemans,1902 |
| <i>P. kembensi</i> Oudemans.,19??
* In decaying wrack.
Br. Isles | <i>Rhodacarineae</i> Oudemans,1902 (sensu Lee,1970) |
| <i>P. flevensis</i> Oudemans,1922
* In wrack
Zuidzee coast | <i>Rhodacarus</i> Oudemans,1902 |
| <i>Cornigamasus</i> Evans & Till,19?? | <i>strenzkei</i> Willmann,1957
* intertidal
Kiel |
| <i>C.</i> sp.
Britain | <i>R. roseus</i> Oudemans,1902
@ "orange lichen zone"
Ireland |
| <i>Gamasodes</i> Oudemans,1939 | <i>R. pallidus</i> Hull,1918
* In intertidal crevices.
Kiel, Northumberland, Scotland, W Ireland, Spain |
| <i>G.</i> sp.
Britain | <i>Rhodacarellus</i> Willmann,1935 |
| <i>Holoparasitus</i> Oudemans,1936 | <i>R. corniculatus</i> Willmann,1935
@ "beach"
Kiel |
| <i>H.</i> sp.
Britain | <i>R. silesiacus</i> Willmann,1936
@ intertidal sand
?Europe? |
| <i>Paragamasus</i> Hull,19?? | <i>Rhodacaropsis</i> Willmann,1935 |
| <i>P.</i> sp. | <i>R. inexpectata</i> Willmann,1935
@ coastal ground water
Kiel |
| | <u>Cyrtolaelapidae</u> |
| | <i>Hydrogammasus</i> Berlese,1902 |

- H. salinus*** (Laboulbène,1851?)
=? *H. littoralis* (G. & R. Canestrini,1885)
* On *Ascophyllum* and in rock-crevices of high water zone; feeds on Collembola.
W Ireland
- H. giardi*** (Berlese & Trouessart,1889)
= *Seius giardi* Berlese & Trouessart,1889
W. Ireland, ?French Atlantic coast
- Eviphidoidea**
- Eviphididae**
- Thinoseius*** Halbert,1920
* In decaying seaweed.
- T. ramsayi*** Evans,19??
Europe
- T. fucicola*** (Halbert,1920)
=? *T. berlexii* Halbert,1920
W Ireland, SW England
- T. spinosus*** Willmann,19??
Europe
- Pachylaelapidae** Vitzthum,1931
- Pachylaelaps*** Berlese,1888
- P. littoralis*** Halbert,1915
W Ireland
- Onchodellus*** Berlese,1904
- O.*** sp.
Britain
- Machrochelidae** Vitzthum,1930
- Machrocheles*** Latreille, in Cuvier,1829
* *Machrocheles* is the only spelling in Neave
- M. marginatus*** var. *littoralis* (Halbert,1915)
= *Holostaspis marginatus* var. *littoralis* Halbert,1915
Ireland
- Ascoidea**
- Halolaelapidae**
- Gamasellus*** Berlese,1892 (TM *G. falciger* (G. & R. Canestrini,1885?))
* This genus is not found in Hayward & Ryland (Eds.) 1990. Synonymized? - Correct family?
- G. inermis*** Halbert,1920
W Ireland
- Gamasus*** Latreille,1802
* This genus is not found in Hayward & Ryland (Eds.) 1990. Synonymized? - Correct family?
- G. (Gamasus) kempersi*** Oudemans,1902
Norway, Netherlands, Northumberland, Ireland
- G. (Gamasus) lunaris*** Berlese,1906
= *G. rubescens* G. & R. Canestrini,1882
* Possibly not a real shore-mite
Northumberland, Ireland, Italy
- G. coleoptratorum*** (Linnaeus,1758)
= *Acarus coleoptratorum* Linnaeus,1758
Ireland
- G. crassipes*** (Linnaeus,1758) var. *longicornis* Berlese,1???
* Possibly not a real shore-mite
W Ireland
- Gamasoides*** Berlese,1903 (TM *Poecilochirus carabi* G. & R. Canestrini,1885)
- G. spinipes*** (C.L. Koch,1844)
= *G. brachiosus* (G. Canestrini,1885-89)
W Ireland
- Halolaelaps*** Berlese & Trouessart,1889
- H. marinus*** (Brady,1875)
= *Gamasus marinus* Brady,1875
=? *H. glabriusculus* Berlese & Trouessart,1889
= *Zercon marinus* : Moniez,1890
= *Parastinus marinus* : Oudemans,1902
W Ireland, Britain, France, Spain
- H. celticus*** Halbert,1915
W Ireland
- H. incisus*** Hyatt,1980?
Br. Isles
- Saprogamasellus*** Berlese & Trouessart,1???
- S.*** sp.
Britain
- Antennoseiidae**
- Antennoseius*** Berlese,1916
- A.*** sp.
Britain
- Ascidae**
- Episeius*** Hull,1918
* This genus is not found in Hayward & Ryland (Eds.) 1990. Synonymized? - Correct family?
- E. grandis*** (Berlese,1916)
= *Lasioseius grandis* Berlese,1916
W Ireland
- Arctoseioides*** Willmann,1949
- A.*** sp.
- Britain
- Arctoseius*** Thor,1930
- A.*** sp.
Britain
- Iphidozercon*** Berlese,1904
- I.*** sp.
Britain
- Leioseius*** Berlese,1917
- L.*** sp.
Britain
- Neojordensia*** Evans,1957
- N.*** sp.
Britain
- Platyseius*** Berlese,1917
- P.*** sp.
Britain
- Phytoseiidae**
- Amblyseius*** Berlese,1915
- A.*** sp.
Britain
- Lasioseius*** Berlese,1917 (TM *Seius muricatus* (C.L. Koch,18??))
- L. salinus*** Halbert,1920
W Ireland
- Seius*** Agassiz,1845
= *Sejus* Koch,1836
- Tychlodromus*** Scheuten,1???
- T.*** sp.
Britain
- Veigaioidae**
- Veigaiidae*** Oudemans,1939
- Cyrthydrolaelaps*** Berlese,1905
- C. hirtus*** Berlese,1905
* In intertidal crevices
Wales, W Ireland, NW France, Spain
- Gamasolaelaps*** Berlese,1904
- G. excisus*** (L. Koch,1879)
= *Seius excisus* L. Koch,1879
= *Cyrtolaelaps (?) aurantiacus* Berlese,1903
= *Metaparastus suboles* Oudemans & Voigts,1904
Bremen, W Ireland, Siberia
- IXODIDA**
- Ixodoidea**
- Ixodidae***
= *Isodidae*
- Ixodes*** Latreille,1795
- I. uriae*** (White,1???)
@ *Uria aalge*
British Columbia
- NOTOSTIGMATIDA**
* Not in Scandinavian maritime or marine zone?
- HOLOTHYRIDAE**
* Not in Scandinavian maritime or marine zone?
- PANTOPODA*** Gerstaecker,1863
= ***PYCGONOIDA*** Latreille,1810
= ***PODOSOMATA*** Leach,1815
- PEGMATA*** Fry,1978
- Nymphonidae** Wilson,1878
- Nymphon*** J.C. Fabricius,1794
- N. brevirostre*** Hodge,1863
= *N. rubrum* Hodge,1865
= *N. gracile* : G.O. Sars
Öresund,Isefjord, Bohuslän, Denmark, all Norway, British Isles, Bretagne
- N. brevitarse*** Kroyer,1844
Greenland, Strait of Matotschkin Sharr
- N. hirtum*** J.C. Fabricius,1794
= *Chaetonyphon hirtum* : Auctt.
Skagerak, Troms, Belgium, Northumberland, Scotland, Shetlands, Iceland, E & W Greenland, Spitzbergen
- N. glaciale*** Lilljeborg,1870
Varangerfjord
- N. gracile*** Leach,1814
non = *N. gracile* : G.O. Sars
Bergen - Nordland, Morocco
- N. grossipes*** (O. Fabricius,1780)
Loftoten - Varangerfjord
- N. mixtum*** Kröyer,1844-45
= *N. grossipes* Auctt. non? (O. Fabricius,1780)
* *N. grossipes*, *N. mixtum* & *N. glacialis* Lilljeborg,1851 seem to be a species complex, of which only *N. mixtum* can be found S. of the Arctic.
Öresund - Bohuslän - Loftoten, Denmark, Shetland

- N. leptochelus** G.O. Sars,1888
Bohuslän-Skagerrak, all Norway, British Isles
- N. gracilipes** Heller,1875
= *N. giganteum* Goodsir,1844
* Is very close to, or possibly identical with *N. stroemi*.
Skagerrak (off Arendal) in 150-250 m - NE Finnmark, Northumberland
- N. stroemi** Krøyer,1844-45
Kattegatt - Bohuslän - W Finnmark
- N. macrum** Wilson,1880
W. Skagerrak, Bergen - NE Finnmark, W Greenland
- N. longitarse** Krøyer,1844-45
Bergen, Varangerfjord, Northumberland, Arctic
- N. longimanum** (G.O. Sars,1888)
W. Skagerrak??, Jan Mayen, Kara Sea
- N. spinosum** Goodsir,1842
= *Chaetonymphon s.*
Bohuslän-Skagerrak, all Norway
- N. spinosissimum** Norman,1894
= *Chaetonymphon s.*
Bergen - Varangerfjord, Spitsbergen, Iceland, S & W Greenland
- N. hirtipes** Bell,1853
= *Chaetonymphon h.*
Lofoten - NE Finnmark, NE of Shetland
- N. macronyx** G.O. Sars,1877
= *Chaetonymphon m.*
NE Finnmark, Faroe-Shetland Channel, Spitzbergen, E & W Greenland
- N. tenellum** (G.O. Sars,1888)
= *Chaetonymphon tenellum* G.O. Sars,1888
W. of Finnmark at 1134m to Faroe Channel
- N. serratum** G.O. Sars,1879
W. Norway, Hudson Bay
- N. megalops** G.O. Sars,1887
Bergen - S Troms, between Norway and Shetland
- N. elegans** H.J. Hansen,1887
N. Norway, Shetland-Faroe Channel
- N. sluteri** Hoek,1881
Varangerfjord - NE Finnmark, Spitzbergen, Arctic
- N. microrhynchum** G.O. Sars,1888
Novaja Zemlya, Arctic
- Boreonymphon** G.O. Sars,1888
- B. abyssorum** (Norman,1873)
= *Nymphon abyssorum* Norman,1873
Møre, Norwegian Sea
- B. ossiansarsi** Knaben, in Just,1972
Off W. Norway
- B. compactum** Just,1972
E & NW Greenland
- B. robustum** (Bell,1855)
= *Nymphon robustum* Bell,1855 (TM)
= *Nymphon hians* Heller,1875
between Faeroes & Iceland, off N Iceland, E & W Greenland, Spitsbergen, Kara Sea
- Acheliidae** Fry,1978
- Achelia** Hodge,1864
- A. echinata** Hodge,1864
@ *Flustra foliacea*
Bohuslän, Hirshals, Helgoland, Bergen - E Finnmark, British Isles
- A. longipes** (Hodge,1864)
= *Anmothea longipes* Hodge,1864
NW Finnmark - Porsangerfjord, British Isles, Atlantic coast of France, W Ireland
- A. hispida** Hodge,1864
* May be synonymous with *A. longipes*.
W Finnmark - Porsangerfjord, SW Britain, W Ireland
- A. laevis** Hodge,1864
Trondelag S of Trondheimsfjord, Helgoland, British Isles
- A. simplex** (Giltay,1934)
Isle of Man, Galway, Millport
- A. vulgaris** (Costa,1861)
Britain
- Ammotheidae** Dohrn,1881
= *Phoxichiliidae* G.O. Sars,1891 (p.p.)
- Ammothea** Leach,1814
- A. hilgendorfi** (Böhm,1879)
Britain (introduced)
- Eurycyde** Schiödte, in Rink,1857
- E. hispida** (Krøyer,1844-45)
Norway (Nordmøre and northwards), Denmark?
- Phoxichilidium** Milne-Edwards,1840
- P. femoratum** (Rathke,1799)
@ reported from cysts in *Syncoyne eximia*
Bohuslän, all Norway, Britain, Greenland, France
- P. tubulariae** Lebour,1947
@ gastric cavity of *Tubularia larynx*
Bohuslän, Holland, British Isles
- P. viresceus** Hodge,1864
- © reported from cysts in *Syncoyne eximia*
British Isles, France
- Paranymphon** Caullery,1896
* Correct family?
- P. spinosum** Caullery,1896
off Wales in >90m
- Tanystylum** Miers,1879
* Correct family?
- T. conirostre** (Dohrn,1881)
?Britain, Mediterranean
- Anoplodactylidae**
- Anoplodactylus** Wilson,1878
- A. angulatus** (Dohrn,1881)
SW Britain, France, Mediterranean
- A. petiolatus** (Krøyer,1844)
@ manubrium of *Obelia* sp. medusae
Öresund - Bohuslän - S Troms, North Sea, British Isles
- A. pygmæus** (Hodge,1864)
@ gastric cavity of *Obelia geniculata* polyps & on *Dynamena pumila*
Denmark, British Isles, Helgoland, Holland, France, Azores
- A. exiguis** (Dohrn,1881)
Bohuslän? (refound recently (Oct. 1994) in 30-38 m. at Lilleskärsslättan), Atlantic coast of France
- A. typhlops** G.O. Sars, 1891?
Norway (More - Nordland), Off W. Ireland
- Endeidae** Norman,1908
- Endeis** Philippi,1843
- E. spinosa** (Montagu,1808)
= *Phoxichilus s.*
= *E. laevis* (Grube,1871)
Bohuslän, Stavanger - Møre, British Isles
- E. charybdaea** (Dohrn,1881)
= *E. spinosa* sensu King, 1974, non (Montagu)
Stavanger area (Dusevigen), Roscoff, British Isles, Mediterranean
- Collosendeidae** Hoek,1881
- Colossendeis** Jarzynsky,1870
- C. proboscidea** (Sabine,1824)
Varangerfjord, Scotland-Faroe Channel
- C. angusta** G.O. Sars,1877
Between Norway and Shetland at 670-820m
- Callipallenidae** Hilton,1942
- Callipallene** Flynn,1929
- C. brevirostris** (Johnston,1837)
Kieler Bucht - Bohuslän - S Nordland, Hirshals, British Isles
- C. phantom** (Dohrn,1881)
= *Pallene producta* G.O. Sars,1888
Bergen - NW Sør-Trøndelag, British Isles
- C. spectrum** (Dohrn,1881)
Britain-Mediterranean
- C. tiberi** (Dohrn,1881)
Britain-Mediterranean
- C. emaciata** (Dohrn,1881)
Atlantic coast of France, England, Wales
- Cordylochele** G.O. Sars,1888
- C. longicollis** G.O. Sars,1888
Skagerrak at 343m (off Arendal) - Lofoten
- C. malleolata** (G.O. Sars,1879)
Off N. Norway at 168-200m
- C. brevicollis** G.O. Sars,1888
Vadsø, Faeroes
- Pseudopallene** Wilson,1878
- P. circularis** (Goodsir,1842)
= *Pallene intermedia* Krøyer,1844-45
= *Pallene discoidea* Krøyer,1844-45
Kullen - Skagerrak - all Norway, Scotland
- P. spinipes** (O. Fabricius,1780)
Bergen - NE Finnmark
- PYCNOGONOMORPHIDA** Fry,1978
- Pycnogonidae** Wilson,1878
- Pycnogonum** Brünlich,1764
- P. littorale** (Ström,1762)
@ *Actinia equina*, *Metridium senile* & *Urticina felina*; may also be associated with *Cucumaria frondosa*
Öresund - Bohuslän - all Norway, British Isles
- P. crassirostre** G.O. Sars,1888
@ Sea anemones
North Sea, Hardangerfjord - Trondheimsfjord - S Troms
- ASCORHYNCHOMORPHIDA** Fry,1978
* Not represented in the area.
- RHOPALORHYNCHOMORPHIDA** Fry,1978
* Not represented in the area.
- RHYNCHOTHORACIMORPIDA** Fry,1978

* Not represented in the area.

UNIRAMIA Snodgrass,1935

* Most taxa of this group treated in this list, are not truly marine. Some Diptera larvae are obligate or facultative estuarine or marine inhabitants. The rest of the species represented in the check-list may be referred to as thalassobionts (bound to the marine littoral fringe) or thalassophilic (facultative inhabitants, of the sea shore, preferring this environment to others). Many other organisms may be characterized as either indifferent sea shore taxa (may as well be found in other environments) or as temporal taxa in the marine littoral. Organisms of indifferent and temporal occurrence in the marine supralittoral zone have, in most cases, been completely left out from this list. Some insect orders, e.g. Hymenoptera, to which some sea shore dwellers belong (e.g. the sand dune species *Pomphilus* Fabricius,1798 *plumbeus* (Fabricius,1787)), have been omitted in this list, on reasons of difficulties in bordering between maritime and non-maritime species. Thus, this tableau must be regarded as very incomplete as a check-list. Certainly, several taxa are missing, and some of the encountered species may possibly show to be temporal visitors to the marine littoral, and should thus perhaps be weeded out.

MYRIPODA "Tusenfotinger"

CHILOPODA "Enkelfotinger"

* Only thalassobionts and a few thalassophilic taxa included

ANAMORPHA

LITHOBIIDA "Stenkrypare"

LITHOBIIDEA

Lithobioidae

Lithobius Leach,1814

L. melanops G. Newport,1845

* Thalassophilic.
Scandinavia, Germany, Iceland

L. forficatus (Linnaeus,1758)

* Thalassophilic.
Scandinavia (e.g. Bohuslän-Uppland), Iceland, France

L. calcaratus ?Koch,1???

* In wrack.
Småland archipelago

L. erythrocephalus ?Koch,1???

* In wrack.
Blekinge-Uppland, Åland

L. curtipes ?Koch,1???

* In wrack.
Uppland, Åland, Finland

Henicopidae

Lamycetes Meinert,1868

* Does this genus belong in that family?

L. fulvicornis Meinert,1868?

* Thalassophilic.

Iceland

EPIMORHA

GEOPHILIDA "Jordkrypare"

Schendylidae

Schendyla Bergsöe & Meinert,1866

S. nemorensis (C.L. Koch,1836)

* Thalassophilic. In Wrack.
Bohuslän

Hydroschendyla Brölemann & Ribaut,1911

H. submarina (Grube,1870)
Sweden, Bretagne, S. England, Mediterranean

Brachyschendyla Brölemann & Ribaut,1911

B. monodi Brölemann,1924
Finistère, Arcachon

Geophilidae

Strigamia Gray,1843

= *Scolioplanes* Bergsöe & Meinert,1866

S. maritima (Leach,1817)

Gotland, Vinga, Bohuslän, S. Norway, Denmark, Helgoland, Br. Isles, France

Geophilus Leach,1814

G. fucorum ?1??? *seurati* Brölemann,1930?

Plymouth, Scotland, Ireland

G. proximus Koch,1???

* In wrack.
Uppland, Åland, Finland

G. algarum ?1??? *decipiens* Brölemann,19??

N France, Arcachon

Necrophloeophagus Newport,1843

N. longicornis (Leach,1815)

= *Geophilus longicornis* Leach,1815

* Thalassophilic.

Scandinavia, Britain

Chilenophiliidae

Pachymerinae

Pachymerium C.L. Koch,1847

P. ferrugineum (C.L. Koch,1835)

* Thalassophilic.

Blekinge, Uppland, Denmark, Åland, Åbo, Germany, Arcachon

SYMPHYLA "Dvärgfotinger"

Scolopendrellidae

Scolopendrella Gervais,1839

S. notacantha Gervais,1839

* Thalassophilic.

Wales

PAUROPODA "Fåfotinger"

TETRAMEROCERATA

ECTOMORPHA

Pauropodidae

Thalassopauropus Bagnall,1935

T. remyi Bagnall,1935

E. Scotland, French Mediterranean

Allopauropus Berlese,1902, ex Silvestri MS

A. thalassophilus Remy,19??

E. Scotland

A. danicus (Hansen,1???)

E. Scotland

A. littoralis Bagnall,1935?

E. Scotland

A. stepheni Bagnall,1935?

E. Scotland

DIPLOPODA "Dubbelfotinger"

* Only thalassobionts and a few thalassophilic taxa included

PENICILLATA

= **PSEPHALOGNATHA** Latzel,1884 "Penselfotinger"

POLYXENIDA

= **SCHIZOCEPHALA** Verhoeff,1926

Polyxenoidea

Polyxenidae Gray & Jones,1842

Polyxenus Latreille,1802

P. lagurus (Linnaeus,1758)

* Thalassophilic.
Bohuslän, Skåne, Åland, Germany, Denmark, Norway, Br. Isles, etc.

PENTAZONIA

* No maritime species

HELMINTHOMORPHA

DIPLOCHETA

JULIDA

Nemasomatidae

Thalassoisobates Verhoeff,1908

= *Isobates* Menge,1861 (p.p.)

T. littoralis (Silvestri,1903)

* In wrack on rocky shores.

Bohuslän, Br. Isles, Mediterranean

Blaniulidae Attems,1909

Choneiulus Brölemann,1921

C. palmatus (Nemec,1895)

* Occasionally in wrack.
Bohuslän, S Norway, Denmark, Finnland, Germany, Netherlands, Britain

Proteroziulus Silvestri,1897

P. fuscus (Am Stein,1857)

* Occasionally in wrack, normally a forest species.
S Sweden, Norway, Åland, S Finnland, Germany, Netherlands, Belgium, France, Britain

Julidae Leach,1815

Cylindroiulidae Verhoeff,1930

Cylindroiulus Verhoeff,1894

C. (Aneuloboiulus) latestriatus (Curtis,1844)

= *C. (Aneuloboiulus) frisius* (Verhoeff,1891)

* Thalassophilic. In wrack.

Gotland, Bohuslän, Norway, Denmark, Germany, Faroes, Netherlands, Br. Isles, Ireland, Arcachon

C. (Aneuloboiulus) punctatus (Leach,1815)

= *C. (Aneuloboiulus) silvarium* (Meinert,1868)

* Occasionally in wrack.

Småland archipelago, Uppland, W Sweden, Norway, S Finnland, Denmark, Netherlands, Belgium, France

Julinae Verhoeff,1930

Julus Linnaeus,1758

J. terrestris Porat,1889

* Thalassophilic

Finland, Åland, Gotland, E. & S. Sweden, Bornholm, Sealand, Bohuslän, Germany

Schizophyllidae Verhoeff,1910

Ommatoiulus Latzel,1884

= *Schizophyllum* Verhoeff,1895 (p.p.)

= *Archiulus* Berlese,1886, non Scudder,1873 (p.p.)

- O. sabulosus** (Linnaeus,1758)
 = *Schizophyllum* Verhoeff,1895 (*Bothrioulus* Verhoeff,1894) *sabulosum* (Linnaeus,1758)
 * Thalassophilic
 S. Finland, Denmark, Germany, Sweden, S Norway, Br. Isles etc.
- Tachypodoiulus** Verhoeff,1894
- T. niger** (Leach,1815)
 = *Julus albipes* C.L. Koch,1838
 * Thalassophilic.
 Normandie, Netherlands, Belgium, Germany, Br. Isles, Ireland
- Brachyiulina** Verhoeff,1910
- Brachyiulus** Berlese,1884
- B. pusillus** (Leach,1814)
 = *B. littoralis* Verhoeff,1898
 Öland, Gotland, Skåne-Blekinge-Småland, Estonia-Germany, Britain, Ireland, France, Spain
- HEXAPODA** Latreille,1825
 * Probably a polyphyletic taxon. Some entognaths may be closer to Myriapoda.
- ENTOGNATHA**
- DIPLURATA** "Larvborstsvansar"
 * Not represented in the area with any sea shore species.
- MYRIENTOMATA**
 = **PROTURA** "Trevfotinger"
 * Not represented in the area with any sea shore species.
- OLIGOENTOMATA**
- COLLEMBOLA** Lubbock,1870 "Hoppstjärtar"
 * Chiefly thalassobionts and a few thalassophilic species have been listed here.
- PODUROMORPHA** Börner,1913
- Hypogastruridae** Börner,1906
- Hypogastrura** Bourlet,1839
- H. viatica** (Tullberg,1872)
 * Thalassophilic. Considered to be cosmopolitan in wrack and salt marshes.
 Bohuslän-all Baltic coasts, Denmark, Norway, German North Sea coast, Br. Isles, France
- H. vernalis** (Carl,1901)
 * Littoral & in wrack.
 Sörmaland-Uppland, Åland
- H. gisini** (Strenzke,1954)
 along sandy sites in Baltic Shores & Sylt
- Ceratophysella** Börner, in Brohmer,1942
- C. armata** (Nicole,1842)
 = *Hypogastrura armata* : Auctt.
 * Ubiquistic & in wrack.
 Bohuslän-Uppland, Finland
- Xenylla** Tullberg,1871
- X. humicola** (O. Fabricius,1780)
 * Thalassophilic. Considered to be presumably cosmopolitan in wrack and salt marshes.
 Bohuslän-all Baltic coasts, Norway, S. North Sea coasts, Br. Isles, France, Mediterranean
- X. maritima** Tullberg,1871
 * Thalassophilic. Considered to be presumably cosmopolitan in wrack and salt marshes.
 Finland, W. Sweden, Uppland, Norway, German North Sea coast, Br. Isles, France, Mediterranean
- Neanuridae** Börner,1901
- Neanurinae** Börner,1901
- Neanura** Macgillivray,1893
 = *Anoura* Gervais,1843, *non* Gray,1838, (Mammalia) *nec* Gray,1840 (Mollusca)
- Pseudachorutinae** Börner,1906
- Anurida** Laboulbène,1865
- A. maritima** (Guérin-Méneville,1836) Laboulbène,1865
 Kattegatt-Bohuslän, Norway, Germany, Netherlands, Belgium, Br. Isles, Arcachon
- A. tullbergi** Schött,1891
 * thalassophilic
 W. Sweden, Sörmaland, Finland, German Baltic & North Sea coasts, Norway, Britain, France, Mediterranean
- A. granaria** (Nicole,1847) Tullberg,1869
 * In wrack & humus
 Skåne-Småland, Finland
- A. thalassophila** (Bagnall,1939) Stach,1949
 = *Anuridella thalassophila* : Auctt.
 Br. Isles
- Pseudachorutella** Stach,1949
- P. remyi** (Denis,1933) Massoud,1967
 = *Anurida remyi* : Auctt.
 Finland, Skåne, German Baltic & North Sea coasts, Jan Mayen, Greenland
- Gastranurida** Bagnall,1949
- G. denisi** (Bagnall,1939) Massoud,1967
 = *Anurida denisi* : Auctt.
 Br. Isles, France
- Anuridella** Willem,1906
- A. marina** Willem,1906
 Br. Isles, France
- A. calcarea** Denis,1925
 Br. Isles, Mediterranean
- A. germanica** (Gisin,1951)
 German Baltic & North Sea coast
- A. submarina** Bagnall,1934
 * A synonym of which other species?
 Br. Isles
- A. immsiana** Bagnall,1939
 * A synonym of which other species?
 Br. Isles
- Frieseinae** Massoud,1967
- Friesea** Dalla Torre,1895
- F. acuminata** (Denis,1925) Goto & Lawrence, in Kloet & Hincks,1964
 Br. Isles, Mediterranean
- F. mirabilis** (Tullberg,1871) Willem,1901
 * In deep wrack layers & humus
 Västra-Åland, Åland, Finland
- Brachystomellidae** Stach,1949
- Brachystomella** Ågren,1903
- Onychiuridae** Börner,1901
- Onychiurus** Gervais,1841
- Protaphorura** Absolon,1901
- P. debilis** (Moniez,1889) Salmon,1964
 = *Onychiurus debilis* : Auctt.
 German Baltic & North Sea coast, Br. Isles, Arcachon, Mediterranean
- P. thalassophila** (Bagnall,1937) Hopkin,1997
 = *Onychiurus thalassophila* : Auctt.
 Scotland, England
- P. halophila** (Bagnall,1937) Salmon,1959
 = *Onychiurus halophila* : Auctt.
 Northumberland, Yorkshire
- P. armata** (Tullberg,1869) Börner,1909
 = *Onychiurus armata* : Auctt.
 * In deep wrack layers.
 Bohuslän-Uppland, Finland
- P. tullbergi** (Bagnall,1935) Kloet & Hincks,1945
 = *Onychiurus tullbergi* : Auctt.
 Scotland
- Tullbergiidae** Bagnall,1935
- Tullbergia** Lubbock,1876
- Odontilidae** Massoud,1967
- ENTOMOBRYOMORPHA** Börner,1913, sensu Mateos,1988
- Isopomoidea** Schäffer,1896 (Börner,1913 (Szeptycki,1979))
- Isotomidae** Schäffer,1896 (Börner,1913)
- Isotominae** Schäffer,1896
- Bagnallella** Salmon,1951
- Isotoma** Bourlet,1839, *non* Dejean,1834 (Coleoptera)
- Desoria** Nicolet, in Desor,1841
- D. tigrina** Nicolet,1842
 * In deep wrack layers.
 Bohuslän-Uppland, Åland, All Europe, Arctic, Antarctic, Australia
- D. olivacea** (Tullberg,1871)
 * In deep wrack layers & eurytopic
 Bohuslän-Uppland, Åland, All Europe, Arctic
- Halisotoma** Bagnall,1949
- H. maritima** (Tullberg,1871) Bagnall,1949
 = *Isotoma maritima* Tullberg,1871
 * In wrack and under stones.
 W. Sweden, Skåne, all Baltic coasts, Norway, S. North Sea coasts, Br. Isles, France, Mediterranean
- H. boneti** (Delamare-Debouteville,1954)
 Portugal-Senegal, S Spain, Mediterranean
- H. poseidonis** (Bagnall,1939)
 = *Isotoma poseidonis* Bagnall,1939
 Br. Isles, Ireland, Biscay
- Isotomiella** Bagnall,1939
- Isotomurus** Börner,1903
- Axelsonia** Börner,1906
- A. littoralis** (Moniez,1890) Handschin,1929
 * In wrack and littoral subsoil.
 North Sea shores, Br. Isles, French North Sea coast, Arcachon, Mediterranean, S America, Australia
- Proisotominae**
- Folsomia** Willem,1902
- F. sexoculata** (Tullberg,1871) Handschin,1929
 Skåne-Uppland, Åland, Norway, German Baltic & North Sea coast, Finland, Denmark, Br. Isles
- F. thalassophila** Bagnall,1940
 Scotland, Ireland
- F. garetti** Bagnall,19??
 * Possibly not a sea shore species? Not recognized in Janssens' check-list.
 Br. Isles
- Proisotoma** Börner,1901
- P. buddenbrocki** Strenzke,1954
 Tärminne, Kieler Bucht

<i>P. admaritima</i> Murphy,1953	Katianidae Börner,1913, sensu Stach,1956
Scotland	
<i>Ballistura</i> Börner,1906	<u>Sturmidae</u> Bretfeld,1994
<i>B. schoetti</i> (von Dalla Torre,1895) Stach,1947	<u>Arrhopalitidae</u> Stach,1956
* Thallasophilic. In wrack and under stones.	<u>Spinophecidae</u> Delamare Deboutteville,1961
W. Sweden, Sörmland-Uppland, Finland, German North Sea coast, Br. Isles	<u>Dicyrtomidae</u> Börner,1906
<i>B. crassicauda</i> Tullberg,1871	<u>NEELIPLLEONA</u> Massoud,1971
* Thallasophilic. In wrack and under stones. Not recognized in Janssens' check-list.	<u>Neelidae</u> Folsom,1896
Finland, German Baltic coast, W. Sweden, Br. Isles	
<u>Auroporphinae</u>	
<i>Archisotoma</i> Axelson,1912	<u>INSECTA</u>
* in littoral subsoil and salt marshes	= <u>ECTOGNATHA</u>
<i>A. besseli</i> (Packard,1877) Willem,1925	<u>ARCHAEOGNATHA</u>
Sweden, ?Norway, German Baltic & North Sea coast, Finland, ?Br. Isles, France, Mediterranean	
<i>A. pulchella</i> (Moniez,1890) da Gama,1968	<u>APTERYGOTA</u>
Denmark, German North Sea coast, Arcachon, ?Br. Isles	= <u>ZYGENTOMA</u>
<i>A. megalops</i> Bagnall,1939	<u>THYSANURA</u> Latreille,1804 "Fjällborstsvansar"
Br. Isles	<u>ZYGOENTOMATA</u>
<i>A. nigricans</i> Bagnall,1939	* Not represented in the marine littoral in the area.
Br. Isles	<u>ARCHEOGNATHA</u>
<i>A. subbrucei</i> Delamare-Debouteville,1954	= <u>MICROCORYPHIA</u>
French Atlantic coast	<u>Machiloidea</u>
<i>Actaletidae</i> Börner,1902	<u>Machilidae</u> Verhoeff,1912 "Nattsmygär", "Silverfiskar"
<i>Actaletes</i> Giard,1889	<u>Machiliinae</u>
<i>A. neptuni</i> Giard,1889	<u>Petrobius</u> Leach,1817 "Kustnattsmygär"
On rocks in the intertidal zone of the French North Sea coast, ?Br. Isles	<u>P. brevistylis</u> Carpenter,1913
<i>Coenaletidiae</i> Bellinger,1985	= <i>P. maritimus</i> Auct., <u>non</u> (Leach,1815)
<i>Isotogastruridae</i> Thibaud & Najt,1992	Baltic-Bohuslän, Helgoland
<i>Entomobryoidea</i> Schött,1891 (Womersley,1934, sensu Szeptycki,1979)	<u>P. maritimus</u> (Leach,1809)
<i>Entomobryidae</i> Schött,1891	= <i>P. lohmanderi</i> Agrell,1944 (synonymized by Davies 1988)
<i>Entomobryinae</i> Schött,1891 (Schäffer,1896)	E. Sweden (on rocky shores), Denmark, Atlantic coasts (on rocky shores), e.g. Br. Isles
<i>Entomobrya</i> Rondani,1861	
= <i>Degerria</i> Nicolet,1841, <u>non</u> Meigen,1838 (Diptera)	<u>PTERYGOTA</u>
= <i>Mydonius</i> Gistl,1848	= <u>PTILOTA</u> "Yrfän"
<i>Seira</i> Lubbock,1870	* The first order of this infraclassis, the Ephemeroptera, has no really marine or maritime species in the area. However larvae of some species within Odonata may be found in estuaries. A list of such species found in the Baltic by Lindberg,1948 thus has been used for making this list.
<i>Willowsia</i> Shoebottom,1917	<u>EPHEMEROPTERA</u> "Dagsländor"
<i>Lepidocyrtinae</i>	<u>Ephemeroidae</u>
<i>Lepidocyrtus</i> Bourlet,1839	<u>Baetidae</u>
<i>L. cyaneus</i> (Tullberg,1871) Tullberg,1871	= <u>Heptagenioidea</u>
* In wrack surface layers.	<u>Baetidae</u>
Bohuslän-Uppland, Finland	<u>Callibaetidae</u> Eaton,1881
<i>L. lanuginosus</i> (Gmelin,1788) Tullberg,1871	<u>ODONATA</u> "Trollsländor"
* In wrack surface layers & ubiquitous.	<u>ZYGOPTERA</u>
Blekinge-Uppland, Åland, Finland	<u>Lestinoidea</u>
<i>Pseudosinella</i> Schäffer,1897	<u>Lestidae</u>
<i>P. halophila</i> Bagnall,1939	<i>Lestes</i> Leach,1815
Silt, England	<i>L. sponsa</i> (Hansemann,1823)
<i>P. maritima</i> Bagnall,1941	N. Baltic
Br. Isles, Ireland	<u>Coenagrionoidea</u>
<i>Orchesellinae</i>	<u>Coenagrionidae</u>
<i>Orchesella</i> Templeton,1835	<i>Ischnura</i> de Charpentier,1840
<i>O. litoralis</i> Brown,1925	<i>I. elegans</i> (P.L. v. d. Linden,1820)
Yorkshire	N. Baltic
<i>Microfalculidae</i> Massoud & Betsch,1966	<i>Enallagma</i> de Charpentier,1840
<i>Paronellidae</i> Börner,1913	<i>E. cyathigerum</i> (de Charpentier,1840)
<i>Cyphoderidae</i> Börner,1913	N. Baltic
<i>Oncopoduridae</i> Carl & Lebedinsky,1905	<i>Erythromma</i> de Charpentier,1840
<i>Tomoceroidea</i> Schäffer,1896 (Szeptycki,1979)	<i>E. najas</i> (Hansemann,1823)
<i>Tomoceridae</i> Schäffer,1896	N. Baltic
<i>METAXYPLEONA</i> Salmon,1964	<u>Agrionoidea</u>
<i>Poduridae</i> Latreille,1802-03	= <u>Calopterygoidea</u>
<i>SYMPHYPLEONA</i> Börner,1901	<u>Agrionidae</u>
<i>Sminthuridae</i> Lubbock,1862	= <u>Calopterygidae</u>
<i>Deuterostinturus</i> Börner,1901	<i>Agrion</i> Fabricius,1775
= <i>Deuterostinturus</i> Börner,1901	= <i>Calopteryx</i> de Charpentier, <u>in</u> Burmeister,1839
<i>Sminthurides</i> Börner,1900	<i>A. armatum</i> de Charpentier,1840
<i>Sminthirinus</i> Börner,1901	N. Baltic
<i>Smynthurus</i> Latreille,1802	<i>A. pulchellum</i> P.L. v. d. Linden,1825
<i>Bourletiellidae</i> Börner,1913	N. Baltic
<i>Bourletiella</i> Banks,1899	<i>A. hastulatum</i> de Charpentier,1825
<i>Mackenziellidae</i> Yosii,1961	N. Baltic
<i>Sminthurididae</i> Börner,1906	<u>ANISOPTERA</u>

AeshnoideaAeshnidae

Aeshna Fabricius,1775
= *Aeschna* Illiger,1802

A. coerulea (H. Ström,1783)
N. Baltic

A. juncea (Linnaeus,1758)
N. Baltic

A. grandis (Linnaeus,1758)
N. Baltic

A. cyanea (O.F. Müller,1764)
N. Baltic

Brachytron Evans,1845

B. hafniense (O.F. Müller,1764)
N. Baltic

LibelluloideaLibellulidae

Libellula Linnaeus,1758

L. quadrimaculata Linnaeus,1758
N. Baltic

Sympetrum Newman,1833

S. striolatum (de Charpentier,1840)
N. Baltic

S. vulgatum (Linnaeus,1758)
N. Baltic

S. flaveolum (Linnaeus,1758)
N. Baltic

S. danae (J.H. Sulzer,1776)
N. Baltic

Leucorrhinia Brittinger,1850

L. rubicunda (Linnaeus,1758)
N. Baltic

Orthetrum Newman,1833

O. cancellatum (Linnaeus,1758)
N. Baltic, SW Sjælland in 10 ppt S

NEOPTERAORTHOPTEROIDEA

* As far as known, by the compiler there are no truly maritime species.

ORTHOPTERA

= *SALTATORIA* "Hopprätvingar"

CAELIFERA

Acrididae "Markgräshoppor"

Omocestus Bolívar,1878

O. haemorrhoidalis (?17??)
Germany

ENSIFERA
= GRYLOOPTERA

Tettigoniidae "Vårtbitare"

Decticus Audinet-Serville,1831

D. verrucivorus (Linnaeus,1758) "Stor vårtbitare"
= *Gryllus verrucivorus* Linnaeus,1758
Sweden, Germany

DERMAPTERAFORFICULINALabiduroideaCarcinophoridaeLabiduridaeHEMIPTEROIDEAPHTIRAPTERA "Löss"

ANOPLURA "Åkta löss"

Echinophthiriidae

Echinophthirius Giebel,1871

E. horridus (von Olfers,1816) "Sällus"

= *E. sericans* Meinteger,18??

= *E. setosus* ??17??

= *E. phocae* Lucas,17??

@ earless seals, e.g. *Phoca vitulina*
Europe; e.g. around Sweden, Denmark, Netherlands, Arctic Seas

Antarctophthirus Enderlein,1906

A. trichechi (Bohemann,1???)
@ *Odobenus rosmarus*
Arctic Seas

A. microchir (Trouessart & Newman,1888)
@ *Otaridae* & *Phoca vitulina*
California

A. callorhini (Osborn,1899)

@ *Callorhinus ursinus*

Alaska

Proechinophthirus Ewing,1923

P. fluctus (Ferris,1916)

@ *Callorhinus ursinus* & *Eumetopias jubatus*

Arctic Seas, Alaska

MALLOPHAGA "Fjäderätere", "Fägellöss" (& "Pälssätare")

@ birds

* No check-list, but names of a few genera that may be found on sea birds, have been listed here.

This is a group, which is considered very difficult, even by specialists.

AMBLYCERAMenoponidae

Eidmanniella Kéler,1938

@ e.g. *Sula*, *Phalacrocorax*

Sweden, etc.

Actornitophilus Ferris,1916

@ e.g. *Larus*, *Rissa*

Sweden, etc.

Colpocephalum Nitzsch,1818

@ e.g. *Larus*, *Rissa*

Sweden, etc.

Comatomenopon Uchida,1920

@ Laridae

Sweden, etc.

Ancistrona Westwood,1874

@ Procellariiformes

Sweden, etc.

Procellariiphaga Eichler,19???

@ Procellariiformes

Sweden, etc.

Piagetiella Neumann,1906

@ e.g. *Phalacrocorax*

?

Austromenopon Bedford,1939

@ e.g. *Larus*, *Rissa*, *Alca*, *Fulmarus*

Sweden, etc.

A. crocatum Nitsch,18??, in Giebel

@ *Haematopus*

Br. Isles

Austromenopea ??,19???

* Is this *Austromenopon* wrongly spelled?

A. uriae Timmermann,1???

@ *Uria aalge*

N Atlantic, British Columbia

ISCHNOCERAPhilopteridae

Anaticola Clay,1935

Anatoecus Cummings,1916

Cummingsiella Ewing,1930

= *Dollabella* Cummings,1916, non de Lamarck,1801 (Mollusca)

Quadraceps Clay & Meinertzhagen,1939

@ e.g. *Larus*, *Rissa*, *Alca*, *Fulmarus*

Sweden, etc.

Q. obliquus (Mjöberg,1???)

@ *Uria aalge*

North Atlantic, British Columbia

Saemundssonia Timmermann,1936

@ e.g. *Larus*, *Rissa*, *Alca*, *Fulmarus*

Sweden, etc.

S. calva (Kellogg,1???)

@ *Uria aalge*

North Atlantic, British Columbia

Perineus Thompson,1936

@ Procellariiformes, Laridae

Sweden, etc.

Halipeurus Thompson,1936

@ Procellariiformes.

Sweden, etc.

Pectinopygus Mjöberg,1910

@ e.g. *Sula*, *Phalacrocorax*

Sweden, etc.

P. bassani (O.F. Müller,1776)

@ *Sula bassana*

Br. Isles

Philopterus Nitzsch,1818

P. gonothonax Giebel,18??

@ *Larus marinus*

Br. Isles

HEMIPTERA= HETEROPTERA "Halvvingar"

* The "most marine" species of this order is to be found in the suborder Gerromorpha, where four families, Gerridae, Veliidae, Mesovelidae and Hermatobatidae together have a dozen genera, e.g. *Halobates*, with both estuarine and oceanic species. However none of these more than 80 species is to be found in Europe. They occur in warmer seas.

GERROMORPHAGerroidea

- Gerridae** "Vattenlöpare"
- Gerris** J.C. Fabricius,1794
- G. odontogaster** (Zetterstedt,1828)
* A typical brackish water species. Limnohaline.
Baltic, Sweden, Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, France, Br. Isles
- G. thoracicus** Schummel,1832, ?non Stephens,1829
* A typical brackish water species.Pseudohalobiont. Halophilic.
Baltic, Sweden, Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, France, Br. Isles
- G. rufoscutellatus** Latreille,1807
Sweden, Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, France, ?Br. Isles
- G. argentatus** Schummel,1832
Sweden, Norway, Finland, Denmark, Germany, North Sea Shores, France, ?Br. Isles
- G. lacustris** (Linnaeus,1758)
Sweden, Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, France, ?Br. Isles
- Veliidae** "Bäcklöpare"
- Microvelia** Westwood,1834
- M. reticulata** (Burmeister,18??)
* A typical brackish water species.
Baltic
- Mesoveliidae** Douglas & Scott,1867
- Mesoveliinae** China & Miller,1959
- Mesovelia** Mulsant & Rey,1852
- M. furcata** Mulsant,1???
* A typical brackish water species. Limnohaline.
Baltic, Sweden, Norway, Finland, Denmark, Germany, North Sea Shores, France, Br. Isles
- NEPOMORPHA**
- Nepoidea**
- Nepidae** "Vattenskorpcioner"
- Nepa** Linnaeus,1758
- N. rubra** Linnaeus,1758
* A typical brackish water species.
Baltic
- N. cinerea** Linnaeus,1758
Baltic
- Notonectoidea**
- Notonectidae** "Ryggssimmare"
- Notonecta** Linnaeus,1758
- N. viridis** Delcourt.,18??
* Halophilic.
Germany, Netherlands, England
- N. glauca** Linnaeus,1758
Sweden, Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, France, Br. Isles
- Corixoidea**
- Corixidae** "Buksimmare"
- Cymatinae**
- Cymatia** Flor,1860
- C. bonsdorffi** (Sahlberg,1819)
Baltic
- C. coleoptrata** (J.C. Fabricius,1777)
Baltic, Sweden, Finland, Denmark, Balticum, Germany, North Sea Shores, France, Br. Isles
- C. rogenhoferi** Fieber,18??
Europe
- Micronectinae**
- Micronecta** Kirkaldy,1897
- M. meridionalis** (A. Costa,1847)
Europe
- M. minutissima** (Linnaeus,1758)
Sweden, Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, France, Br. Isles
- Corixinae**
- Arctocoris** Wallengren,1894
- A. carinata** (Sahlberg,1819)
Baltic
- Callicorixa** B. White,1873
- C. praeusta** (Fieber,1848)
Baltic, Sweden, Norway, Denmark, Finland, Balticum, Germany, North Sea Shores, France, Br. Isles
- C. producta** (Reuter,1???)
Baltic, Sweden, Finland
- Corixa** Geoffroy,1762
- C. affinis** Leach,1817
German Baltic coast, England
- C. panzeri** (Fieber,1848)
Sweden, Denmark, Germany, North Sea Shores, France, Br. Isles
- Hesperocorixa** Kirkaldy,1908
* Should spelling be *Hesperocrixa* ?
- H. sahlbergi** (Fieber,1848)
Sweden, Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, France, ?Br. Isles
- Sigara** J.C. Fabricius,1775
- S. assimilis** (Fieber,1848)
Europe
- S. concinna** (Fieber,1848)
Europe
- S. dorsalis** (Leach,1817)
England
- S. falleni** (Fieber,1848)
England, German Baltic coast
- S. lateralis lateralis** (Leach,1817)
Europe; e.g. Baltic
- S. selecta** (Fieber,1848)
Sweden, Norway, Denmark, Germany, North Sea Shores, France, Br. Isles
- S. stagnalis** (Leach,1817)
= *S. lugubris* Fieber,18??
* A typical brackish water species. Halophilic-Halobiont.
Baltic, Netherlands, Br. Isles
- S. distincta** (Fieber,1848)
* Halophilic.
Sweden, Denmark, Germany, Netherlands, Belgium, France, Br. Isles
- S. striata** (Linnaeus,1758)
* A typical brackish water species. Limnohalin.
Baltic, Sweden, Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, France, Br. Isles
- S. venusta** Douglas & Scott,1???
France
- LEPTOPODOMORPHA**
- Leptopodoidea**
- Salidae** "Strandstinkflyn"
- Saldinae**
- Salda** Fabricius,1803
- S. littoralis** (Linnaeus,1758)
* Halophilic.
Sweden, Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, France, Br. Isles
- Chartoscirta** Stål,1868
- C. cincta** (Herrick-Schaeffer,1841)
Europe; e.g. Sweden
- C. elegantula** (Fallén,1807)
Sweden
- Halosalda** Reuter,1912
- H. lateralis** Reuter,1912
* Halobiont.
Sweden, Norway, Finland, Denmark, Germany, North Sea Shores, France, Br. Isles
- Saldula** Van Duzee,1914
- S. pilosella** (Thomson,18???)
* Halobiont.
Sweden, Norway, Finland, Balticum, Br. Isles
- S. scotica** (Curtis,18???)
* Halophilic.
Sweden, Norway, Finland, Balticum, Br. Isles
- S. saltatoria** (Linnaeus,1758)
Sweden, Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, France, ?Br. Isles
- S. pallipes** (J.C. Fabricius,1781)
Sweden, Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, France, Br. Isles
- Micracanthia** Reuter,1912
* Right subfamily?
- M. fennica** Reuter,1912?
* Halobiont.
Sweden, Finland, Balticum, ?Norway
- Chiloxyanthinae**
- Chiloxyanthus** Reuter,1891
- C. pilosus** (Fallén,1807)
* Halobiont.
Sweden, Denmark, Germany, Netherlands, France, England
- Pentacora** Reuter,1912
- Aepophilinae** "Havstinkflyn"
- Aepophilus** Signoret,1879
- A. bonnairei** Signoret,1879, in Puton
* Halobiont.
Br. Isles, N. France, Ireland, Iceland
- CIMICOMORPHA**
- Miroidea**
- Miridae**
= *Capsidae* Burmeister "Ängsstinkflyn"
- Mirinae** Dohrn,1858
- Mirini** Dohrn,1858
- Poeciloscytus** Fieber,1858
- P. vulneratus** (Panzer,1798)

- = *Polymerus* Hahn,1831 (*Poeciloscytus*) *vulneratus* (Panzer,1798)
 @ e.g. *Salsola*, *Galium*, *Corispermum*, etc. * Halophilic.
 Sweden, Denmark, Germany, Netherlands, France, England
- Stenodemini* China,1943
Teratocoris Fieber,1858
T. saundersi Douglas & Scott,1867
 @ *Phragmites* * Halophilic.
 Sweden, Norway, Finland, Denmark, Balticum, Germany, Netherlands, Br. Isles
- Trigonotylus* Fieber,1858
T. psammaeicolor Reuter,1885
 @ *Ammophila* & *Elymus* * Halophilic.
 Sweden, Denmark, Germany, Netherlands, England
- Phylinae* Handlirsch,1925
Phylini Douglas & Scott,1845
Conostethus Fieber,1858
C. salinus (J. Sahlgren,1870)
 @ Halophytes
 Sweden, Denmark, Netherlands, England, German Baltic coast
- C. frisicus* E. Wagner,1952
 @ *Limonium*
 German North Sea coast
- C. brevis* Reuter,1877
 @ *Limonium*
 Scotland
- PENTATOMOMORPHA
- Piesmatoidea
Piesma Peletier & Serville,1825
P. quadrata (Fieber,1844?)
 * Halophilic.
 Sweden, Norway, Finland, Denmark, Germany, North Sea Shores, England
- P. salsolae* Becker,18??
 * Halophilic.
 Finland, France
- Lygaeoidea
Lygaeidae
Henestaris Spinola,1837
H. halophilus (Burmeister,1835)
 * Halobiont.
 Germany, France, England
- H. laticeps* (Curtis,1836)
 * Halobiont.
 France, England
- Trapezonotus* Fieber,1860
T. distinguendus Flor,18??
 * Halophilic.
 Sweden, Norway, Finland, Denmark, Balticum, Germany, Netherlands, England
- Bosus* Amyet & Serville,1843
B. maritimus (Scopoli,1763)
 * Halophilic.
 Germany, Netherlands, France, England
- HOLOMETABOLA
- NEUROPTERA "Nätvingar"
- MEGALOPTERA
- Sialidae "Sävsländor"
Sialis Latreille,1802
S. sp.
 Baltic
- TRICHOPTERA "Nattsländor"
 * Some species represented by larvae only in very "diluted" estuarine water, have not been listed here.
- Hydropsychoidea
Hydropsychidae
Hydropsyche Pictet,1834
H. sp.
 Baltic
- Polycentropodidae
Polycentropus Curtis,1835
P. flavomaculatus Pictet,18??
 Baltic
- Limnephiloidea
Limnephilidae
Limnephilus Leach,1815
L. affinis (Curtis,1827?)
 * The only true European estuarine-marine species. Tolerates a salinity of up to 24 ppt.
 Baltic, Br. Isles
- L. marmoratus* Curtis,1830
 N. Baltic
- L. decipiens* (Kolenati,1848)
 N. Baltic
- L. lunatus* Curtis,18??
 N. Baltic
- L. ignavus* McLachlan,18??
 N. Baltic
- Phryganeidae
Phryganea Linnaeus,1758
P. grandis Linnaeus,1758
 N. Baltic
- P. striata* Linnaeus,1758
 N. Baltic
- Agrypnia* Curtis,1835
A. pagetana Curtis,1835
 N. Baltic
- Agrypnetes* McLachlan,1876
A. sp.
 Baltic
- Molannidae
Molanna Curtis,1834
M. angusta (Curtis,1828?)
 N. Baltic
- Sericostomatidae
Lepidostoma Rambur,1842
L. hirtum (Fabricius,17??)
- Leptoceridae
Atripsodes Billberg,1820
A. aterrimus (Stephens,18???)
 Baltic
- Ocetus* McLachlan,1877
O. ochracea Curtis,1925
 SW Sjælland, in 10 ppt S.
O. furva (Rambur,1842?)
 N. Baltic
- ?Superfamilies? & ?Families?
Grammotaulius Kolenati,1848
G. atomarius (J.C. Fabricius,1793), ?non *Phryganea atomaria* J.F. Gmelin,1790
 N. Baltic
- Colpotaulius* Kolenati,1848
C. incisus (Curtis,1834)
 N. Baltic
- Ptoceridae
 * Taxa from a "Baltic Invertebrate Code-list". Not listed in Parker (Ed.),1982
- Lycentropus* ?,1???
- L.* sp.
 Baltic
- Raylea* ?,1???
- R.* sp.
 Baltic
- Agraylea Curtis,1834
A. sp.
 Baltic
- LEPIDOPTERA "Fjärilar"
 * This part of the list is certainly much incomplete. Only some, more or less obligate, associates of maritime herbs are listed.
- DITRYSIA
- Tortricoidae "Vecklare"
- Tortricidae Stephens,1829
- Olethreutinae Walsingham,1897
Olethreutini Obraztsov,1946
- Bactra* Stephens,1834 (TM *Olethreutes lancealana* Hübner,1796-99)
- B. robustana* Christoph,1872
 @ *Scirpus maritimus*
 Bohuslän, Halland, Skåne, Denmark, Germany
- Lobesia* Guenée,1845 (TM *Asthenia reliquana* Hübner,1825)
- L. (Lobesia) littoralis* Westwood & Humphreys,1845
 @ *Armeria maritima*
 Bohuslän, Halland, Skåne, Denmark, Norway, Germany, Netherlands
- Eucosmini* Obraztsov,1946
- Eucosma* Hübner,1823 (TM *E. circulana* Hübner,1823)
- E. (Phaneta* Stephens,1852) *tripoliana* Barrett,1880 (TM of *Phaneta* is *Cochylis pauperana*
 @ *Aster tripolium* Duponchel,1842)
 W Jylland, N. Germany

- E. (Phaneta** Stephens,1852) *krygeri* Rebel,1937
@ *Artemisia maritima*
Sjælland (Amager)
- E. (Phaneta** Stephens,1852) *maritima* Westwood & Humphreys,1845
@ *Artemisia* spp.
N Germany
- Tortricinae
- Aethes** Billberg,1820
- A. dilucidana** ?,1???
@ *Angelica archangelica litoralis*
Bohuslän, Halland, Netherlands, Britain
- Phalonidia** Le Marchand,1933
* Placed in correct superfamily & family?
- P. affinitana** Dgl.,1???
@ *Aster tripolium*
Bohuslän, Halland, Skåne, Denmark
- P. vectisana** Humphreys & Westwood,1845
@ *Triglochin maritimum*
Halland, Skåne, Denmark, Finland
- Pterophoroidea
- Pterophoridae "Fjädermott"
- Agdistis** Hübner,1825
- A. benneti** (Curtis,18???)
@ *Limonium*
S. Sjælland
- Pyraloidea
- Pyralidae** "Mott"
- Acentropus** Curtis,1834
- A. niveus** (Olivier,1791)
* in sheltered, vegetation-rich brackish water (up to 20 ppt S) lagoons (& in fresh water)
Baltic, Danish estuaries
- Crambinae
- Pediasia** Hübner,1825
- P. aridella** (Thunberg,1788) "en gräsmotts-art"
* Halophilic.
Halland
- Tineoidea "Malar"
- Bucculatrigidae
- Bucculatrix** Zeller,1839
- B. maritima** Stainton,1851
@ *Aster tripolium*
Bohuslän, Halland-S Baltic shores, Denmark, Finland
- Gelechioidea
- Oecophoridae
- Depressiniae
- Agonopterix** Hübner,1825
- A. zephyrella** ?,1???
@ *Angelica archangelica litoralis*
Halland, Germany, Netherlands, Britain
- Gelechiidae
- Gelechiinae
- Scrobipalpa** Janse,1951
- S. salicorniae** (E. Hering,1???)
@ *Salicornia*, *Suaeda*, etc.
Halland, central Baltic shores, Denmark, Finland
- S. stangei** (E. Hering,1889)
@ *Triglochin maritimum*
Bohuslän, Halland-central Baltic Shores, Denmark, Finland
- S. plantaginella** (Stainton,18???)
@ *Plantago maritima*
Bohuslän, Halland-S Baltic shores, Denmark, Finland
- S. nitentella** (Fuchs,1902)
@ *Arriplex*, *Suaeda* & *Chenopodium*
Göteborg archipelago, Skåne-central Baltic shores, Denmark, Finland
- Aristolochiinae
- Monochroa** Heinemann,1870
- M. tetragonella** (Stainton,1885)
@ *Glaux maritima*
Bohuslän, Halland, Skåne, S Baltic shores, Denmark, Finland
- Aristolelia** Hübner,1825
- A. brizella** Treitschke,1833
@ *Armeria maritima*
Halland-central Baltic shores, Norway, Denmark, Finland
- Coleophoridae "Stäckdrägmalar"
- Coleophora** Hübner,1822
- C. atriplicis** Meyrick,1928
@ *Arriplex*, *Suaeda* & *Chenopodium*
Halland-central Baltic shores, Finland
- C. adjunctella** Hodgkinson,1882
@ *Juncus gerardi*
Halland-S & N Baltic shores, Denmark, Finland
- C. salicorniae** Wocke,1876
@ *Salicornia*
Halland, S Baltic shores, Denmark, Finland
- C. asteris** Mühlig,18???
@ *Aster tripolium*
Göteborg archipelago, Skåne, Denmark
- C. suaedivora** Durrant,1???
@ *Suaeda maritima*
E Småland, England
- Elachistidae
- Elachista** Treitschke,1833
- E. scirpi** Stainton,1???
@ *Scirpus maritimus* & *Juncus gerardi*
Bohuslän-S Baltic shores, Denmark
- DIPTERA "Tvåvingar"
* See also Yerbury, J. W., 1919. Seashore Diptera. *J. mar. biol Ass. U.K.* **12** : 141-145.
- NEMATOCERA "Myggor"
- Tipuloidea
- Tipulidae** "Harkrankar"
- Tipulinae "Egentliga harkrankar"
- Tipula** Linnaeus,1758
- T. solstitialis** ?,1???
* In brackish water ditches.
Britain
- Limoniiinae** "Småharkrankar"
- Limonia** Meigen,1803
- L. (Geranomyia** Haliday,1833) *unicolor* Haliday,1833
* Larvae in tufts of *Lichina pygmaea* and *Catenella repens*
S. England
- L. (Geranomyia** Haliday,1833) *bezzii* A. & L.,19??
* Larvae assumed to feed on *Enteromorpha*
Suffolk-Dorset, N. Africa
- L. (Dicranomyia** Stephens,1829) *goritiensis* (Mik,1???)
* In tiny streams, which make the cliff wet.
S. English coast
- L. sera** ?,19??
* Salt-marshes.
Britain
- L. complicata** ?,19??
* Salt-marshes.
Britain
- L. ventralis** ?,1???
* In brackish water ditches.
Britain
- Eriopterini
- Erioptera** Meigen,1803
- E. strictita** ?,1???
* In salt-marshes.
Britain
- E. bivittata** ?,1???
* In brackish water ditches.
Britain
- Gonomyia** Meigen,1818
- G. conoviensis** ?,19??
* In tiny streams, which make the cliff wet.
Britain
- Psychodoidea
- Ptychopteridae
= *Liriopaeidae* "Glansmyggor"
- Ptychoptera** Meigen,1803
- P. contaminata** van der Wulp,18???
* In brackish water ditches.
Sweden, Britain
- Culicoidea
- Chironomidae
= *Tendipedidae* "Fjädermyggor"
- Telmatogotoninae Brundin,1966
- Thalassomyia** Schiner,1856
- T. frauendorfii** Schiner,1856
Br. Isles, France, Mediterranean
- Psamathomyia** Deby,1889
- P. pectinata** Deby,1889
Br. Isles, French Mediterranean coast
- Telmatogeton** Schiner,1866
=? *Parachunio* Kieffer,1911

- T. remanei* Remmert,1963
Kiefer Bucht
- Orthocladiinae* Edwards,1929
- Clunio* Haliday,1855
- C. marinus* Haliday,1855
Baltic-Bohuslän, Belgium, France, Br. Isles
- Halocladius* Hirvenoja,1973
* Includes some former *Cricotopus* species with larvae in saline waters. Probably some of those beneath.
- H. fucicola* (Edwards,1926)
= *Cricotopus fucicola* (Edwards,1926)
* Intertidal, on rocky shores.
Br. Isles
- H. varians* (Staeger,1839)
= *Cricotopus halophilus* Kieffer,1909
= *Cricotopus marinus* Alverdes,1911
* Brackish water species. Muus,1967 found *Chironomus halophilus* Kieffer (Strenzke) in most Danish estuaries, between 1-16 ppt S. Is he referring to this species?
Kieler Bucht, Öresund, Bergen, Helgoland, Br. Isles, Belgium, Normandie, ?Faroes, ?Iceland
- Cricotopus* van der Wulp,1874
- C. obnixus* (Walker,1856)
SW. England
- C. annulator* Goetghebuer,1927
Baltic, Belgium
- C. bicinctus* (Meigen,1818)
= *Trichocladus dizonias* (Meigen,1830)
Baltic, Belgium
- C. caducus* Hirvenoja,19??
Baltic
- C. festivellus* (Kieffer,1906)
Baltic, Belgium
- C. flavocinctus* (Kieffer,1924)
Baltic, Belgium
- C. laetus* Hirvenoja,19??
Baltic
- C. obtusus* Hirvenoja,19??
Baltic
- C. ornatus* (Meigen,1818)
Baltic, Belgium
- C. pilatensis* (Zetterstedt,1850)
Baltic, Belgium
- C. pilosellus* Brundin,19??
Baltic
- C. sylvestris* (Fabricius,1794)
Baltic, Belgium
- C. triannulatus* (Macquart,1826)
Baltic, Belgium
- C. trifascia* Edwards,1929
Baltic, Belgium
- Psectrocladius* Kieffer,1906
- P. barbimanus* Edwards,1929
* Brackish water species
England
- P. sordidellus* (Zetterstedt,1838)
Baltic, Belgium
- P. ventricosus* Kieffer,19??
Baltic
- Smittia* Holmgreen,1869
- S. rupicola* (Kieffer, in Mercier,1923)
Germany, Belgium, France
- S. bacilligera* (Kieffer,1924)
German Baltic
- Pseudosmittia* Goetghebuer,1932
- P. brevifurcata* (Edwards,1926)
England
- Thalassosmittia* Strenzke & Remmert,1957
- T. thalassophila* (Bequaert & Goetghebuer,1913)
Belgium, England
- Corynoneura* Winnertz,1846
- C. marina* Kieffer,1924
Germany
- C. scutellata* Winnertz,1846
Baltic
- Metricnemus* van der Wulp,1874
- M. picipes* (Meigen,1818)
Scandinavia, Germany, Netherlands, Belgium, France, SW. England
- Paraphaenocladius* Thienemann,1924
- P. impensus* (Walker,1856)
SW. England
- Abiskomyia* Edwards,1937
- A. virgo* Edwards,19??
Baltic
- Acricotopus* Kieffer,1921
- A. thienemanni* Goetghebuer,19??
Baltic
- Camptocladius* van der Wulp,1874
- C. sp.*
Baltic
- Eukiefferiella* Thienemann,1926
- E. sp.*
Baltic
- Heterotanytarsus* Spärck,1923
- H. apicalis* Kieffer,19??
Baltic
- Heterotriassocladus* Spärck,1923
- H. grimshawi* (Edwards,1929)
Baltic, Belgium
- H. marcidus* (Walker,1856)
Baltic, Belgium
- H. maeae* Brundin,19??
Baltic
- H. subpilosus* (Kieffer,19??)
Baltic
- Limnophyes* Eaton,1875
- L. sp.*
Baltic
- Microcricotopus* Thienemann,1932
- M. balticus* Palmen,19??
Baltic
- M. bicolor* (Zetterstedt,18??)
Baltic
- Orthocladius* van der Wulp,1874
- O. excavatus* Brundin,19??
Baltic
- Parakiefferiella* Thienemann,1936
- P. bathopila* Kieffer,19??
Baltic
- Rheorthocladius* Thienemann,1935
- R. sp.*
Baltic
- Syncretotopus* Brundin,1956
- S. rufiventris* (Meigen,18??)
Baltic
- Trissocladius* Kiefer,1908
- T. sp.*
Baltic
- Tanypodinae Thienemann & Zavrel,1916
- Ablabesmyia* Johannsen,1905
- A. monilis* (Linnaeus,1758)
Baltic
- A. barbitarsis* (Zetterstedt,18??)
Baltic
- A. cingulata* (Walker,1???)
Baltic
- A. melanops* (Meigen,18??)
Baltic
- Anatopynia* Johannsen,1905
- A. plumipes* (Fries,1823)
Baltic, Belgium
- Clinotanytarsus* Kieffer,1913
- C. nervosus* (Meigen,1818)
Baltic, Belgium
- Macropelopia* Thienemann, in Thienemann & Kieffer,1916
- M. nebulosa* (Meigen,1804)
Baltic, Belgium
- Arctopelopia* Fittkau,1962
- A. griseipennis* (Van der Wulp,1858)
Baltic
- Procladius* Skuse,1889
- P. breviatus* Remmert,1???
- Baltic*
- P. barbatus* Brundin,19??
Baltic
- P. crassinervis* (Zetterstedt,18??)
Baltic
- P. flavifrons* Edwards,19??
Baltic

- P. lundstroemi* Goetghebuer,18??
Baltic
- P. nigriventris* Kieffer,1???
Baltic
- P. nudipennis* Brundin,19??
Baltic
- Psectrotanypus* Kieffer,1909
- P. varius* (Fabricius,1787)
Baltic, Belgium
- Holotanypus* Roback,1982
- H. choreus* (Meigen,1804)
Baltic, Belgium
- H. signatus* (Zetterstedt,1850)
Baltic, Belgium
- Psilotanypus* Kieffer,1906
- P. sp.*
Baltic
- Tanypus* Meigen,1803
= *Tanipus* Rondani,1856
- T. punctipennis* Meigen,1818
Baltic
- Pentaneurini
- Thienemannimyia* Fittkau,1957
- T. fusciceps* Edwards,1???
Baltic
- T. lentiginosa* Fries,1???
Baltic
- T. vitellina* (Kieffer,18???)
Baltic
- Diamesinae* Kieffer,1923
- Diamesa* Meigen,1830
- D. sp.*
Baltic
- M. ekmani* Brundin,19??
Baltic
- Protanypus* Kieffer,1906
- P. morio* (Zetterstedt,18???)
Baltic
- Pothastia* Kieffer,1922
- P. sp.*
Baltic
- Prodiamesinae* Sæther,1976
- Prodiamesa* Kieffer,1906
- P. olivacea* (Meigen,1818)
Baltic, SW England
- Monodiamesa* Kieffer,1921
- M. bathyphila* Kieffer,1???
Baltic
- Chironominae* Macquart,1838
- Campochironomus* Kieffer,1918
- C. pallidivittatus* Malloch,1???
Baltic
- C. tentans* (Fabricius,1805)
Baltic, Belgium
- Chironomus* Meigen,1803
= *Tendipes* Meigen,1800
- C. salinarius* Kieffer,19??
* The larvae thrive in laboratory cultures with up to 50 ppt S.
Baltic, dominating species in Danish estuaries.
- C. halophilus* Meigen,18??
Baltic, Westfalen
- C. bicornutus* ?,1???
Westfalen
- C. aprilinus* Hoffmannsegg, in Meigen,1830
?Br. Isles
- C. annularius* Meigen,1818
= *C. annularis* (De Geer,1776)
Baltic, Belgium
- C. plumosus* (Linnaeus,1758)
Baltic
- C. anthracinus* Zetterstedt,18??
= *C. bathophilus* Kieffer,19??
Baltic
- C. thummi* Kieffer,1911
=? *C. gregarius* Meigen,18??
Baltic
- C. dorsalis* Meigen,18??
Baltic
- C. tenuistylus* Brundin,19??
Baltic
- Cryptochironomus* Kieffer,1918
- C. supplicans* (Meigen,1830)
Baltic, Belgium
- C. albifasciatus* (Staeger,1839)
Baltic, Belgium
- C. psittacinus* (Meigen,1830)
Baltic, Belgium
- Cryptocladopelma* Lenz,1941
- C. viridula* (Fabricius,17???)
Baltic
- C. edwardsi* Krus.,19??
Baltic
- Cryptotendipes* Lenz,1941
- C. usmaensis* (Pagast,1931?)
* Is this not a synonym of *Tanytarsus usmaensis* Pagast,1931?
Baltic
- C. paripes* Edwards,19??
Baltic
- Demicryptochironomus* Lenz,1941
- D. vulneratus* (Zetterstedt,1838)
Baltic, Belgium
- Einfeldia* Kieffer,1924
- E. insolita* Kieffer,19??
=? *E. dissidens* (Walker,1856)
Baltic
- Endochironomus* Kieffer,1918
- E. dispar* (Meigen,1830)
Baltic, Belgium
- E. impar* (Walker,18???)
Baltic
- E. intextus* (Walker,18???)
Baltic
- E. tendens* (Fabricius,1775)
Baltic, Belgium
- E. albibennis* (Meigen,1830)
=? *E. marinimus* Kieffer,1921
Baltic
- Fleuria* Kieffer,1924
- F. sp.*
Baltic
- Harnischia* Kieffer,1921
- H. curtilamellata* (Malloch,1915)
= *H. pseudosimplex* Goetghebuer,1923
Baltic, Belgium
- H. viridula* (Linnaeus,1767)
Baltic
- Kribioxenus* Kieffer,1921
- K. brayi* Goetghebuer,19??
Baltic
- Lauterborniella* Thienemann & Bause,1913
- L. agrayloides* (Kieffer,1911)
Baltic
- Lenzia* Kieffer,1922
- L. flavipes* (De Geer,1776)
Baltic
- L. punctipes* Wiedemann,1???
Baltic
- Dicrotendipes* Kieffer,1913
= *Limnochironomus* Kieffer,1920
- D. nervosus* (Staeger,1839)
Baltic, Belgium
- D. pulsus* (Walker,1856)
Baltic
- Microchironomus* Kieffer,1918
=? *Leptochironomus* Pagast,1931
- M. tener* (Kieffer,1918)
= *Leptochironomus balticus* Pagast,1931?
Baltic, Belgium
- Microtendipes* Kieffer,1915
- M. brevitarsis* Brundin,19??
Baltic
- M. chloris* (Meigen,1818)
Baltic, Belgium
- M. confinis* (Meigen,1830)
Baltic, Belgium
- M. pedellus* (De Geer,1776)
Baltic

<i>Pagastiella</i> Brundin,1949	<i>T. brundtini</i> Lindeberg,19?? Baltic
<i>P. orophila</i> (Edwards,1929) Baltic, Belgium	<i>T. chinyensis</i> Goetghebuer,1934 Baltic, Belgium
<i>Parachironomus</i> Lenz,1921	<i>T. curticornis</i> Kieffer,19?? Baltic
<i>P. arcuatus</i> (Goetghebuer,1919) Baltic, Belgium	<i>T. decipiens</i> Lindeberg,19?? Baltic
<i>P. mauricii</i> (Kruseman,1933) Baltic, Belgium	<i>T. eminulus</i> (Walker,1856) Baltic, Belgium
<i>P. swammerdami</i> (Kruseman,19??) Baltic	<i>T. gregarius</i> Kieffer,1909 Baltic, Belgium
<i>P. varus</i> (Goetghebuer,1921) Baltic, Belgium	<i>T. heusdensis</i> Goetghebuer,1923 Baltic, Belgium
<i>Paracladopelma</i> Harnisch,1923	<i>T. usmaensis</i> Pagast,1931 Baltic, Belgium
<i>P. camptolabis</i> (Kieffer,1913) Baltic	<i>T. holochlorus</i> Edwards,1929 Baltic, Belgium
<i>P. nigritula</i> Goetghebuer,19?? Baltic	<i>T. lestagei</i> Goetghebuer,1922 Baltic, Belgium
<i>P. obscura</i> Brundin,19?? Baltic	<i>T. lugens</i> Kieffer,19?? Baltic
<i>Paratendipes</i> Kieffer,1911	<i>T. norvegicus</i> Kieffer,19?? Baltic
<i>P. albimanus</i> (Meigen,1818) Baltic	<i>T. signatus</i> (van der Wulp,1859) Baltic, Belgium
<i>P. nudisquama</i> Edwards,19?? Baltic	<i>T. socialis</i> Lindeberg,19?? Baltic
<i>Paralauterborniella</i> Lenz,1941	<i>T. telmaticus</i> Lindeberg,19?? Baltic
<i>P. nigrohalteralis</i> Malloch,19?? Baltic	<i>Cladotanytarsus</i> Kieffer,1922
<i>Pentapedium</i> Kieffer,1913	<i>C. atridorsum</i> Edwards,19?? Baltic
<i>P. sordens</i> (van der Wulp,1874) Baltic, Belgium	<i>C. mancus</i> (Walker,1856) Baltic, Belgium
<i>P. tritum</i> (Walker,1856) Baltic, Belgium	<i>C. wexionensis</i> Brundin,19?? Baltic
<i>Polypedilum</i> Kieffer,1912	<i>Constempellina</i> Brundin,1947
<i>P. bicrenatum</i> Brundin,19?? Baltic	<i>C. brevicosta</i> Edwards,19?? Baltic
<i>P. convictum</i> (Walker,1856) Baltic, Belgium	<i>Corynocera</i> Zetterstedt,1837
<i>P. cultellatum</i> Goetghebuer,1931 Baltic, Belgium	<i>C. ambigua</i> Zetterstedt,18?? Baltic
<i>P. laetum</i> (Meigen,1818) Baltic, Belgium	<i>Lauterbornia</i> Kieffer,1911
<i>P. nubeculosum</i> (Meigen,1804) Baltic, Belgium	<i>L. coracina</i> (Zetterstedt,18???) Baltic
<i>P. pullum</i> (Zetterstedt,1838) Baltic, Belgium	<i>Micropsectra</i> Kieffer,1909
<i>P. scalaenum</i> (Schrank,1803) Baltic, Belgium	<i>M. groenlandica</i> Andersen,1937? Baltic
<i>Pseudochironomus</i> Malloch,1915	<i>M. monticola</i> Edwards,19?? Baltic
<i>P. prasinatus</i> (Staeger,1839) Baltic, Belgium	<i>M. praecox</i> (Meigen,18???) = <i>M. inermipes</i> Kieffer,19?? Baltic
<i>Sergentia</i> Kieffer,1922	<i>Paratanytarsus</i> Tienemann & Bause,1913
<i>S. coracina</i> (Zetterstedt,18???) Baltic	<i>P. confusus</i> Palmen,19?? Baltic
<i>S. longiventris</i> Kieffer,19?? Baltic	<i>P. inopertus</i> (Walker,1856) Baltic, Belgium
<i>Stenochironomus</i> Kieffer,1919	<i>P. intricatus</i> (Goetghebuer,1921) Baltic, Belgium
<i>S. sp.</i> Baltic	<i>P. laccophilus</i> Edwards,19?? Baltic
<i>Stictochironomus</i> Kieffer,1919	<i>P. natvigi</i> Goetghebuer,19?? Baltic
<i>S. crassiforceps</i> Kieffer,19?? Baltic	<i>P. penicillatus</i> Goetghebuer,19?? Baltic
<i>S. histrio</i> (Fabricius,1794) Baltic, Belgium	<i>P. tenuis</i> (Meigen,1830) Baltic, Belgium
<i>S. pictulus</i> (Meigen,1830) Baltic, Belgium	<i>Rheotanytarsus</i> Thienemann & Bause,1913
<i>S. rosenschoeldi</i> (Zetterstedt,18???) Baltic	<i>R. sp.</i> Baltic
<i>Xenochironomus</i> Kieffer,1921	<i>Stempellina</i> Thienemann & Bause,1913
<i>X. xenolabis</i> (Kieffer,1916) Baltic, Belgium	<i>S. bausei</i> (Kieffer,19???) Baltic
<i>Glyptotendipes</i> Kieffer,1913	<i>S. subglabripennis</i> Brundin,19?? Baltic
<i>G. barbipes</i> (Staeger,1839) Baltic, Norway	<i>Stempellinella</i> Brundin,1947
<i>Tanytarsus</i> van der Wulp,1874	<i>S. brevis</i> Edwards,19??
<i>T. brunnipes</i> (Zetterstedt,1850) Scandinavia, SW. England	

Baltic	Sweden, Br. Isles, European mainland
<i>S. minor</i> (Edwards,1929) Baltic, Belgium	<i>C. (Culiseta) subochrea</i> (Edwards,1921) * Larvae may tolerate slightly brackish water? Scandinavia, Br. Isles, European mainland
<i>Zavrelia</i> Kieffer, in Thienemann & Bause, 1913	<u>Anophelinae</u>
<i>Z. pentatoma</i> Kieffer, 1913 Baltic, Belgium	<i>Anopheles</i> Meigen, 1818
<u>Heleidae</u> = Ceratopogonidae "Svidknott"	<i>A. (Anopheles) atroparvus</i> van Thiel, 1927 * The only North European malaria vector. Earlier considered to be a subspecies of <i>A. maculipennis</i> Meigen, 1818. Coastal distribution. Tolerates high salinities. Sweden, S. Baltic, German North Sea coast, Netherlands
<i>Culicoides</i> Latreille, 1809	<i>A. (Anopheles) algeriensis</i> Theobald, 1903 * Not coastal, but larvae may develop in low salinity brackish water. Br. Isles, N. Germany, Mediterranean countries
<i>C. furens</i> (Poey, 18??) Br. Isles	<i>A. (Anopheles) claviger</i> (Meigen, 1804) * Not coastal, but larvae may develop in low salinity brackish water. Br. Isles, Ireland, European mainland
<i>C. sp.</i> Baltic	<u>Chaoboridae</u> = Corethridae "Tofsmyggor"
<i>Dasyhelea</i> Kieffer, 1911	<u>Chaoborinae</u>
<i>D. flavoscutellata</i> (Zetterstedt, 1850) Scandinavia, Britain	<i>Chaoborus</i> Lichtenstein, 1800
<i>Monoculicoides</i> Khalaf, 1954	<i>C. (C.) crystallinus</i> (De Geer, 1776) = <i>C. plumicornis</i> (Fabricius, 1794) = <i>C. fuscus</i> (Staeger, 1839) Baltic, Belgium
<i>M. nubeculosus</i> (Meigen, 18??) Britain	<i>C. (C.) flavidans</i> (Meigen, 1830) Baltic, Belgium
<i>Bezzia</i> Kieffer, 1899	<u>BRACHYCERA</u> "Flugor" * Almost only halobionts have been listed
<i>B. sp.</i> Baltic	<u>ORTHORRHAPHA</u>
<i>Palpomyia</i> Meigen, 1818, ex Megerle MS	<u>Tabanoidea</u>
<i>P. sp.</i> Baltic	<i>Tabanidae</i> "Bromsar"
<i>Sphaeromais</i> ?,1???	<u>Tabaninae</u>
<i>S. sp.</i> Baltic	<i>Tabanus</i> Linnaeus, 1758
<i>Leptoconops</i> Skuse, 1889	<i>T. bovinus</i> Linnaeus 1758 "Fäbroms", "Nötbroms" Sweden, German Baltic coast
<u>Dixidae</u>	<i>Hybomitra</i> Enderlein, 1922
<i>Dixa</i> Meigen, 1818	<i>Chrysopsinae</i> "Blinningar"
Subgen.: <i>Dixella</i> Dyar & Shannon, 1924	<i>Chrysops</i> Meigen, 1800
<i>D. (Dixella) attica</i> Pandazis, 19?? Britain	<i>C. relicta</i> Hoffmannsegg, in Meigen, 1820 "Blindbroms", "Guldögonbroms" * In salt-marshes Sweden, Germany
<i>Culicidae</i> "Stickmyggor"	<i>Haematopota</i> Meigen, 1803 "Egentliga blinningar"
<u>Culicinae</u>	<i>H. grandis</i> ?Macquart, 1834? * In salt-marshes Britain
<i>Aedes</i> Meigen, 1818, ex Hoffmannsegg MS	<i>H. bigota</i> ?,1???
<i>A. detritus</i> (Haliday, 1833) = <i>A. salinus</i> Ficalbi, 1896 = <i>Ochlerotatus</i> Lynch Arribálzaga, 1891 <i>detritus</i> (Haliday, 1833) * Chiefly coastal. Tolerates high salinities. S. Baltic, Denmark, Br. Isles, Ireland, France	* In salt-marshes Britain
<i>A. caspius</i> (Pallas, 1771) = <i>Ochlerotatus</i> Lynch Arribálzaga, 1891 <i>caspicus</i> (Pallas, 1771) * A salt tolerant species which also commonly is found in non-saline inland waters. along the European coast from Finland-England-Atlantic-Mediterranean	<i>Silvius</i> Meigen, 1820
<i>A. dorsalis</i> (Meigen, 1830) = <i>Ochlerotatus</i> Lynch Arribálzaga, 1891 <i>dorsalis</i> (Meigen, 1830) * Not coastal, but larvae may develop in low salinity brackish water. Br. Isles, N. & central Europe	<u>Stratiomyidae</u> "Vapenflugor"
<i>A. flavescentes</i> (O.F. Müller, 1764) * Not coastal, but larvae may develop in low-medium salinity brackish water. Br. Isles, N. & central Europe	<u>Stratiomyinae</u>
<i>A. leucomelas</i> (Meigen, 1804) * Not coastal, but larvae may develop in low-medium salinity brackish water. Br. Isles, N. Europe	<i>Stratiomys</i> Geoffroy, 1762
<i>Culex</i> Linnaeus, 1758	<i>S. longicornis</i> ?,1???
<i>C. (Barraudius</i> Edwards, 1921) <i>modestus</i> Ficalbi, 1889 * Not coastal, but larvae may develop in low salinity brackish water. Br. Isles, Mediterranean countries	* In brackish water ditches. Britain
<i>C. (Culex) pipiens</i> Linnaeus, 1758 * Not coastal, but larvae may develop in low salinity brackish water. Sweden, Br. Isles, temperate areas of the world	<i>S. furcata</i> Fabricius, 1???
<i>C. (Culex) torrentium</i> Martini, 1925 * Not coastal, but larvae may develop in low salinity brackish water. Br. Isles, N. & central Europe	* In brackish water ditches. Sweden, Britain
<i>C. (Neoculex</i> Dyar, 1925) <i>territans</i> Walker, 1856 * Not coastal, but larvae may develop in low salinity brackish water. Baltic area, Br. Isles, European mainland	<u>Clitellarinae</u>
<i>Culiseta</i> Felt, 1904 = <i>Theobaldia</i> Neveu-Lemaire, 1902	<i>Nemotelus</i> Geoffroy, 1762
<i>C. (Allotheobaldia</i> Brolemann, 1919) <i>longiareolata</i> (Macquart, 1838) * Larvae may develop in both fresh and brackish water. Br. Isles, Mediterranean countries	<i>N. uliginosus</i> ?,1???
<i>C. (Culisella</i> Felt, 1904) <i>liorea</i> (Shute, 1928) * Larvae tolerates slightly brackish water. Br. Isles, France, Spain	* Larvae develop in brackish water. Sweden, Britain
<i>C. (Culisella) morsitans</i> (Theobald, 1901) * Larvae tolerates slightly brackish water. Scandinavia, Br. Isles, European mainland	<i>N. notatus</i> ?,???
<i>C. (Culiseta) annulata</i> (Schrank, 1776) * Larvae tolerates brackish water. Our largest Culicidae species.	* Larvae develop in brackish water. Britain
	<u>Asiloidea</u>
	<i>Asilidae</i> Leach, 1819 "Rovflugor"
	<i>Asilinae</i> Macquart, 1838, emend. Schiner, 1862
	<i>Philonicus</i> Loew, 1849
	<i>P. albiceps</i> (Meigen, 1820) * Coastal sand dunes. Prey is peculiar to seashore. ?Scandinavia, Britain, Germany
	<u>Empidoidea</u>
	<i>Empididae</i> "Dansflugor"
	<i>Hilara</i> Meigen, 1822
	<i>H. lundbecki</i> Frey, 19??

Sweden, Denmark, Germany, Netherlands, Br. Isles

Cersodromia Walker,1849, ex Haliday MS

* In wrack-beds (in moist wrack).

C. arenaria (Haliday,1833)

Sweden, Denmark, Norway, Finland, Germany, Netherlands, S. England

C. difficilis Lundbeck,1910

Sweden, Denmark, Finland, France, S. England

C. incana Walker,18??

Sweden, Denmark, Netherlands, France, S. England

C. speculifera Walker,18??

Denmark, Germany, France, S. England

C. cursitans (Zetterstedt,1842)

Bohuslän, Denmark, Finland, Netherlands, France, S. England

Coloboneura Melander,1902

C. hirta (Walker,18???)

= *Cersodromia hirta* Walker,18???

Sweden, Denmark, Germany, Netherlands, France, S. England

Dolichopodidae "Stylflugor"

Dolichopodinae

Dolichopus Latreille,1796

D. sabinus ?,1???

* In salt-marshes

Britain

D. plumipes (Scopoli,1763)

= *Musca plumipes* Scopoli,1763

* In salt-marshes

Sweden, Britain

Hygrocoleuthus Loew,1857

H. latipennis (Fallén,1823)

Sweden, Norway, Finland, Balticum, Denmark, Germany, Netherlands, Belgium

H. diadema (Haliday,1832)

Sweden, Finland, Balticum, Denmark, Germany, Netherlands, Belgium, France, Br. Isles

Hydrophorinae

Hydrophorus Fallén,1823

H. bisetus Loew,18???

Denmark, Germany, Netherlands, France, Br. Isles

H. praecox (Lehmann,18???)

Sweden, Finland, Balticum, Denmark, Germany, Netherlands, Belgium, France, S. England

H. oceanus (Macquart,1???)

* Bare mud in salt-marshes.

S. England

Liancalus Loew,1857

L. virens (Scopoli,1763)

= *Musca virens* Scopoli,1763

* In tiny streams, which make the cliff wet.

Sweden, Britain

Thinophilus Wahlberg,1844

T. flavipalpis Zetterstedt,1844, ?non Schiodte,1844

Sweden, Finland, Balticum, Germany, Netherlands, France, Br. Isles

T. ruficornis Haliday,18???

Sweden, Finland, Balticum, Denmark, Germany, Netherlands, France, Br. Isles

Schoenophilus Mik,1878

S. versutus Walker,18???

Sweden (Skåne), Denmark, Germany, Netherlands, Belgium, France, Br. Isles

Subfamily?

Muscidideicus Becker,1917

M. praetextatus Haliday,18???

German North Sea coast, Belgium, France, Br. Isles

Orthoceratium Schrank,1803

O. lacustre (Scopoli,1763)

German North Sea coast, Belgium, Br. Isles

Aphrosylus Haliday, in Walker,1851

* Larvae may be predators on barnacles

A. ferox Haliday, in Walker,1851

Netherlands, France, S. England

A. mitis Verrall,18???

France, S. England

A. celtiber Haliday,1851?

* Is the name *A. celticer*?

France, S. England

A. raptor Haliday, in Walker,1851

France, S. England

A. venator Loew,18???

France

Machaerium Haliday,1832

M. maritimae Haliday,1832

* In salt-marshes

Denmark, Germany, France, S. England

CYCLORRAPHA

ASCHIZA

Phoroidea

Phoridae "Puckelflugor"

Syrphoidea

Syrphidae "Blomflugor"

Lathyrophthalmus Mik,1897

L. aeneus (Scopoli,1763)

= *Musca aena* Scopoli,1763

= *Eristalinus* Rondani,1845 *aeneus* (Scopoli,1763)

* In salt-marshes. Larvae probably in small pools in the splash zone.
Britain, Germany

Eristalis Latreille,1804 "Slamflugor"

E. abusivus ?,1???

* In salt-marshes.

Britain

E. sepulchralis (Linnaeus,1758)

= *Musca sepulchralis* Linnaeus,1758

@ on thrift blossoms (*Armeria maritima*)

Sweden, S. England

Helophilus Leach,1817 (Emend. pro *Elophilus* Meigen,1903)

H. parallelus ?,1???

* In brackish water ditches.

Britain

Lejops Rondani,1857

= *Liops* Verrall,1901, *non* Fieber,1870 (Hemiptera)

L. vittata (Meigen,18???)

* In brackish water ditches.

Sweden, Britain

Tropidia Meigen,1822

T. scita ?,1???

* In brackish water ditches.

Britain

CHIZOPHORA

Tephritoidea

Otitidae

= *Ortalidiidae* "Fläckflugor"

Melieria Robineau-Desvoidy,1830

= *Meliera* Becker,1903

M. omissa Meigen,18???

Finland, Denmark, Germany, Netherlands, Belgium, France, Br. Isles

M. picta (Meigen,1826)

Germany, Netherlands, France, Br. Isles

M. cana Loew,18???

Germany, Netherlands, France, Br. Isles

Ceroxys Macqurt,1835

C. urticae ?,1???

* In brackish water ditches.

Britain

Tephritidae

= *Trypetidae*

= *Trypaneidae* "Fruktflugor", "Borrflugor"

Paroxyna Hendel, in Lindner,1927

P. plantaginis ?,1???

@ *Aster tripolium*

Britain

Sciomyzoidea

Sciomyzidae "Kärrflugor"

Salicella Robineau-Desvoidy,1830

S. fasciata (Meigen,1830)

N. France, Ireland, S. Europe

Helcomyzidae

Helcomyza Curtis,1825

H. ustulata Curtis,1825, ex Meigen MS

* One of the main species in wrack-beds (in fairly dry wrack)

Sweden, Finland, Denmark, Germany, Netherlands, Belgium, France, Scotland

Heterochaeta Rondani,1857

H. buccata (Fallén,1820)

* One of the main species in wrack-beds (in moist wrack).

Skåne, Norway, Finland, Balticum, Denmark, Germany, Netherlands, Belgium, France, Br. Isles

Chamaemyiidae "Markflugor"

Chamaemyia Meigen,1803

C. maritima (Zetterstedt,1846)

Sweden, Denmark, Germany, Netherlands, Belgium, France, Br. Isles

C. flavipalpis Haliday,18???

Britain

Coelopidae "Tångflugor"

* 6 Swedish species.

Coelopa Meigen,1830

= *Fucomyia* Haliday, in Curtis,1837

- C. frigida** (J. C. Fabricius,1805)
 = *C. parvula* Haliday,18??
 = *C. eximia* Stenhammar,18??
 * One of the main species in wrack-beds (in moist wrack).
 Sweden, Norway, Finland, Balticum, Denmark, Germany, Netherlands, Belgium, France, Br. Isles
- C. pilipes** Haliday,1838
 * One of the main species in wrack-beds (in moist wrack).
 Sweden, Finland, Germany, Belgium, France, S. England, Scotland
- Malacomysia** Westwood,1840, *ex* Haliday MS
 = *Malacomysza* Curtis,1837, *ex* Haliday MS, *non* Wesmael,1836 (Neuroptera)
- M. sciomyzina** (Haliday,1833)
 = *Coelopa furcata* Zetterstedt,1847
 * One of the main species in wrack-beds (in moist wrack).
 Bohuslän, Balticum, Germany, Netherlands, France, S. England
- Oedoparea** Loew,1862
 = *Heteromyza* Fallén,1820 (p.p.)
- O. buccata** (Fallén,1820)
 = *Heteromyza buccata* Fallén,1820
 * One of the main species in wrack-beds. Is this species possibly identical with *Heterochaeta b.* ?
 Sweden, S. England, Scotland
- Sepsidae** "Svängflugor"
- Orygmatinae**
- Orygma** Meigen,1830
- O. luctuosa** Meigen,1830
 * One of the main species in wrack-beds (in moist wrack).
 W. Sweden, Norway, Finland, Denmark, Germany, Netherlands, Belgium, France, S. England, Scotland
- Drosophiloidae**
- Drosophilidae**
- Drosophila** Fallén,1823
- D. funebris** (J.C. Fabricius,1787) "Stora ättikflugan"
 * In banks of decaying seaweed.
 Sweden, Scotland
- D. phalerata** Meigen,18??
 S. England
- Ephydriidae** "Vattenflugor"
- Hecamede** Haliday, *in* Curtis,1838
- H. albicans** (Meigen,1830)
 Sweden, Denmark, Balticum, Germany, Netherlands, Belgium, France, S. England
- Clanoneurum** Becker,1903
- C. cimiciformae** (Haliday,18??)
 Germany, Belgium, France, England, Mediterranean
- Atissa** Haliday, *in* Curtis,1837
- A. limosina** Becker,1???
 Norway, Finland, France, Br. Isles
- A. pygmaea** (Haliday,1833)
 ?Denmark, Germany, Netherlands, Belgium, France, Br. Isles
- Glenanthe** Haliday,1839
- G. fuscinervis** Becker,1???
 Norway (Molde), Finland, Germany
- G. ripicola** Haliday,1839
 Finland, Germany, Belgium, France, Br. Isles
- Philygria** Stenhammar,1844
- P. ochracea** Oldenberg,1???
 Borkum
- Ephydria** Fallén,1810 "Saltflugor"
- E. riparia** Fallén,1813
 Sweden, Norway, Finland, Balticum, Denmark, Germany, Netherlands, Belgium, France, Br. Isles
- E. salinea** Zetterstedt,1846
 Sweden (so far only in inland salt water)
- Scatella** Robineau-Desvoidy,1830
- S. stagnalis** Fallén,18??
 Sweden, Norway, Finland, Balticum, Denmark, Germany, Netherlands, Belgium, France, Br. Isles
- S. subguttata** (Meigen,1830)
 Sweden, Norway, Finland, Balticum, Denmark, Germany, Netherlands, Belgium, France, Br. Isles
- Halmopota** Haliday,1856
- H. salinarum** Bché,1???
 Germany, Br. Isles
- Lamproscatella** Hendel,1917
- L. quadrisetosa** (Becker,1896)
 European littoral
- Parydroptera** Collin,1913
- P. discomyzina** ?,1???
 * In salt-marshes.
 Britain
- Chloropoidea**
- Tethinidae**
- Pelomyia** Williston,1893
- P. cinerella** (Haliday,18???)
- Finland, ?Denmark, Germany, Netherlands, France, Br. Isles
- P. angustifacies** de Meij,1???
 Germany, Netherlands, Br. Isles
- Tethnia** Haliday, *in* Curtis,1836
 = *Tethina* Haliday,1838
- T. cinerea** Loew,18???
 ?Denmark, Germany, Netherlands, France, Br. Isles
- T. grisea** (Fallén,1810?)
 Sweden, Finland, Germany, Netherlands, Br. Isles
- T. illota** (Haliday,1836?)
 Finland, ?Denmark, Amrum, Netherlands, France, Br. Isles
- T. griseola** v. d. Wulp,18???
 Netherlands, Belgium, France, Br. Isles
- T. pictipes** Becker,1???
 Br. Isles
- T. nigripes** Cz.,1???
 Germany
- T. longirostris** Loew,18???
 France
- Canaceidae** "Surfflugor"
- Canace** Haliday, *in* Curtis,1838
- C. nasica** Haliday,18???
 * Breeding in *Enteromorpha*
 Belgium, France, Br. Isles
- Xanthocanace** Hendel.,1914
- X. ranula** Loew,18???
 ?Denmark, Germany, Netherlands, Belgium, France, Br. Isles
- Chloropidae** "Fritflugor"
- Calamonocosis** Enderlein,1911
- C. tomentosa** (Macquart,1???)
 S. England
- Muscoidea**
- Scathophagidae**
 = *Cordyluridae* "Kolvflugor"
- Cordylurinae**
- Scopeuma** Meigen,1800
 = *Scathophaga* Meigen,1803
- S. litoreum** Fallén,1823?
 * In moist wrack. ?Identical with *Scatomyza l.* ?
 Sweden, Norway, Finland, Balticum, Denmark, Germany, Netherlands, Belgium, France, Br. Isles
- S. decipiens** (Haliday, *in* J. Curtis,1832)
 Germany, Netherlands, Br. Isles, Ireland, N. France
- S. islandicum** Becker,18???
 Norway, Iceland, N. France
- S. villipes** (Zetterstedt,1846)
 * Under dead seaweed.
 Sweden, Norway, S. England, Scotland
- S. rufiventre** Villeneuve,19???
 France
- S. calida** ?,1???
 * In moist wrack.
 Britain
- Ceratinostoma** Meade,1885
- C. ostiorum** (Haliday, *in* Curtis,1832)
 * In moist wrack.
 Sweden, Norway, ?Denmark, Germany, Netherlands, Belgium, France, S. England
- Calliphoridae** "Spyflugor"
- Sarcophaga** Meigen,1826
- S. (Bellieria)** Robineau-Desvoidy,1863 *melanura* Meigen,1826
 S. England
- Muscidae** "Egentliga flugor"
- Mydaeinae**
- Limnophorinae**
- Limnophora** Robineau-Desvoidy,1830
- L. (Neolimnophora** Schnabl,1902) *maritima* (v. Röder,1887)
 ?Denmark, German North Sea coast, France, Br. Isles
- L. (N.) virgo** Villeneuve,1906
 German North Sea coast, France, Br. Isles
- L. (Spilogona** Schnabl,1911) *baltica* Ringdahl,1918
 Sweden, Finland, ?Denmark, Germany, Netherlands, Br. Isles
- L. (S.) glauca** Stein,1916
 Sweden, Finland, Germany, Belgium
- L. (S.) aerea** (Fallén,1825)
 Sweden, Norway, Finland, Denmark, Germany, Netherlands, France, S. England
- L. (S.) biseriata** Stein,1916
 Sweden, ?Denmark, Borkum, Netherlands, France, Br. Isles
- L. (S.) marina** Collin,18???
 ?Denmark, Germany, Netherlands, France, S. England

- L. (Villeneuvia Schnabl,1911) aestuum** Villeneuve,1902
?Denmark, German North Sea coast, Belgium, France, Br. Isles
- L. (Limnophora) setinervia** Schnabl,1911
= *L. excusaria* Stein,1916
France, S. England
- L. insularis** Collin,18??
Norway, Br. Isles
- Lispe** Latreille,1796
= *Lispa* Fallén,1810
- L. caesia** Meigen,1826
Sweden, Balticum, ?Denmark, German North Sea coast, Netherlands, Belgium, France, Br. Isles
- L. hydromyzina** Fallén,1825
Sweden, Finland, Balticum, ?Denmark, German coasts, Netherlands, Br. Isles
- L. litorea** Fallén,1825
Sweden, Finland, Balticum, Denmark, German North Sea coast, Netherlands, Belgium, France, Br. Isles
- L. loewi** Ringdahl,1922
Sweden, ?German North Sea coast, France, Br. Isles
- L. tentaculata** (C. De Geer,1776)
Britain
- Mydaeini
- Helina** Robineau-Desvoidy,1830
- H. protuberans** (Zetterstedt,1845)
Sweden, Norway, Finland, Denmark, Germany, Netherlands, Belgium, France, Br. Isles
- Coenosiiiae**
- Chelisiini
- Limnospila** Schnabl,1902
- L. albifrons** (Zetterstedt,1849)
Sweden, Denmark, Germany, France, Br. Isles
- Dexiopsis** Pokorný,1893
- D. lacustris** Karl,1???
Amrum, Br. Isles
- Coenosiiini
- Coenobia** Meigen,1826
- C. salinarum** Stein,1897
Germany, Netherlands, France, Br. Isles
- Anthomyiidae**
- Fucellini
- Fucellia** Robineau-Desvoidy,1842
- F. maritima** (Haliday,1838)
* One of the main species in wrack-beds
Sweden, Norway, Finland, Balticum, Denmark, Germany, Netherlands, Belgium, France, Br. Isles
- F. griseola** (Fallén,1819)
Sweden, Finland, Balticum, Denmark, German Baltic coast,
- F. fulciformis** (Fallén,1819)
Sweden, Norway, Finland, Denmark, German Baltic coast, S. England
- F. marina** ?1???
* Maybe just a lapsus for *F. maritima* ?
Sweden, Denmark
- Families yet lacking undoubted systematic placement
- Heleomyzidae** "Mylflugor"
= **Helomyzidae**
- Neoleria** Malloch,1919
- N. maritima** Villeneuve,19??
N. France
- Sphaeroceridae** "Hoppflugor"
= **Borboridae**
- Thoracochaeta** Duda,1918
- T. zosterae** (Haliday,1833)
= **Leptocera** Olivier,1813 *zosterae* (Haliday,1833)
* One of the main species in wrack-beds (in moist wrack).
Bohuslän, Norway, ?Denmark, Germany, Netherlands, France, S. England, Scotland
- T. brachystoma** (Stenhammar,18???)
* One of the main species in wrack-beds (in moist wrack).
S. England
- Scatomyza** Fallén,1810
- S. litorea** Fallén,1810
Sweden, S. England
- Collinellula** Strand,1928
- C. limosa** (Fallén,1820?)
= *Limosina limosa* Fallén,1820?
Balticum, ?Denmark, Germany, France, S. England
- Coprophila** Duda,1918
- C. vagans** (Haliday,1833)
S. England
- COLEOPTERA** "Skalbaggar"
* See also Keys, James H., 1918. A List of the Maritime, Sub-Maritime and Coast-frequenting Coleoptera of South Devon and South Cornwall, with especial reference to the Plymouth District. *J. mar. biol. Ass. U.K.*, 11 : 497-513
- Carabidae** "Jordlöpare"
* Around 50 British coastal species
- Cicindelinæ** Leconte & Horn,1883 "Sandjägare"
- Cicindela** Linnaeus,1758
- C. maritima** Latreille,1822-24 "Strandsandjägaren"
= *C. hybrida* Linnaeus,1758 var. *maritima* Latreille,1822-24
* Halophilic
Skåne, Halland, Värmland, Dalarna, Medelpad-Norrbotten, Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, France, Br. Isles
- C. hybrida** Linnaeus,1758 "Brun sandjägare"
Skåne-Bohuslän, Denmark
- Carabinae** Horn,1881 & Harpalinae "Egentliga jordlöpare"
- Broscus** Panzer,1813
- B. cephalotes** (Linnaeus 1758) "Läderbagge"
* Coastal. In maritime sand dunes.
Skåne-Gästrikland, Denmark, Germany, S. England
- Dyschirius** Panzer,1813, ex Bonnelli
- D. obscurus** (Gyllenhal,1827)
* Halophilic
Sweden, Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, France, ?Br. Isles
- D. thoracicus** (P. Rossi,1797)
= *D. arenosus* Stephens,1827
* On sand
Sweden, Denmark, Finland, German North Sea coast
- D. impunctipennis** Daws.,1854
* Halobiont
Sweden, Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, France, Br. Isles
- D. extensus** Putzeys,1846
* Halobiont
England, Germany
- D. chalcurus** Erichson,1837
* Halobiont
Sweden, Denmark, Germany, Netherlands, Belgium
- D. salinus** Schaub,1843
* Halobiont
Sweden, Norway, Denmark, Balticum, Germany, Netherlands, Belgium, Br. Isles, Arcachon
- D. globosus** Herbst,1???
* Eurytopic with preference for sandy shores.
Bohuslän-Uppsala, Åland, Finland
- Bembidion** Latreille,1802
= **Bembidium** Gyllenhal,1810
- B. (Actedium** Motschoulsky,1864 *pallidipenne* (Illiger,1802)
* Halophilic
Sweden, Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, France, Br. Isles
- B. (Peryphus** Stephens,1828 *saxatile* Gyllenhal,1827
* Halophilic
Sweden, Norway, Finland, Denmark, Balticum, Germany, Netherlands, Br. Isles
- B. (Peryphus** Stephens,1828 *femoratum* (Sturm,1825)
= *B. maritimum* Stephens,18???
* Coastal.
S. England, N Germany
- B. (Notaphus** Stephens,1828 *varium* (Olivier,1795)
* Sub-maritime
S. England, Germany
- B. (Emphanes** Motschoulsky,1864 *normannum* (Dejean,1821)
* Halobiont
Denmark, Germany, Netherlands, Belgium, S. England
- B. (Ocys** Stephens,1828 *harpalooides* (Serville,1821)
S. England, Germany
- B. (Notaphemphanes** Netolitzky,1920 *ephippium* (Marsham,1802)
* Halobiont
Denmark, Germany, Netherlands, Belgium, England
- B. (Talanes** Motschoulsky,1864 *aspericollis* (Germar,1829)
* Halobiont
Germany, Belgium, France
- B. (Philochthus** Stephens,1828 *aeneum* (Germar,1824)
* Halobiont
Sweden, Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, Br. Isles
- B. (Diplocampa** Bedel,1896 *fumigatum* (Creutzer, in Duftschmid,1812)
* Halobiont
Sweden, Denmark, Germany, Netherlands, Belgium, England
- B. concinnum** (Stephens,18???)
* Halobiont
Germany, Netherlands, Belgium, Br. Isles
- B. iricolor** Bedel,18??
* Halobiont
Netherlands, Belgium, France, England
- B. assimile** Gyllenhal,18???
* Hygrophilous. Especially abundant on sea shores.
Bohuslän-Skåne, Finland
- Clivina** Latreille,1802
- C. fossor** (Linnaeus,1758)
Bohuslän, Finland
- Trechus** Schellenberg,1806
- T. quadristratus** Schrank,1???
* Eurytopic on heath & shore.
Bohuslän, Finland
- ADEPHAGA**

<i>Cillenus</i> Samouelle,1819, <u>ex</u> Leach MS	<i>A. ruficornis</i> (Goeze,1777) = <i>Platynus</i> Stephens,1827, <u>ex</u> Bonelli,1810 (<i>Anchomenus</i> Stephens,1827, <u>ex</u> Bonelli,1810) <i>ruficornis</i> Bohuslän, S. England (Goeze,1777)
<i>C. lateralis</i> Samouelle,1819	
* An amphipod hunter; e.g. <i>Corophium</i> & <i>Talitrus</i> . Halobiont. Fully intertidal. Denmark, German North Sea coast, Netherlands, Br. Isles, Arcachon	
<i>Aepus</i> Samouelle,1819, <u>ex</u> Leach MS	<i>Microlestes</i> Schmidt-Goebel,1846
<i>A. marinus</i> (Ström,1783)	<i>M. maurus</i> (Sturm,1827)
* Maritime. Halobiont. Lives permanently between the tidemarks in crevices & beneath stones. S. Norway, Denmark, S. England, Scotland	* Coastal. S. England, Germany
<i>Dicheirotrichus</i> J. Du Val,1855	<i>Lionychus</i> Wissmann, <u>in</u> Schmidt-Goebel,1846
<i>D. gustavii</i> Crotch,18??	<i>L. quadrillum</i> (Creutzer, <u>in</u> K. Duftschmidt,1812)
= <i>Carabus pubescens</i> G. von Paykull,1798, non O.F. Müller,1776	* Sub-maritime. Sweden, S. England
* Sub-maritime. Halobiont. Sweden, Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, Br. Isles	
<i>D. obsoletus</i> (Dejean,1829)	<i>Anisodactylus</i> Dejean,1829
* Halobiont. Germany, Netherlands, Belgium, S. England	<i>A. pocilloides</i> (Stephens,1828)
<i>Aepopsis</i> Jeannel,1922	* Halobiont. Sweden, Denmark, Germany, Netherlands, Belgium, England
<i>A. robinii</i> (Laboulbène,1850)	<i>Acupalpus</i> Latreille, <u>in</u> Cuvier,1829
* Maritime. Halobiont. Lives permanently between the tidemarks in crevices & beneath stones. Br. Isles, France, Spain, Morocco	<i>A. (Manicellus</i> V. v. Motschulsky,1864) <i>elegans</i> (Dejean,1829)
<i>Lymaniaeum</i> Stephens,1828	* Halobiont. Denmark, Balticum, Germany, Netherlands, Belgium, England
<i>L. nigropiceum</i> (Marsham,1802)	<i>Nebria</i> Latreille,1802
* At marine shores beneath stones. Germany, Netherlands, S. England	<i>N. complanata</i> (Linnaeus,1767)
<i>Tachys</i> Melsheimer,1806	* Sandy shores Britain, Spain
* 3 British species at marine shores beneath stones.	<i>N. (Nebria) livida</i> (Linnaeus,1758) var. <i>lateralis</i> ?!???
<i>T. (Tachys) scutellaris</i> Germar,1829	German Baltic & North Sea coast
= <i>T. taurica</i> V. v. Motschulsky,1850	<i>Haliplidae</i> Aube,1836 "Vattentrampare"
* Halobiont. German North Sea coast, Netherlands, Belgium, France, England	<i>Haliplus</i> Latreille,1802 (TM <i>Dytiscus impressus</i> Fabricius,1787 = <i>Haliplus</i> (<i>Haliplus</i>) <i>ruficollis</i> (De Geer,1774))
<i>T. (Tachys) parvulus</i> (Dejean,1831)	Subgen.: <i>Haliplus</i> (s. str.)
* Sub-maritime S. England, Germany	<i>H. lineolatus</i> Mannerheim,1844
<i>Trogophloeus</i> Mannerheim,1831	* A typical brackish water species. Limnohalin. N. Baltic
<i>T. schneideri</i> Ganglbauer,18??	<i>H. immaculatus</i> Gerhardt,1877
* Halobiont. Denmark, Borkum, Netherlands	* A typical brackish water species. Pseudohalobiont. Baltic, Norway
<i>T. halophilus</i> von Kiesenwetter,1844	<i>H. apicalis</i> Thomson,1868
* Halobiont. Sweden, Denmark, Germany, Netherlands, France, England	* A typical brackish water species. Pseudohalobiont. Baltic, Britain
<i>Perileptus</i> Schaum,1860	<i>H. fluvialis</i> Aubé,1836, <u>in</u> Dejean & Boisduval German Baltic coast
<i>P. areolatus</i> (Creutz,1799)	Subgen.: <i>Haliplidius</i> Guignot,1928
* Coastal. Sweden, S. England, Germany	<i>H. obliquus</i> (J.C. Fabricius,1787)
<i>Pogonus</i> Dejean,1821, <u>ex</u> Ziegler MS	* A typical brackish water species. Pseudohalobiont. Baltic
<i>P. luridipennis</i> (Germar,1822-25)	<i>H. confinis</i> Stephens,1829 <i>pallens</i> Fowler,1887
* Under pieces of algae cast ashore on salt marsh shores. Halobiont. Bohuslän, Denmark, Germany, Netherlands, France, ?Scotland	* A typical brackish water species. Pseudohalobiont. Baltic
<i>P. chaleucus</i> (Marsham,1802)	Subgen.: <i>Liaphlus</i> Guignot,1928
* Sub-maritime. Halobiont. Denmark, Germany, Netherlands, Belgium, S. England, Scotland, Arcachon	<i>H. fulvus</i> (J.C. Fabricius,1801) <i>lapponum</i> Thomson,1854
<i>P. iridipennis</i> E.A. Nicolai,1822	<i>H. flavicollis</i> Sturm,1834
* Halobiont. Germany	* A typical brackish water species. Pseudohalobiont. Baltic, Norway
<i>P. litoralis</i> (Megerle, <u>in</u> Duftschmid,1812)	Subgen.: <i>Neohaliplus</i> Netolitzky,1911
* Halobiont. Netherlands, Br. Isles	<i>H. lineatocollis</i> (Marsham,1802) German Baltic coast
<i>Panagaeus</i> Latreille,1802	<i>Dytiscidae</i> De Geer,1774 "Dykarbagger"
<i>P. bipustulatus</i> (J.C. Fabricius,1775) "Mindre korslöpare"	<i>Hygrotus</i> Stephens,1828
* Coastal. Skåne-Östergöt., Öland, Gotland, Nericke, S. England	<i>H. (Coelambus</i> Thomson,1860) <i>inaequalis</i> (J.C. Fabricius,1777) * A typical brackish water species. Limnohaline. * Coastal. Baltic, S. England
<i>Chlaenius</i> Dejean,1826, <u>ex</u> Bonelli	<i>Agabus</i> Leach,1817
<i>C. vestitus</i> (G. von Paykull,1790)	<i>A. conspersus</i> (Marsham,1802)
* Coastal. S. England	* Coastal. A typical brackish water species. Pseudohalobiont. Baltic, S. England
<i>Harpalus</i> Latreille,1802	<i>A. nebulosus</i> Forster,1???
<i>H. tenebrosus</i> (Dejean,1821)	* A typical brackish water species. Pseudohalobiont. Baltic, Norway
* Coastal. S. England	<i>A. undulatus</i> Schrank,1???
<i>H. attenuatus</i> Stephens,1828	German Baltic coast
* Coastal. S. England	<i>Ilybius</i> Erichson,1832
<i>H. tardus</i> (Panzer,1796)	<i>I. fuliginosus</i> (J.C. Fabricius,1792)
Sweden, S. England	* A typical brackish water species. Limnohaline. Baltic
<i>Amara</i> Stephens,1827, <u>ex</u> Bonelli,1810	<i>I. subaeneus</i> Erichson,1837
<i>A. ovata</i> (J.C. Fabricius,1792)	* A typical brackish water species. alitic
* Coastal. On gravel. Sweden, S. England	<i>Noterus</i> Schellenberg,1806
<i>A. lucida</i> (Andersch, <u>in</u> Duftschmid,1812)	<i>N. clavicornis</i> (C. De Geer,1774)
* Coastal. S. England	* A typical brackish water species. Baltic, Skåne-Bohuslän, SE. Norway
<i>A. convexiuscula</i> (Marsham,1802)	
* Halophilic. Sweden, Norway, Finland, Denmark, Balticum, North Sea Shores, France, Br. Isles	
<i>Agonum</i> Stephens,1827, <u>ex</u> Bonelli,1810	

<i>Laccophilus</i> Leach,1817	<i>Actinopteryx</i> Mathews,1872
<i>L. variegatus</i> Germar,1???	<i>A. fucicola</i> (Allibert,1844)
* A typical brackish water species	* Halobiont. Beneath seaweed.
Baltic, SE. Norway	Skåne, ?Norway, Denmark, Netherlands, England
<i>Bidessus</i> Sharp,1881	<i>Staphylinidae</i> Latreille,1802 "Kortvingar"
<i>B. geminus</i> (J.C. Fabricius,1798)	<i>Aleocharinae</i>
* A typical brackish water species. Limnohaline.	<i>Actocharis</i> Fauvel,1870
Baltic, Norway	<i>A. marina</i> Fauvel,1870? = <i>A. readingi</i> Shp.,1???
<i>Coelambus</i> Thomson,1860	* Maritime. S. England
<i>C. impressopunctatus</i> Schaller,18??	<i>Phytosus</i> Curtis,1838, ex Rudd MS
* A typical brackish water species. Limnohaline	<i>P. balticus</i> Kraatz,1???
Baltic	* Sub-maritime. Halobiont. Sweden, Denmark, Germany, Netherlands, Belgium, S. England, Scotland
<i>C. parallelogrammus</i> (Ahrens,1812)	<i>P. spinifer</i> G.T. Rudd, in J. Curtis,1838
* A typical brackish water species. Pseudohalobiont. Halophilic.	* Sub-maritime. Halobiont. Denmark, German North Sea coast, Netherlands, France, S. England, Scotland
Bohuslän, Norway, Finland, Denmark, Balticum, North Sea Shores, France, Br. Isles	<i>P. nigriventris</i> (Chevrolat,1843)
<i>Deronectes</i> Sharp,1881	* Sub-maritime. Halobiont. Germany, Netherlands, Belgium, France, S. England
<i>D. depressus</i> (J.C. Fabricius,1794) <i>tatescens</i> Falkenström,1???	<i>Arena</i> Fauvel,1862
* A typical brackish water species. Pseudohalobiont.	<i>A. octavii</i> Fauvel,1862? * Halobiont. Netherlands, S England
Baltic, North Sea coasts	<i>Atheta</i> Thomson,1858
<i>Platambus</i> Thomson,1860	<i>A. thinobiooides</i> (Kraatz,1???)
<i>P. maculatus</i> (Linnaeus,1758)	* Coastal. S. England
Baltic, Skåne-Torne Lappmark	<i>A. (Aloconota</i> Thomson,1858) <i>sulcifrons</i> (Stephens,1829)
POLYPHAGA Emery,1886	S. England
<i>Staphylinidae</i> Latreille,1802	<i>A. (Pseudopasilia</i> Ganglbauer,1895) <i>tabida</i> (Kiesenwetter,18???)
<i>Hydraenidae</i> Mulsant,1844	* Sub-maritime. Halobiont. Germany, S. England, Scotland
<i>Ochthebiinae</i> Thomson,1859	<i>A. (Halobrecta</i> Thomson,1858) <i>flavipes</i> Thomson,18???
<i>Ochthebius</i> Leach,1815	* Sub-maritime. Halobiont. Sweden, Germany, Netherlands, France, S. England, Scotland
<i>O. dilatatus</i> Stephens,1829, ex Leach MS	<i>A. (Halobrecta</i> Thomson,1858) <i>puncticeps</i> Thomson,18???
* Coastal. S. England	* Coastal. Halobiont. Sweden, Norway, Germany, Netherlands, S. England, Scotland
<i>O. metallescens</i> Rosenhauer,1847	<i>A. (Microdota</i> Mulsant & Rey,1873) <i>indubia</i> (Sharp,18???)
* Coastal. S. England	* Coastal. S. England
<i>O. marinus</i> (G. von Paykull,1798?)	<i>A. triangulum</i> (Kraatz,1???)
* A typical brackish water species. Sub-maritime. Halophilic-Halobiont.	* Coastal. ?Halophilic. Sweden, German Baltic, Netherlands, France, Br. Isles
Bohuslän, S. Norway, Finland, Denmark, Balticum, North Sea Shores, France, Br. Isles	<i>A. setigera</i> Sharp.,18???
<i>O. viridis</i> Peyron,1???	* Halophilic. Sweden, Finland, Balticum, Germany, England
* Coastal. Halophilic. S. England, France, Germany	<i>A. marina</i> Rey,18???
<i>O. lejolisi</i> Mulsant & Rey,18???	* Halobiont. Sweden, Finland, Germany, Netherlands, France, England
* Sub-maritime. S. England	<i>A. meridionalis</i> Rey,18???
<i>O. auriculatus</i> Rey,18???	* Halobiont. Sweden, Denmark, Germany, Netherlands, England
* Halobiont. Should the species name be <i>auricularius</i> ?	<i>A. varendorffiana</i> Bernhauer,18???
Denmark, Germany, Netherlands, France, England	* Halobiont. German North Sea coast
<i>O. impressicollis</i> Dejean,1821	<i>A. (Thinobaena</i> Thomson,1859) <i>vestita</i> (Gravenhorst,1???)
* Halophilic. Germany, Netherlands, France, Br. Isles	* Sub-maritime. Halobiont. Sweden, Norway, Denmark, Finland, Germany, Netherlands, S. England, Scotland
<i>O. punctatus</i> Stephens,1829	<i>A. (Acrotona</i> Thomson,1859) <i>fungi</i> (Knob, in Gravenhorst,1806) <i>orbata</i> (Erichson,1837)
* Halophilic. Netherlands, France, England	* Coastal. S. England
Leiodidae	<i>Heterota</i> Mulsant & Rey,1873
<i>Leiodes</i> Latreille,1796	<i>H. plumbea</i> (Waterhouse,1???)
<i>L. ciliaris</i> (W.L.E. Schmidt, in Germar,1841)	* Sub-maritime; under seaweed. Halobiont. Hallands Väderö, Germany, France, S. England
* Halophilic. Sweden, Denmark, Finland, Germany, Netherlands	<i>Myrmecopora</i> Sauley,1865
<i>L. dubia</i> (Kugelann,1794?)	<i>M. lohmanderi</i> Bernhauer,19??
* Sandy coastal habitats. Britain	* Under wrack Bohuslän
<i>Hydnobius</i> Schmidt, in Germar,1841	<i>M. brevipes</i> Butler,18???
<i>H. perrisi</i> (Fairmaire,18???)	<i>M. sulcata</i> (Kiesenwetter,1???)
* Sandy coastal habitats. Britain	* Halobiont. S. England, France
<i>Silphidae</i> "Skinnarbaggar", "Asbaggar"	<i>M. uvida</i> Erichson,18???
<i>Ablattaria</i> Reitter,1885	* Halobiont. Netherlands, Belgium, England
<i>A. laevigata</i> (J.C. Fabricius,1775)	<i>Diglotta</i> Champion,1899
* Coastal. S. England	= <i>Diglossa</i> Haliday,1837, non Wagler,1832
<i>Ptiliidae</i> Erichson 1845/Motschulsky 1845 "Fjädervingar"	* With 2 maritime Scandinavian species.
<i>Ptenidium</i> Erichson,1845	<i>D. meresa</i> (Haliday,1837)
<i>P. punctatum</i> (Gyllenhaal,1827)	* Maritime. Halobiont.
* Sub-maritime. Halobiont. In deep wrack layers. Bohuslän-Uppland, Åland, Finland, Norway, Denmark, Balticum, Germany, Netherlands, Belgium, S. England	Denmark, Germany, Netherlands, France, S. England, Scotland
<i>Acrotrichis</i> Motschulsky,1848	<i>D. marina</i> (Haliday,18???)
<i>A. fratercula</i> Matthews,18???	
* In deep wrack layers. Bohuslän-Uppland, Åland, Finland	
<i>Actidiump</i> Matthews,1868	
<i>A. coarctatum</i> (Haliday,1839?)	
* Halobiont. Sandy shores. Skåne, Denmark, Germany, Netherlands, France, England	

- * on rocky & sandy shores
Britain
- D. submina** Fairmaire,1???
- Arcachon
- Ocalea** Erichson,1837
- O. rivularis** Miller,1???
- S. England
- O. picata** Stephens,18??
- Uppland
- Drusilla** Samouelle,1819, ex Leach MS
- D. canaliculata** Dillwyn,18???
- Br. Isles
- Oxypoda** Mannerheim,1831
- O. tarda** Sharp,1???
- England
- O. exoleta** Erichson,1839
- * Coastal.
S. England
- Aleochara** Gravenhorst,1802
- A. algarum** Fauvel,18???
- * Sub-maritime. Halobiont.
W. Sweden, Norway, Denmark, Germany, Netherlands, Belgium, France, S. England
- A. grisea** Kraatz,1???
- * Sub-maritime. Halobiont.
Sweden, Norway, Finland, Germany, Netherlands, S. England, Scotland
- A. obscurella** Gravenhorst,1806
- * Sub-maritime. Halobiont.
Sweden, Norway, Denmark, Finland, Germany, Netherlands, Belgium, France, S. England, Scotland
- Omalinae**
- Omalium** Gravenhorst,1802
- O. rivulare** (G. von Paykull,1789)
- Sweden, Denmark
- O. laeviusculum** Gyllenhal,1827
- * Sub-maritime. Halophilic.
Bohuslän, Norway, Finland, Germany, Netherlands, S. England, Scotland
- O. riparium** Thomson,18???
- * Sub-maritime. Halobiont.
Skåne, Norway, Denmark, Finland, Germany, Netherlands, Belgium, France, S. England, Scotland
- O. rugulipenne** Rye,18???
- * Halobiont.
Netherlands, France, Br. Isles
- Micralymma** Westwood,1837
- M. marinum** (Ström,1783)
= *M. brevipenne* (Gyllenhal,1810)
* Maritime. Halobiont.
Bohuslän, W. Norway, Denmark, German North Sea coast, Netherlands, Belgium, France, Br. Isles
- Oxytelinae**
- Cryptobium** Mannerheim,1831
- C. fracticornis** (G. von Paykull,1790)
- Sweden, Denmark
- Bledius** Samouelle,1819, ex Leach MS
- B. spectabilis** Kraatz,1???
- * Halobiont.
Norway, Denmark, Germany, Netherlands, Arcachon, S. England, Scotland
- B. arenarius** (G. von Paykull,1800)
- * Halophilic.
Sweden, Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, Arcachon, Br. Isles
- B. tricornis** (Herbst, in Fuessly,1784)
- * Halophilic.
Sweden, Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, France, England
- B. taurus** (Dejean,1821)
= *B. furcatus* (Olivier,1???)
* Halobiont.
Halland, Bohuslän, Denmark, Germany, France, England
- B. diota** ?,18???
- Sweden, Denmark
- B. bicornis** (?L.H. v. Bloch, in Becker,1799?)
- * Halobiont.
Denmark, Germany, Netherlands, England
- B. unicornis** (Dejean, in Germar,1825?)
- * Halobiont.
Germany, Belgium, Arcachon, England
- Oxytelus** Gravenhorst,1802
- O. perrisi** Fauvel,18???
- * Sub-maritime. Halobiont.
Sweden, Norway, Denmark, Germany, Netherlands, Belgium, France, S. England, Scotland
- O. complanatus** Erichson,1839
- * Coastal.
S. England
- Proteininae**
- Proteinus** Latreille,1796
- P. brachypterus** Fabricius,1???
- Sweden, Denmark
- Xantholininae**
- Hypongryus** Tottenham,1940
- * Should it be spelled *Hyponygrus* ?
- H. punctulatus** (G. von Paykull,1789)
- Denmark
- Quediinae**
- Heterothops** Stephens,1829, ex Kirby MS
- H. binotatus** (Gravenhorst,1806?)
- * Sub-maritime. Halobiont.
Sweden, Norway, Denmark, Germany, Netherlands, France, S. England, Scotland
- Steninae**
- Stenus** Latreille,1796
- S. atratulus** Erichson,1839
- * Coastal
Sweden, S. England
- S. nanus** Stephens,1829
- S. England
- Paederinae**
- Medon** Stephens,1833
- M. brunneus** Erichson,18???
- Denmark
- M. ripicola** (Kraatz,1???)
- * Sub-maritime.
S. England
- M. pociferus** Peyron,18???
- * Sub-maritime. Halobiont.
S. England
- Astenus** Dejean,1833
- A. filiformis** (Latreille,18???)
- * Coastal
S. England
- Lathrobium** Gravenhorst,1802
- L. angustatum** Boiduval & Lacordaire,18???
- S. England
- Tachyporinae**
- Tachyporus** Gravenhorst,1802
- T. chrysomelinus** (Linnaeus,1758)
- all of Sweden, S. England
- Staphylininae** Latreille,1802
- Cafius** Stephens,1829, ex Leach MS
- C. xantholoma** (Hellwig, in J.L.C. Gravenhorst,1806)
- * Sub-maritime. Halobiont.
Uppland-Skåne-Bohuslän, Norway, Denmark, Finland, Germany, S. North Sea coast, France, Br. Isles
- C. fucicola** (Curtis,1???)
- * Sub-maritime. Halobiont.
S. England, Scotland
- Philonthus** Stephens,1829
- P. salinus** von Kiesenwetter,1844
- * Halobiont.
Denmark, Germany, Netherlands
- Orthidus** Mulsant & Rey,1876
- O. cribratus** Erichson,1840
- Europe
- Xantholina** Serville,1825
- X. longiventris** Heer,1839
- S. England
- Remus** F. Holme,1838
- R. sericeus** F. Holme,1838
- * Under wrack. Halobiont.
Swedish westcoast, Norway, Denmark, Germany, Netherlands, S. England
- Quedius** Stephens,1829
- Q. tristis** (Gravenhorst,18???)
- S. England
- Q. umbrinus** Erichson,1839
- * Coastal.
Sweden, S. England
- Q. semiaeneus** (Stephens,1829)
- * Coastal.
S. England
- Staphylinus** Linnaeus,1758
- S. ater** Gravenhorst,1802
- * Coastal.
S. England
- S. globulifer** Geoffroy, in A.F. de Fourcroy,1785
- S. England
- Ocypus** Samouelle,1819
- O. oleans** Müller,1???
- Amrum
- ?Subfamilies?

- Olophrum** Erichson,1839
- O. piceum** (Gyllenhal,1810)
Sweden, S. England
- O. consimile** (Gyllenhal,1810)
* Halophilic.
Sweden, Norway, Germany
- Trogophloeus** Mannerheim,1831
- T. halophilus** H. v. Kiesenwetter,1844
* Maritime.
S. England
- T. unicolor** Sharp,1???
= *T. anglicanus* Sharp,1???
* Sub-maritime.
S. England
- Pselaphidae** "Klubbhornsbaggar"
- Brachyluta** Thomson,1859
- B. simplex** (Waterhouse,1???)
* Sub-maritime.
S. England
- B. helferi** Schm.-Göb.,18??
* Halobiont.
Sweden, Norway, Denmark, Germany, Netherlands, England
- Reichenbachia** Leach,1826
- R. waterhousi** Rye,18??
* Halobiont.
England
- Hydrophiloidae** De Geer,1774
- Hydrophilidae** De Geer,1774 "Vattenbaggar", "Palpbaggar"
- Sphaeridiinae** Latreille,1802
- Cercyon** Leach,1817
- C. littoralis** (Gyllenhal,1808)
* Sub-maritime. Halobiont. In deep wrack layers.
Bohuslän-Uppland, Åland, Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, France, Britain
- C. depressus** Stephens,1829
* Sub-maritime. Halobiont. In deep wrack layers.
Bohuslän-Uppland, Finland, Norway, Denmark, Germany, Netherlands, S. England, Scotland
- C. marinus** Thomson,1???
* Littoral.
Bohuslän
- Hydrophilinae** De Geer,1774
- Hydrophilini** De Geer,1774
- Enochrus** Thomson,1859
- E. bicolor** (J.C. Fabricius,1792)
* A typical brackish water species. Halophilic-Halobiont.
Sweden, Norway, Finland, Denmark, Balticum, North Sea Shores, France, Br. Isles
- E. melanoccephalus** (Olivier,1792)
* A typical brackish water species. Pseudohalobiont.
Baltic- Bohuslän, S. Norway
- E. testaceus** (J.C. Fabricius,17??)
* A typical brackish water species. Limnohaline.
Baltic
- E. quadripunctatus** (Herbst,1797) *fuscipennis* Thomson,18??
* A typical brackish water species. ?Pseudohalobiont.
Baltic, Norway
- Berosini Mulsant,1844
- Berosus** Leach,1817
- B. spinosus** Stephens,18??
* A typical brackish water species. Halobiont.
Sweden (Skåne-Gotland), S. Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, Br. Isles
- Oocylini Hansen,1991
- Laccobius** Erichson,1837
- L. decoratus** (Gyllenhal,1810)
* A typical brackish water species. Halobiont.
Baltic
- L. minutus** (Linnaeus,1758)
* A typical brackish water species. Limnohaline.
Baltic
- Elaphorus** J.C. Fabricius,1775
= *Helophorus* Illiger,1801
- E. minutus** J.C. Fabricius,1775
* A typical brackish water species.
Baltic
- Anacaenini Hansen,1991
- Paracymus** Thomson,1867
- P. aeneus** Germar,18??
* Brackish water species. Halobiont.
Skåne, Germany, Netherlands, France, England
- Histeridae** "Stumpbaggar", "Pansarbaggar"
- Halacritus** Schmidt,1893
- H. punctum** (Aubé,1843)
- = *Acritus* Leconte,1853 *punctum* (Aubé,1843)
* Sub-maritime. Halobiont.
German North Sea coast, Netherlands, S. England
- Baeckmanniolas** Reichardt,1926
- B. maritimus** (Stephens,1829, ex Kirby MS)
= *Pachylopus* Erichson,1834 *maritimus* (Stephens,1829, ex Kirby MS)
* Sub-maritime. Halobiont.
Denmark, Germany, Netherlands, France, S. England, Scotland
- Scarabaeoidea**
- Scarabaeidae** "Bladhorningar"
* 7 coastal species occurs in Britain
- Aphodius** Illiger,1798
- A. nitidulus** (J.C. Fabricius,1792)
* Coastal.
S. England
- Psammobius** Heer,1841
- P. porcicollis** (Illiger,1803)
* Coastal.
S. England
- Aegialia** Latreille,1807
- A. arenaria** (J.C. Fabricius,1787)
* Sub-maritime. Halobiont.
Sweden, Norway, ?Finland, Denmark, Balticum, Germany, North Sea Shores, France, Br. Isles
- Byrrhoidea** Latreille,1804
= *Dryopoidea*
- Heteroceridae** MacLeay,1825 "Strandgrävbaggar"
- Heterocerinae** MacLeay,1825
- Heterocerus** Fabricius,1792
- H. flexuosus** Stephens,1828
* Halobiont.
S. Sweden, Norway, Denmark, Balticum, Germany, Netherlands, Arcachon, Br. Isles
- H. (Littorimus** Gozis,1885) *maritimus* Guérin-Menéville,1844
* Halobiont.
Netherlands, Arcachon
- H. obsoletus** Curtis,1828
* Halophilic.
Sweden, Denmark, Germany, Belgium, Br. Isles
- Elateroidea** Leach,1815
- Elateridae** Leach,1815
= *Dicronychidae*
= *Lissomidae*
= *Plastoceridae* "Knäppare"
- Agriotes** Eschscholtz, in Thon,1829
- A. sordidus** (Illiger,1807)
* Sandy coasts.
Britain
- Throscidae**
= *Trixagidae*
- Trixagus** Kugelann,1794
- T. elateroides** (Heer,1841)
* Salt-marshes.
Britain
- Cantharoidea**
- Cantharidae** "Flugbaggar"
- Cantharis** Linnaeus,1758
- C. rufa** Linnaeus,1758
* Beneath strandline algae
Sweden, Britain
- C. (Telephorus** Schaeffer,1766) *darwiniana* Crotch,18??
* Halobiont.
Germany, Belgium, Scotland
- Paratinus** Abeille de Perrin,1891
- P. femoralis** (Erichson,18??)
* Halophilic.
Sweden, Denmark, Finland, Germany
- Bostrichoidea**
- Ptinidae** "Tjuvbaggar"
- Ptinus** Linnaeus,1767
- P. palliatus** (Perris,18??)
Britain
- Cleroidea** Latreille,1802
- Melyridae** Leach,1815
- Malachius** Fabricius,1775
- M. sardous** Erichson,1840
* Halobiont.
Germany, Netherlands
- M. vulneratus** (Abeille,18??)
* Salt-marshes
Britain
- Dolichosoma** Stephens,1830

<i>D. lineare</i> (Rossi,17??) * Salt-marshes Britain	<i>A. humilis</i> Germar,1825 * Halobiont. Sweden, Norway, Denmark, Germany, Netherlands, England
<u>Cucujoidea</u> = <u>Clavicornia</u>	<i>A. crotchi</i> Pic.,19?? * Halobiont. England, France
<u>Corylophidae</u> = <u>Orthoperidae</u>	<i>A. bimaculatus</i> (Illiger,1801) * Halophilic. Sweden, Finland, Denmark, Balticum, Germany, Netherlands, France
<i>Corylophus</i> Leach, in Stephens,1833	<i>A. flavigipes</i> (Panzer,17??) * Halophilic. Sweden, Finland, Denmark, Balticum, Germany, Netherlands, France
<i>C. cassidoides</i> (Marsham,1802)? * "Only British species occurring on the coast" according to Hayward & Ryland (Eds.),1990. Synon. of next?	<i>A. antherinus</i> (Linnaeus,1761) * In wrack. Sweden
Britain	<u>Oedemeridae</u> "Blombaggar"
<i>C. sublaevipennis</i> Du Val,18?? * Coastal. S. England	<i>Nacerdes</i> Dejean,1834, ex Steven MS = <i>Nacerda</i> Faldermann,1837
<u>Cryptophagidae</u> "Fuktbaggar"	<i>N. melanura</i> (Linnaeus,1758)? * Coastal. Sweden, S. England
<i>Atomaria</i> Stephens,1829, ex Kirby MS	<u>Mordellidae</u> "Tornbaggar"
<i>A. fuscipes</i> (Gyllenhal,1808) Sweden, Britain	<i>Mordellistena</i> A. Costa, in O. Costa,1854
<u>Nitidulidae</u> "Glansbaggar"	<i>M. parvula</i> (Gyllenhal,18??) <i>inaequalis</i> Mulsant,1???
<i>Meligethes</i> Stephens,1829, ex Kirby MS	* Coastal. Sweden
<i>M. exilis</i> Sturm,1845 * Coastal. S. England	<u>Chrysomeloidea</u> Latreille,1802
<u>Lathridiidae</u> "Mögelbaggar"	<i>Chrysomelidae</i> Latreille,1802 "Bladbaggar"
<i>Corticaria</i> Marsham,1802 * 4 coastal British species	<i>Macrolea</i> Samouelle,1819, ex Hoffmannsegg MS = <i>Haemona</i> Dahl,1823, ex Megerle MS
<i>C. crenulata</i> (Gyllenhal,18??) * Coastal. W. Sweden, S. England	<i>M. mutica</i> Fabricius,1??? * Lives on <i>Ruppia</i> , <i>Zostera</i> etc.. A typical brackish (mesohaline) water species. Halobiont. Uppland-Bohuslän, Finland, Danish estuaries, Balticum, Germany, North Sea Shores, France, Br. Isles
<i>C. impressa</i> (Olivier,1790) * Coastal. Uppland, S. England	<i>M. pubipennis</i> Reuter,1??? * A typical brackish water species. Halobiont. Baltic
<u>Coccinellidae</u> "Nyckelpigor"	<i>Chrysolina</i> Motschulsky, in Schrenck,1860
<i>Subcoccinella</i> Guérin-Méneville,1842, ex Huber	<i>C. haemoptera</i> (Linnaeus,1758) * on sandy coasts. S. England
<i>S. vigintiquatuorpunctata</i> (Linnaeus,1758) "Härig (24-prickig) nyckelpiga" * Coastal. Skåne-Gästrikland, S. England	<i>C. banksi</i> (J.C. Fabricius,1775) * Coastal. S. England
<i>Micraspis</i> Dejean,1835, ex Chevrolat MS	<u>Psylliodes</u> Berthold, in Latreille,1827
<i>M. sexdecimpunctata</i> (Linnaeus,1758) = <i>Typhaspis</i> Crotch,1874 <i>sexdecimpunctata</i> (Linnaeus,1758) * Coastal. S. England	<i>P. marcida</i> (Illiger,1807) * Coastal, Halobiont. Sweden, Denmark, Finland, Balticum, Germany, Netherlands, Belgium, France, S. England
<i>Coccinella</i> Linnaeus,1758	<i>P. conannus</i> (Stephens,18??) * Salt-marshes. Britain
<i>C. undecimpunctata</i> Linnaeus,1758 * Halophilic. Sweden, Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, France, Br. Isles	<i>Crepidodera</i> Dejean,1835, ex Chevrolat MS
<i>C. septempunctata</i> Linnaeus,1758 "7-prickig nyckelpiga", "Jungfru Maria Nyckelpiga" * Ubiquitous & often shore occurring Sweden, Britain	<i>C. impressa</i> (Fabricius,17???) * Salt-marshes. Britain
<u>Phalacridae</u>	<i>Cassidinae</i> "Sköldbaggar"
<i>Phalacrus</i> G. von Paykull,1800	<i>Cassida</i> Linnaeus,1758
<i>P. brunnipes</i> (Brisout,18??) * Coastal & in banks of rivers Britain	<i>C. nobilis</i> Linnaeus,1758 * Coastal. Sweden, S. England
<u>Tenebrionoidea</u> Latreille,1802	<i>C. vittata</i> C. de Villers,1789 * Salt-marshes. S. England
<i>Tenebrionidae</i> Latreille,1802 "Svarbbaggar"	<u>Curculionoidea</u> Latreille,1802
<i>Phaleria</i> Latreille,1802	<u>Curculionidae</u> Latreille,1802 "Vivlar"
<i>P. cadaverina</i> (J.C. Fabricius,1792) * Sub-maritime. Halobiont. Sweden, Denmark, Germany, Netherlands, Belgium, France, S. England	<i>Apion</i> Herbst, in Jablonsky,1797 "Spetsvivlar"
<i>Melanimon</i> Steven,1829	<i>A. atomarium</i> Kirby,1808 * Coastal. S. England
<i>M. tibialis</i> (Fabricius,1781?) S. England	<i>A. confluens</i> Kirby,18?? * Coastal. S. England
<i>Opatrium</i> J.C. Fabricius,1775	<i>A. hookeri</i> Kirby,1808 * Coastal. S. England
<i>O. sabulosum</i> (Linnaeus,1758) * Coastal. Skåne-Västmanland, S. England	<i>A. schoenherri</i> Boheman,18?? * Coastal. S. England
<i>Phylan</i> Dejean,1821	<i>A. laevicolle</i> Kirby,1808 * Coastal. S. England
<i>P. gibbus</i> (J.C. Fabricius,1775) * Coastal. Halobiont. Sweden, Norway, Denmark, Finland, Germany, Netherlands, Belgium, France, S. England, Scotland	<i>A. ononicola</i> Bach,18?? * Coastal. S. England
<i>Cylindrinotus</i> Faldermann,1837	
<i>C. pallidus</i> (Curtis,18??) Britain	
<u>Anthicidae</u> Latreille,1802 "Snabbaggar"	
<i>Anthicus</i> G. von Paykull,1798	
<i>A. angustatus</i> Curtis,1838, ?non Dejean,1821 * Sub-maritime. Halobiont. S. England, France	

- A. limonii* Kirby,1808
 * Halobiont.
 France, ?Belgium
- Otiorhynchus* Germar,1824 "Öronvivlar"
- O. rugifrons* (Gyllenhaal,1813)
 * Coastal.
 Sweden, S. England
- O. atroapterus* (C. de Geer,1775)
 * Coastal. Halobiont.
 Sweden, Denmark, Germany, Netherlands, Belgium, S. England, Scotland
- O. ligneus* (Olivier,1807)
 * Halophilic.
 Sweden, Denmark, Germany
- Bagous* Germar,1817
- B. argillaceus* Gyllenhaal, in Shoeherr,1836
 * Halophilic.
 Sweden, Denmark, Germany, England
- Cathormiocerus* Shoeherr,1843
- C. attaphilus* Brisout,18??
 * Coastal.
 S. England
- C. myrmecophilus* (Seidlitz,18??)
 * Coastal.
 S. England
- Polydrusus* Germar,1817 "Glansvivlar"
 = *Polydrosus* Shoeherr,1826
- P. chrysomela* (Olivier,1807)
 * Sub-maritime.
 S. England
- Philopedon* Stephens,1831, non Stephens,1829
 =? *Cneorhinus* Shoeherr,1823
- P. plagiatus* (Schaller,18??)
 * Coastal.
 S. England
- Sitona* Germar,1817 "Ärtvivlar"
- S. waterhousei* J. Walton,1846, ex Shoeherr MS
 * Coastal.
 S. England
- Phloeophagia* Aurivillius,1924
- P. spadix* (Herbst, in Jablonsky,1795)
 * Sub-maritime.
 S. England
- Smicronyx* Shoeherr,1843
- S. jungermanniae* (Reich,1797)
 * Coastal.
 S. England
- Tychius* Germar,1817
- T. schneideri* (Herbst, in Jablonsky,1795)
 * Coastal.
 S. England
- Limnobaris* Bedel,1885
- L. talbum* (Linnaeus,1758)
 * Sub-maritime.
 S. England
- Ceuthorhynchidius* J. Du Val,1854
- C. dawsoni* Brisout,18??
 * Coastal. Halobiont.
 S. England, ?France, ?Belgium
- Ceutorhynchus* Germar,1824, ex Schueppel MS
 = *Ceutorhynchus* Gemminger & Harold,1871
- C. terminatus* (Herbst, in Jablonsky,1795)
 * Coastal.
 S. England
- Sirocalus* Heyden, in Heyden, Reitter & Weise,1906
- S. cecilis* Hansen,18??
 * Halobiont.
 Denmark, Germany, Netherlands
- Mecinus* Germar,1821
- M. circulatus* (Marsham,1802)
 * Coastal
 S. England