

South Scandinavian Marine Protoctista

Provisional Check-list compiled at the Tjärnö Marine Biological Laboratory by:

TJARNOLAB. / Hans G. Hansson - 1991-07 - 1997-04-02

Species found in South Scandinavia, as well as from neighbouring areas, chiefly the British Isles, have been considered, as some of them may show have a slightly more northern distribution, than what is known today. However, species with a typical Lusatian distribution, with their northern distribution limit around France or Southern British Isles, have as a rule been omitted here, albeit a few species with probable northern limits around the British Isles are listed here until distribution patterns are better known. The compiler would be very grateful for every correction of presumptive lapses and omissions an initiated reader could make.

Denotations: (TM) = Genotype @ = Associated to * = General note

P.S. This list is a very unfinished manuscript. Chiefly flagellated organisms have yet been considered. This provisional PDF-file is so far only published as an Intranet file within TMBL:s domain.

PROTOCTISTA

= PROTISTA

PROTOZOA Goldfuss,1818 (Emend. von Siebold,1846)

Gregarella Chatton & Brachon,1936

* Is this a much regressed flagellate?

G. fabreum Chatton & Brachon,1936

@ on surface of or within food vacuoles of the ciliate *Fabrea salina* Henneguy
?France

Abollifera Vørs,1992

* Taxon incertae sedis

A. prolabilens Vørs,1992 (TM - Tvarminne)

Tvarminne, Denmark

Allantion Sandon,1924

* Taxon incertae sedis

A. tachyploosa Sandon,1924

Tvarminne, Denmark, Greenland, European & Arctic fresh waters

Ancyromonas Kent,1880

* Taxon incertae sedis

A. sigmoides Kent,1880

= *Bodo cephaloporus* Larsen & Patterson,1990

Tvarminne, Denmark, Jersey, England, Greenland, midAtlantic, Arctic Canada, Fiji, Hawaii, Panama, Australia, Antarctica

Bodomorpha Hollande,1952 (TM *Pseudobodo minima* Hollande,1942)

* Taxon incertae sedis.

Bodopsis Lemmermann,1913 (TM *Dimorpha alternans* Klebs,1892)

* Taxon incertae sedis.

B. gobboldi Lackey,1933

E North America

Caecitellus Patterson, Nygaard, Steinberg & Turley,1993

* Of uncertain affinities.

C. parvulus Griessmann,1913 Patterson, Nygaard, Steinberg & Turley,1993

= *Bodo parvulus* Griessmann,1913 (TM)

Oslofjord, midAtlantic

Chlorarachnion Geitler,1930

* Marine? Taxon incertae sedis. Genus transferred to new Divisio: Chlorarachniophyta and new Classis: Chlorarachniophycace by Hibberd & Norris,1984

Cladomonas von Stein,1878

* Marine? Taxon incertae sedis.

Colpodella Cienkowsky,1865

* Probably related to *Perkinsus* & Apicomplexa, according to Patterson,1989

C. perforans (Hollande,1938) Patterson & Zölfel,1992

= *Spiromonas perforans* (Hollande,1938)

= *Bodo perforans* Hollande,1938

@ *Chilomonas paramecium*

?

C. gonderi (Foisner & Foisner,1984) Patterson & Zölfel,1992

= *Spiromonas gonderi* Foisner & Foisner,1984

@ invading pellicle of ciliate genus *Colpoda*

?

Colponema von Stein,1878

* Taxon incertae sedis.

?C. sp. : Paperna & Laurencin,1979

@ *Dicentrarchus labrax*

French mariculture

Cruzella de Faria, da Cunha & Pinto,1922

* Taxon incertae sedis.

Cryothecomonas Thomsen, Buck, Bolt & Garrison,1991 (TM *C. armigera* Thomsen, Buck, Bolt &

Garrison,1991 - Antarctic)

* Taxon incertae sedis. 3 antarctic species

C. inermis Thomsen, Buck, Bolt & Garrison,1991

?Kattegatt

Cryptaulax Skuja,1948 (TM *C. akpos* Skuja,1948 - fresh water)

= *Spiromonas* Skuja,1939, non Perty,1852

* Taxon incertae sedis. May belong among Kinetoplastida

C. marina Thronsdælen,1969

* May be identical with the freshwater living species *C. thiophila* Skuja,1956

Gulf of Finland, Tvarminne, Oslofjord, S & W Norway

Cyclomonas Fromentel,1874

= *Astasiopsis* Bütschli,1885 (TM *Cyclidium distortum* Dujardin,1841)

* Marine? Belongs possibly among euglenids

C. distortum (Dujardin,1841) Fromentel,1874

= *Cyclidium distortum* Dujardin,1841 (TM)

= *Astasiopsis distortum* (Dujardin,1841) Bütschli,1885

?

Dingensia Patterson & Zölfel,1992, in Patterson & Larsen (TM *Heteromita angusta* Dujardin,1841)

* Taxon incertae sedis. Very similar to *Cryptaulax* Skuja

Dinomonas Kent,1880

* Taxon incertae sedis.

Diphylliea Massart,1920

* Marine? Incertae sedis.

Diplocalium Grassé & De Landre,1952 (TM *Bicosoeca inopinatum* ??,???)

* Marine? Incertae sedis.

Diplomita Fromentel,1874 (TM *Diplomita insignis* Fromentel,1874)

* Marine? Incertae sedis.

Diplonema Griessmann,1913, non Berndt,1845 (Diptera), nec Greene,1857 (Coel.)

= *Isonema* ??,???, non Meek & Worthen,1865 (Mollusca), nec Maas,1909 (Coel.)

= *Flagellomonas* Skvortzow,19?? = *Lackeymonas* Skvortzow,19??

= *Lowymonas* Skvortzow,19?? = *Milaneciamonas* Skvortzow,19??

= *Spina* Skvortzow,19?? = *Teixeiramonas* Skvortzow,19??

= *Kolbeana* Skvortzow,19??

* Genus incertae sedis. Ultrastructural affinities with euglenids.

D. ambulator Larsen & Patterson,1990

W Scotland

Discocelis Vørs,1988 (TM - Denmark)

Tvarminne, Denmark, Panama, Brazil, Australia

D. saleuta Vørs,1988 (TM - Denmark)

Tvarminne, Denmark, Panama, Brazil, Australia

Entomosigma Schiller,1925

* Marine? Genus incertae sedis.

Errera Schouteden,1907

* Marine? Genus incertae sedis.

Fromentella Patterson & Zölfel,1992 (TM *Pleuromonas granulosa* Fromentel,1874)

* Marine? Genus incertae sedis.

Gymnophrys Cienkowski,1876

* Marine? Genus incertae sedis.

Heliobodo Valkanov,1928

* Marine? Genus incertae sedis.

Helkesimastis Woodcock & Lapage,1915

* Taxon incertae sedis

H. faecicola Woodcock & Lapage,1915?

Plymouth

Hemistasia Griessmann,1913

* Genus incertae sedis.

Histonia Voigt,1902

* Marine? Genus incertae sedis.

Hyperamoeba Aléxieff,1924, non Goodkov & al.,1982

* Marine? Genus incertae sedis.

Kamera (Woodcock,1916) Patterson & Zölfel, in Patterson & Larsen,1992 (TM *Heteromastix lens* Woodcock,1916?)

* Marine? Genus incertae sedis.

Kathablepharis Skuja,1939

K. ovalis Skuja,1948

* Fresh water genus incertae sedis.

Gulf of Finland

Kittokskia Vørs,1992

* Taxon incertae sedis

K. ystava Vørs,1992 (TM - Tvarminne)

Tvarminne, Denmark, Arctic Canada

Luffisphaera Belcher & Swale,1975

* Taxon incertae sedis

L. turiformis Vørs,1992

Denmark

L. cucumiformis Belcher & Swale,1975

Tvarminne, English fresh water

L. myosurus Belcher & Swale,1975

Tvarminne, English fresh water

L. longihastis Vørs,1992

?Denmark

L. "sp. B" Belcher & Swale,1975

Tvarminne, English fresh water

L. "striata" Thomsen (nomen nudum) : Vørs,1992

Denmark

Metopion Larsen & Patterson,1990

* Taxon incertae sedis

M. fluens Larsen & Patterson,1990

Tvarminne, Denmark, Canaries, Fiji, Brazil

Metromonas Larsen & Patterson,1990

* Taxon incertae sedis. *Clastrivira parva* Schouteden,1907 may be the gliding stage of one species.

M. simplex (Griessmann,1913) Larsen & Patterson,1990

= *Phylomonas simplex* Griessmann,1913 (TM)

Tvarminne, Denmark, Wadden Sea, Germany, France, England, Ireland, Greenland, Arctic Canada, Brazil, Fiji, Hawaii, Australia, Antarctica

M. grandis Larsen & Patterson,1991

N midAtlantic

Micromonas Borrel,1902, non Manton & Parke,1960

* Marine? Genus incertae sedis.

- Ministeria** Patterson, Nygaard, Steinberg & Turley,1993
* Genus incertae sedis.
- M. marisola** Patterson, Nygaard, Steinberg & Turley,1993
N midAtlantic
- Monochrysis** Skuja,1948
* Marine? Genus incertae sedis.
- Multicilia** Cienkowsky,1881
?= *Polymastix* Gruber,1884
* Genus incertae sedis.
- M. marina** ??,???
- Nephromonas** Droop,1953
* Genus incertae sedis.
- N. hyalina** Droop,1953 (TM - Finland)
* Recorded from low salinity pools.
Finland
- Parabodo** Skuja,1939
* Marine? Genus incertae sedis.
- Paramastix** Skuja,1948
* Marine? Genus incertae sedis.
- Petasaria** Moestrup,1979
* Genus incertae sedis.
- P. heterolepis** Moestrup,1979 (TM)
Denmark
- Phylloimitus** von Stein,1878
* Genus incertae sedis.
- P. granulatus** Larsen & Pattersen,1990
Denmark
- Phylomonas** Klebs,1892
* Marine? Genus incertae sedis.
- Pirsonia** Schneppf, Drebes & Elbrächter,1990
* Genus incertae sedis.
- P. guinardiae** Schneppf, Drebes & Elbrächter,1990 (TM - SE North Sea)
@ penetrating the valvae & girdle region of the pelagic diatom *Guinardia flaccida* (Castr.) Peragallo (main host) & *Rhizosolenia delicatula*
SE North Sea
- P. verrucosa** Kühn, in Kühn, Drebes & Schneppf,1996
@ phagotrophic on *Rhizosolenia delicatula*
German Bight
- P. formosa** Kühn, in Kühn, Drebes & Schneppf,1996
@ phagotrophic on *Rhizosolenia setigera* (main host), *R. schrubsolaei*, *R. pungens*, *R. stolterfothii*, *Guinardia flaccida*, *Thalassionema nitschiae* & *Stephanopyxis turris*
German Bight
- P. diadema** Kühn, in Kühn, Drebes & Schneppf,1996
@ phagotrophic on *Coscinodiscus granii*, *C. wailesii* (main hosts) & *C. concinnus*
German Bight
- P. eucampiae** Kühn & Schneppf, in Kühn, Drebes & Schneppf,1996
@ phagotrophic on *Eucampia zodiacus* (main host), *Cerataulina pelagica* & *Leptocylindrus danicus*
German Bight
- P. mucosa** Drebes, in Kühn, Drebes & Schneppf,1996
@ phagotrophic on *Rhizosolenia schrubsolaei* (main host), *R. delicatula*, *R. setigera*, *Cerataulina pelagica*, *Eucampia zodiacus* & *Chaetoceras decipiens*
German Bight
- Platytilomonas** Larsen & Patterson,1990
* Marine? Genus incertae sedis.
- Pleurostomum** Namyslowsky,1913
* Marine? Genus incertae sedis.
- Quadriciilia** Vørs,1992
* Taxon incertae sedis
- Q. rotundata** (Skuja,1948) Vørs,1992 (TM)
= *Tetramitus rotundatus* Skuja,1948
= *Collodictyon sphaericum* Norris,1964
Tvärrimme, Denmark, New Zealand, Antarctica, Swedish fresh water
- Rhizomonas** Kent,1880 (TM *R. verrucosa* Kent,1880 - ?fresh water)
= *Solenicola* Pavillard,1916
* Belonging where?
- R. setigera** (Pavillard,1916) Patterson, Nygaard, Steinberg & Turley,1993
= *Solenicola setigera* Pavillard,1916 (TM of *Solenicola* Pavillard,1916)
@ covering the diatom *Dactyliosolen mediterraneus* (H. Perag.) H. Perag.
Mediterranean, N midAtlantic
- Rigidomastix** Aléxéeff,1929
* Marine? Genus incertae sedis.
- Salpingorhiza** Klug,1936
* Marine? Genus incertae sedis.
- Schewiakoffia** Corliss,1960
= *Maupaxia* Schewiakoff,1893, non Viguer,1886 (Polychaeta)
* Marine? Genus incertae sedis.
- Stenocodon** Pascher,1942
* Marine? Genus incertae sedis.
- Stephanomonas** Kent,1881
* Marine? Genus incertae sedis.
- Stomatochone** Pascher,1942
* Marine? Genus incertae sedis.
- Streptomonas** Klebs,1892
* Marine? Genus incertae sedis.
- Telonema** Griessmann,1913
* Systematic position problematic, but an affinity to *Cyathomonas* may possibly be assumed. Not mentioned in Chrétiennet-Dinet,1990
- T. subtle** Griessmann,1913 (TM)
Tvärrimme, Denmark, Oslofjord, S Norway, Bergen, Tromsö, Greenland, Arctic Canada, France, North Atlantic, Mediterranean, Belize, Japan
- T. sp.** : Vørs,1992
* Will be described by Thomsen as *T. antarctica* (nom. nudum)
Tvärrimme, Denmark, Antarctic
- Tetracilia** Valkanov,1970
* Brackish water. Genus incertae sedis.
- Toshiba** Patterson & Zöllfler,1991 (TM *Dimastigamoeba vorax* Massart,1920)
= *Dimastigamoeba* Massart,1920, non Blochmann,1894
* Marine? Genus incertae sedis.
- Trichonema** Fromentel,1872
* Marine? Genus incertae sedis.
- Trimastix** Kent,1880
* Genus incertae sedis.
- T. ciona** (Parona Corrado,1886) Grassé,1952
= *Elvirea ciona* Parona Corrado,1886
@ *Ciona intestinalis*
Mediterranean
- Apusomonadidae** Karpov & Mylnikov,1989
* Taxon incertae sedis
- Apusomonas** Aléxéeff,1929
Amastigomonas de Saedeleer,1931
= *Thecamonas* Larsen & Patterson,1990
- A. debruynei** de Saedeleer,1931
= *Thecamonas trahens* Larsen & Patterson,1990
Tvärrimme, S Baltic, temperate & tropical Atlantic, Greenland, Arctic Canada
- A. mutabilis** (Griessmann,1913) Molina & Nerad,1991
= *Rhynchomonas mutabilis* Griessmann,1913
= *Thecamonas mutabilis* (Griessmann,1913) Larsen & Patterson,1990
Tvärrimme, Denmark, Wadden Sea, England, France, Greenland, midAtlantic, Brazil, Australia
- A. bermudensis** Molina & Nerad,1991
Bermudas
- Jacobidae** Patterson,1990 (ICZN) = *Jacobaceae* Patterson,1990 (ICBN)
* Family incertae sedis.
- Jakoba** Patterson,1990
* Genus incertae sedis.
- J. libera** (Ruin,1938) Patterson,1990
= *Cryptobia libera* Ruin,1938 (TM)
Kattegatt, SE North Atlantic, N midAtlantic, India, Australia
- Proteomyxidae**
* Marine? Taxon incertae sedis. Correct taxon niveau? Polyphyletic group.
@ Usually phyco- or phytoparasitic.
- Amoeboaphelidium** Scherffel,1925
* Marine?
- Amylophagus** Scherffel,1925
* Marine?
- Aphelidiopsis** Scherffel,1925
* Marine?
- Barbeta** Dangeard,1886
* Marine?
- Cibdelia** Juel,1925
* Marine?
- Diplophysalis** Zopf,1885
* Marine?
- Ectobiella** de Bruyne,1890
* Marine?
- Endospora** Scherffel,1925
* Marine?
- Endobella** de Bruyne,1888
* Marine?
- Endomonas** Zopf,1887
* Marine?
- Gymnococcus** Zopf,1885
* Marine?
- Polysporella** Zopf,1887
* Marine?
- Protomonas** Zopf,1885 (TM *Monas amyli* Cienkowsky,1865)
= *Quasimonas* Gobi,1915 (TM *Monas amyli* Cienkowsky,1865)
* Marine?
- Pseudospora** Cienkowsky,1865
* Marine?
- Pseudosporopsis** Scherffel,1925
* Marine?
- Woronina** Cornu,1872
* Marine?
- Pseudodendromonadidae**
* Marine? Taxon incertae sedis. Correct taxon niveau?
- Cyathobodo** Petersen & Hansen,1961
* Marine?
- Pseudodendromonas** Bourrelly,1953
* Marine?
- Thaumatomastigidae** Patterson & Zöllfler,1991 = *Thaumatomastigaceae*
* Taxon incertae sedis. Correct taxon niveau?
- Gyromitus** Skuja,1939 (TM *G. disomatus* Skuja,19???)
* Marine?

- Hyaloselene** Skuja,1956
* Marine?
- Protaspis** Skuja,1939
- P. glans** Skuja,1939
Tåvrinne, Latvia, Denmark, Wadden Sea, Arctic Canada, Hawaii
- P. simplex** Vørs,1992
Tåvrinne, Denmark, Arctic Canada
- P. verrucosa** Larsen & Patterson,1990
Tåvrinne, Denmark, England, Arctic Canada, Fiji
- P. gemmifera** Larsen & Patterson,1990
N midAtlantic
- Rhizaspis** Skuja,1948
* Marine?
- Synoikomonas** Skuja,1969
= *Gomesiamonas* Skvortzow & Noda,1969 = *Guttula* Skvortzow & Noda,1969
= *Hortobagiomonas* Skvortzow & Noda,1969 = *Oyemonas* Skvortzow & Noda,1969
= *Refractocystis* Skvortzow & Noda,1969 = *Refractodes* Skvortzow & Noda,1969
= *Refractomonas* Skvortzow & Noda,1969 = *Stromia* Skvortzow & Noda,1969
= *Synoikomonas* Skvortzow & Noda,1969
* Marine?
- Thaumatomonas** de Saedeleer,1931
* Marine?
- Thaumatomastix** Lauterborn,1899
= *Thaumatomema* Lauterborn,1896 (homonymity with a fossil diatom)
?= *Reckeritia* Conrad,1920 (may be a synonym of *Thaumatomonas* de Saedeleer instead)
?= *Tentaculomonas* Skvortzow & Noda,1969
- T. setifera** Lauterborn,1899 (TM)
Plymouth
- T. salina** (Birch-Andersen,1973) Beech & Moestrup,1986
= *Chrysosphaerella salina* Birch-Andersen,1973
Tåvrinne, Denmark, a widespread euryhaline & eurythermal species
- T. "punctata"** Thomsen, unpublished (nomen nudum)
Tåvrinne, S Baltic
- T. triangulata** (Balanov,1980) Beech & Moestrup,1986
= *Chrysosphaerella triangulata* Balanov,1980
Tåvrinne, Denmark, Russia
- T. sagittifera** (Conrad,1920) Beech & Moestrup,1986
= *Reckeritia sagittifera* Conrad,1920
Denmark
- Katablepharidae**
* Belonging in which higher taxon?
- Katablepharidae**
- Leucocryptos** Butcher,1967
* Colourless. Taxon incertae sedis. Closely related to freshwater genus *Katablepharis* Skuja,1939, both previously classified in Cryptophyceae.
- L. marina** (Braarud,1935) Butcher,1967 (TM)
= *Bodo marina* Braarud,1935
= *Chimonomas marina* (Braarud,1935) Halldal,1953
Tåvrinne, Belt Sea, Skagerrak, Oslofjord, S & W Norway, North Sea, Plymouth, Arctic Seas, N midAtlantic, cosmopolitan
- L. remigera** Vørs,1992
Tåvrinne, Denmark, Faeroes, Greenland, Arctic Canada
- SPONGOMONADIDA** Hibberd,1983
* Marine? Taxon incertae sedis.
- Rhipidodendron** von Stein,1878
* Marine?
- Spongomonas** von Stein,1878 (TM) *Phalansterium intestinalum* Cienkowsky,1870
* Marine?
- SARCOMASTIGOPHORA** Honigberg & Balamuth,1963
- MASTIGOPHORA** Diesing,1866
- PHYTOMASTIGOPHORA** Calkins,1909
- CRYPTOMONADIDA Senn,1900 "Rekylalger"
= CRYPTOMONADES Engler,1903 = CRYPTOPHYCEAE Pascher,1914
All Pankow,1990 & Parke & Green, in Parke & Dixon,1976 included.
- CRYPTOMONADINA Senn,1900
= CRYPTOMONADES Engler,1903
- Goniomonadidae** Hill,1992 (ICZN)
= Goniomonadaceae Hill,1992 (ICBN) = *Cyathomonadaceae* sensu Pringsheim,1944
- Goniomonas** von Stein,1878
= *Cyathomonas* Flamentel,1874 (p.p.)
* Colourless.
Tåvrinne, Denmark, Belize, European fresh water
- G. truncata** (Fresenius,1858) von Stein,1878 (TM)
= *Monas truncata* Fresenius,1858
= *Cyathomonas truncata* (Fresenius,1858) Fish,1885
* sand- & mud-living. Usually a fresh water species.
Tåvrinne, Denmark, Belize, European fresh water
- G. amphinema** Larsen & Patterson,1990
Tåvrinne, Denmark, England, midAtlantic, Australia, Fiji, Panama, Belize
- G. pacifica** Larsen & Patterson,1990
Tåvrinne, Denmark, Wadden Sea, England, midAtlantic, Belize, Panama, Brazil, Hawaii
- Cryptomonadaceae** Ehrenberg,1838, sensu Butcher,1967
= Cryptomonadidae
- Chilomonas** Ehrenberg,1832
* Colourless.
- C. paramecium** Ehrenberg,1832 (TM)
brackish water from many localities, e.g. Britain, but normally freshwater
- Chroomonas** Hansgirg,1885 (TM) *C. nordstedtii* Hansgirg,1885 - fresh water)
* Some of the species of subgenus *Cryptochrysis* should possibly go to *Pyrenomonas*.
- C. (Chroomonas) raphanoidea** Conrad & Kufferath,1954
Bad description.
Belgium
- C. (Chroomonas) baltica** (Büttner,1911) N. Carter,1938
= *Cyanomonas baltica* Büttner,1911
Kieler Bucht, Denmark, Belgium, Isle of Wight
- C. (Chroomonas) vectensis** N. Carter,1937
S Baltic, Kattegatt, Vadehavet, Belgium, Isle of Wight
- C. (Chroomonas) extensa** Butcher,1967
Isle of Wight
- C. (Chroomonas) monococca** Butcher,1967
Essex salt-marshes
- C. (Chroomonas) virescens** (Butcher,1952) Butcher,1967
= *Cryptochrysis virescens* Butcher,1952
E Britain
- C. (Chroomonas) diplococca** Butcher,1967
Yorkshire, Isle of Wight
- C. (Chroomonas) dispersa** Butcher,1967
Severn river salt-marsh lagoons
- C. (Chroomonas) plurococca** Butcher,1967
Essex, Kent & Isle of Wight salt-marshes
- C. (Chroomonas) mesostigmatica** Butcher,1967
Essex, Guernsey
- C. (Chroomonas) collegioides** Butcher,1967
S England pools & salt-marshes
- C. (Chroomonas) placoidea** Butcher,1967
Oslofjord, Whitby rockpool
- C. (Cryptochrysis** Pascher,1911) *nana* Butcher,1967
Puffin Island rockpool
- C. (Cryptochrysis) marina** (Büttner,1911) Butcher,1967
= *Cryptomonas marina* Büttner,1911
Kieler Bucht
- C. (Cryptochrysis) esopus** (Conrad & Kufferath,1954) Butcher,1967
= *Cryptomonas esopus* Conrad & Kufferath,1954
* Inadequate description.
Belgium
- C. (Cryptochrysis) pseudocaudata** Conrad & Kufferath,1954 Butcher,1967
= *Cryptomonas pseudocaudata* Conrad & Kufferath,1954
Belgium
- C. guttala** Conrad
Parke & Dixon,1976 area
- C. africana** Meyer & Pienaar,19??
?
- C. sp.** : Edler, Hällfors & Niemi,1984
Gulf of Finland
- Cryptomonas** Ehrenberg,1832, emend. von Stein,1878
* Probably confined to fresh water only. Marine species to be transferred to other genera.
- C. (Cryptomonas) curvata** (Ehrenberg,1832) Penard,1921
= *C. ovata* var. *curvata* Ehrenberg,1832
?
- C. (Cryptomonas) diplastida** Butcher,1967
Engl. Channel
- C. (Cryptomonas) erosa** Ehrenberg,1832, emend. von Stein,1878
* Fresh & brackish water
Gulf of Finland
- C. (Cryptomonas) irregularis** Butcher,1967
Engl. Channel
- C. (Cryptomonas) tilloensis** Conrad & Kufferath,1954
Belgium
- C. (Cryptomonas) marssonii** Skuja,1948
= *C. erosa* Ehrenberg var. *reflexa* Marsson,1904
Latvia, S Baltic, Belgium
- C. (Cryptomonas) ovata** Ehrenberg,1832, emend. von Stein,1878 (TM)
* Fresh water & brackish water ditches etc.
Gulf of Finland, S Baltic, Oslofjord, Britain (almost cosmopolitan)
- C. (Cryptomonas) profunda** Butcher,1967
Oslofjord, Thames estuary
- C. (Cryptomonas) rhynchophora** (Conrad,1939) Butcher,1967
= *Rhodomonas rhynchophora* Conrad,1939
Belgium
- C. (Cryptomonas) stigmatica** Wislouch,1924
Oslofjord, Engl. Channel, Black Sea
- C. (Cryptomonas) suberosa** Butcher,1967
Yorkshire
- C. (Cearulomonas) magnifica** Butcher,1967
Isle of Wight saltmarsh
- C. (Cearulomonas) cyanomagna** Butcher,1967
Cornwall rockpools
- C. (Cearulomonas) synechia** (Skuja,1939) Butcher,1967
= *Chroomonas synechia* Skuja,1939
Riga Bay, Belgium
- C. torta** Conrad & Kufferath,1954
* Species of uncertain relationship, sensu Butcher,1967
Belgium

- C. calciformis* Lucas,1970
Parke & Dixon,1976 area
- C. rostellata* Lucas,1970
Parke & Dixon,1976 area
- Komma* Hill,1991
- Rhinomonas* Hill & Wetherbee,1988
- R. pauca* Hill & Wetherbee,1988?
- R. fulva* (Butcher) Hill & Wetherbee,1988?
= *Chroomonas (Cryptochrysis* Pascher,1911) *fulva* (Butcher,1952) Butcher,1967
= *Cryptochrysis fulva* Butcher,1952
Wales
- R. lateralis* (Butcher) Hill & Wetherbee,1988?
= *Chroomonas (Cryptochrysis* Pascher,1911) *lateralis* (Butcher,1952) Butcher,1967
= *Cryptochrysis lateralis* Butcher,1952
- R. fragarioides* (Butcher) Hill & Wetherbee,1988?
= *Chroomonas (Cryptochrysis* Pascher,1911) *fragarioides* Butcher,1967
Engl. Channel
- R. reticulata* (Lucas,1970) Novarino,19??
= *Cryptomonas reticulata* Lucas,1970
Parke & Dixon,1976 area
- Rhodomonas* Karsten,1898, emend. Hill & Wetherbee,1989
= *Pyrenomonas* Santore,1984
= *Pyrenomonas* Leedale & Hibberd,in Lee, Hutner & Bovee,1985
- R. baltica* Karsten,1898 (?)
= *Cryptomonas baltica* (Karsten,1898) Butcher,1967
= *Chroomonas pseudobaltica* Butcher,1967
Gulf of Finland, Belt Sea - Kattegatt area, Bohuslän, Oslofjord, Bergen, Tromsö, Isle of Wight, Yorkshire, Wales
- R. marina* (Dangeard,1892) Lemmermann,19??
= *Cryptomonas marina* Dangeard,1892
= *Chroomonas marina* (Büttner) Butcher,19??
= *Rhodomonas pelagica* Lohmann,1908
= *Cryptomonas pelagica* (Lohmann,1908) Butcher,1967
W Baltic, Kattegatt, E North Sea
- R. salina* (Wislouch,1924) Hill & Wetherbee,1989
= *Cryptomonas salina* Wislouch,1924 (?) of *Pyrenomonas* Leedale & Hibberd,in Lee, Hutner & Bovee,1985
= *Pyrenomonas* (Wislouch,1924) Leedale & Hibberd,in Lee, Hutner & Bovee,1985
= *Chroomonas (Cryptochrysis* Pascher,1911) *salina* (Wislouch,1924) Butcher,1967
* With great form variability.
Kattegatt, Tromsö, all around Britain, Mediterranean
- R. lacustris* var. *lacustris* Pascher & Ruttner, in Pascher,1913
= *Cryptomonas pusilla* Bachmann,1923
= *Rhodomonas minuta* Skuja,1948
= *Rhodomonas pusilla* (Bachmann,1923) Javornicky,1967
* Should probably be removed from *Rhodomonas*, according to Hill & Wetherbee,1989
All Baltic, Denmark
- R. lacustris* var. *nannoplantica* (Skuja,1948) Javornicky,1976
= *Rhodomonas minuta* var. *nannoplantica* Skuja,1948
S Baltic
- R. ruttneri* Schiller,1925
* Species of uncertain relationship, sensu Butcher,1967
?
- R. gracilis* Schiller,1925
* Species of uncertain relationship, sensu Butcher,1967
Belgium, Adriatic
- R. abbreviata* Butcher, ex Hill & Wetherbee,1989?
= *Cryptomonas (Cryptomonas) abbreviata* Butcher,1967
Guernsey rockpool
- R. atrorosea* Butcher, ex Hill & Wetherbee,1989?
= *Chroomonas (Cryptochrysis* Pascher,1911) *atrorosea* Butcher,1967
Sussex, Isle of Wight, Bristol Channel
- R. chrysoidae* Butcher, ex Hill & Wetherbee,1989?
= *Cryptomonas (Cryptomonas) chrysoidae* Butcher,1967
Essex
- R. duplex* Hill & Wetherbee,1989?
= *Cryptomonas (Cryptomonas) appendiculata* Butcher,1967
Isle of Man
- R. falcata* Butcher, ex Hill & Wetherbee,1989?
= *Chroomonas (Cryptochrysis* Pascher,1911) *falcata* Butcher,1967
Wales
- R. heteromorpha* Butcher, ex Hill & Wetherbee,1989?
= *Chroomonas (Cryptochrysis* Pascher,1911) *heteromorpha* Butcher,1967
Britain
- R. maculata* Butcher, ex Hill & Wetherbee,1989?
= *Cryptomonas (Cryptomonas) maculata* Butcher,1967
Engl. Channel
- Geminigera* Hill,1991
- G. cryophila* (Taylor & Lee) Hill,1991?
?
- Proteomonas* Hill & Wetherbee,1986
- P. sulcata* Hill & Wetherbee,1986?
?
- Teleaulax* Hill,1991
- T. acuta* (Butcher,1952) Hill,1991
= *Cryptomonas acuta* Butcher,1952 (?)
= *Cryptomonas prora* Conrad & Kufferath,1954
Kattegatt, Oslofjord, Bergen, Belgium, England, Wales
- T. amphioxidea* (Conrad,1939) Hill,1991
= *Rhodomonas amphioxidea* Conrad,1939
= *Chroomonas amphioxidea* (Conrad,1939) Butcher,1967
Gulf of Finland, Kattegatt, Belgium, E North America, Japan
- Plagioselmis* Butcher,1967
- P. prolonga* Butcher,1967 (?)
Kattegatt, Bergen, Suffolk
- P. punctata* Butcher,1967
England (St. Germans River)
- Storeatula* Hill,1991
- S. major* Butcher, ex Hill,1991?
= *Cryptomonas (Cryptomonas) major* Butcher,1967
Essex, Isle of Wight
- Ioselmis* Butcher,1967
- L. obconica* Butcher,1967 (?)
Bergen, Plymouth
- Falcomonas* Hill,1990
- F. daucoides* (Conrad & Kufferath) Hill,1990?
= *Hillea daucoides* (Conrad & Kufferath,1954) Butcher,1967
= *Chroomonas daucoides* Conrad & Kufferath,1954
Belgium
- Hemiselmidaeae* Butcher,1967 ex Silva,1980
- Hemiselmis* Parke,1949
= *Sennia* : Skuja,1948, non Pascher
= *Nephroselmis* : Pascher,1912, non von Stein,18
=? *Protocryphsis* Pascher,1911 (p.p.)
- H. (Plagiomonas* Butcher,1967) *anomala* Butcher,1967
Oslofjord, Wales
- H. (Hemiselmis) brunnescens* Butcher,1967
Oslofjord, Engl. Channal, Wales
- H. (Hemiselmis) cyclocea* Butcher,1967
Oslofjord, England, Wales
- H. (Plagiomonas* Butcher,1967) *simplex* Butcher,1967
Oslofjord, Bergen, England, Wales
- H. (Plagiomonas* Butcher,1967) *virescens* Droop,1955
Øregrend, S Kattegatt, Oslofjord, Bergen, Tromsö, Yorkshire, Scotland, Japan, Australia
- H. (Hemiselmis) rufescens* Parke,1949 (?)
North Sea, Isle of Man
- H. (Hemiselmis) amylifera* Butcher,1967
Suffolk
- H. (Hemiselmis) oculata* Butcher,1967
Engl. Channel
- H. (Plagiomonas* Butcher,1967) *rotunda* Butcher,1967
Isle of Wight
- H. vinososa* (Conrad,1939) Chrétiennet-Dinet-1990
= *Protochrysis vinososa* Conrad,1939
Belgium
- Hilleaceae* Butcher,1967
- Hillea* Schiller,1925, emend. Butcher,1967
- H. fusiformis* (Schiller,1913) Schiller,1925 (?)
= *Chlamydomonas fusiformis* Schiller,1913
Adriatic
- H. caerulea* (Schiller,1925) Butcher,1967
= *Rhodomonas caerulea* Schiller,1925
* Possibly identical with *H. fusiformis* (Schiller).
?
- H. phaselos* (Conrad & Kufferath,1954) Butcher,1967
= *Chroomonas phaselos* Conrad & Kufferath,1954
Belgium
- H. cyanea* (Lackey,1932) Butcher,1967
= *Chroomonas cyanea* Lackey,1932
Belgium, New Jersey
- H. minuscula* (Conrad & Kufferath,1954) Butcher,1967
= *Rhodomonas minuscula* Conrad & Kufferath,1954
Belgium
- H. fusulina* (Conrad & Kufferath,1954) Butcher,1967
= *Rhodomonas fusulina* Conrad & Kufferath,1954
Belgium
- H. serpens* (Conrad & Kufferath,1954) Butcher,1967
= *Cryptomonas serpens* Conrad & Kufferath,1954
Belgium
- H. semilunaris* (Conrad & Kufferath,1954) Butcher,1967
= *Cryptomonas semilunaris* Conrad & Kufferath,1954
Belgian brackish water
- H. akrobelos* (Conrad & Kufferath,1954) Butcher,1967
= *Cryptomonas akrobelos* Conrad & Kufferath,1954
Belgian brackish water
- H. marina* Butcher,1962
Wales
?
- DINOFLAGELLIDA* Bütschli,1885
= *DINOPHYCEAE* West & Fritsch,1927
- Karyoclastis* Campbell,1926
* Correct order? Which family? Related to *Duboscquella* ?
- K. tintinni* Campbell,1926
@ *Tintinnopsis nucula*
?California
- DESMOMONADALES* Pascher,1914
- Adinomonadaceae* Schiller,19??
- Adinomonas* Schiller,1928

- * Regarded as a doubtful dinoflagellate by Sournia,1986.
* Not mentioned by Larsen & Sournia,1992, in Patterson & Larsen
- Haplodiiniaceae** Lindemann,1928 = Desmomonadaceae
- Desmopastix**
* Not mentioned by Larsen & Sournia,1992, in Patterson & Larsen
- Pleromonas** Pascher,1914 (TM *P. erosa* Pascher,1914)
- Haplodinium** Klebs,1912 (TM *H. anjoliense* Klebs,1912)
- Desmocapsaceae** Pascher,1914
- Desmocapsa** Pascher,1914
@ Rhizophyceae epiphyte
- D. gelatinosa** Pascher,1914 (TM)
?
- OXYRRHINALES Sournia,1984
* Taxon not recognized by Taylor,1987.
- Oxymniaeae** Sournia,1984
- Oxrrhis** Dujardin,1841
- O. marina** Dujardin,1841 (TM)
= *O. tentaculifera* Conrad,1939
= *O. maritima* Van Meel,1969
* Main occurrence in rock pools.
Gulf of Finland, S & W Baltic, Oresund, Kattegatt, Oslofjord, Bergen, Tromsö, Britain, (almost universal distribution)
- PROROCENTRALES Lemmermann,1910
- Prorocentridae** = Prorocentraceae von Stein,1883
- Prorocentrum** Ehrenberg,1834
= *Cercaria* Michaelis,1830
= *Exuviaella* Cienkowski,1881
= *Postprorocentrum* Gourret,1883
- P. micans** Ehrenberg,1834 (TM)
= *P. schilleri* Bohm, in Schiller,1933
= *P. levartinoides* Bures,1959
= *P. pacificum* Wood,1963
cosmopolitan neritic, W & S Baltic, Kattegatt, Skagerrak, Oslofjord, North Sea, Britain
- P. balticum** (Lohmann,1908) Loeblich III,1970
= *Exuviaella baltica* Lohmann,1908
= *Prorocentrum pomoideum* Bursa,1959
= *Exuviaella aequatorialis* Hasle,1960
all Baltic but Bothnian Bay, Kattegatt, Skagerrak, Oslofjord, North Sea, Plymouth
- P. aporum** (Schiller,1918) Dodge,1975
= *Exuviaella apora* Schiller,1918
= *Exuviaella granii* Gaarder,1938
= *Exuviaella antarctica* Hada,1970
Norway, North Sea, Adriatic
- P. cassubicum** (Woloszynska,1928) Dodge,1975
= *Exuviaella cassubica* Woloszynska,1928
* Litoral
central Baltic?, Isle of Wight, Woods Hole
- P. lima** (Ehrenberg,1859) Dodge,1975
= *Cryptomonas lima* Ehrenberg,1859
= *Exuviaella marina* Cienkowski,1881
= *Dinophysix laevis* von Stein,1883
= *Exuviaella cincta* Schiller,1918
= *Exuviaella osterfeldii* Schiller,1933
= *Exuviaella caspica* Kiselev,1940
* Litoral & benthic. Toxic to mammals.
Baltic, Oslofjord, Trondheimsfjord, North Sea, Britain, except NE England, Europe
- P. nanum** Schiller,1918
= *Exuviaella pusilla* (Schiller) Schiller,1928
off Plymouth
Atlantic, Adriatic
- P. compressum** (Bailey,1850) Abe,1967
= *Pyxidicula compressa* Bailey,1850
= *Prorocentrum bidentatum* Schiller,1928
= *Prorocentrum lebouriae* Schiller,1928
= *Exuviaella oblonga* Schiller,1928
= *Exuviaella tenuicula* Matzenauer,1933
= *Exuviaella elongata* Rampi,1951
Limfjorden, Kattegatt, North Sea, off N & NW Britain, ?eastward Gulf stream drift, Mediterranean
- P. triestinum** Schiller,1918
= *P. redfieldii* Bursa,1959
= *P. pyrenoideum* Bursa,1959
Limfjorden, Netherlands, Plymouth, Mediterranean
- P. gracile** Schutt,1895
= *P. macrurus* Anthanassopoulos,1931
= *P. hentscheli* Schiller,1933
= *P. sigmoides* Bohm,1933
= *P. diamantinae* Wood,1963
E Scotland, Atlantic, Mediterranean
- P. scutellatum** Schröder,1900
= *P. sphaeroideum* Schiller,1928
= *P. robustum* Tafall,1942
N Kattegatt, Skagerrak, N & off SW Scotland, Atlantic, Arctic Sea
- P. dentatum** von Stein,1883
= *P. obusidens* Schiller,1928
= *P. veloi* Tafall,1942
= *P. monacense* Kufferath,1957
North Sea, Plymouth, Galway
- P. minimum** (Pavillard,1916) Schiller,1933
= *Exuviaella minima* Pavillard,1916
= *Prorocentrum triangulatum* Martin,1929
= *Exuviaella mariae-lebouriae* Parke & Ballantine,1957
= *Prorocentrum cordiformis* Bursa,1959
* Certain clones toxic to mice.
W & S Baltic, Öresund, S & central Kattegatt, S North Sea, Sussex, N Scotland, Portugal, Mediterranean
- Mesoporus** Lillick,1937
= *Porella* Schiller,1928
- = *Dinoporella* Halim,1919
= *Porothece* Silva,1960
- M. perforatus** (Gran,1915) Lillick,1937 (TM)
= *Exuviaella perforata* Gran,1915
= *Porella perforata* : Schiller,1928
= *Porella asymmetrica* Schiller,1933
= *Porella adriatica* Schiller,1928
= *Porella bisimpresa* Schiller,1928
= *Porella globulus* Schiller,1928
SW & N Kattegatt, Skagerrak, Oslofjord, off Norway, North Sea, Britain, Adriatic
- DINOPHYSIALES Lindemann,1928
- Citharistaceae**
- Citharistes** von Stein,1883 (TM *C. regius* von Stein,1883)
- Dinophysiidae** Bergh,1882 = *Dinophysiaceae* Bütschli,1885
- Dinophysis** Ehrenberg,1839
= *Phalacroma* von Stein,1883 = *Pseudopalacroma* Jörgensen,1923
= *Prodinophysis* Balech,1944 = *Metadinophysis* ??,??
= *Pseudopalacroma* Jörgensen, ex Lebour,19??
- D. acuta** Ehrenberg,1839 (TM)
* DSP-producing, toxic to mammals.
worldwide distribution, W & S Baltic, Öresund, Kattegatt, Skagerrak, Oslofjord, North Sea, N Scotland
- D. acuminata** Claparède & Lachmann,1859
= *D. borealis* Paulsen,1949
= *D. lachmanni* Paulsen,1949
= *D. boehni* Paulsen,1949
* DSP-producing, toxic to mammals.
all Baltic, Öresund, Kattegatt, Skagerrak, Oslofjord, North Sea, around Britain
- D. arctica** Mereschkowsky,1879
Belt Sea
- D. baltica** (Paulsen,1908) Kofoid & Skogsberg,1928
= *D. ovum* var. *baltica* Paulsen,1908
= *D. cassubica* Woloszynska,1928
All Baltic except Bothnian Sea & Bay
- D. caudata** Saville-Kent,1881
= *D. tripos* Gurrell,1883
= *D. homunculus* von Stein,1883
SE Kattegatt, off N Scotland, off SW England
- D. contracta** (Kofoid & Skogsberg,1928) Dodge,1982
= *Phalacroma contractum* Kofoid & Skogsberg,1928
=? *Phalacroma ruudi* Braarud,1935
Norway, Oslofjord, North Sea, off N Iceland, E England, NW Spain
- D. dens** Pavillard,1915
* Probably a development stage of *D. acuta* Ehrenberg
Kattegatt, off Norway, off W Britain, Danish Strait, Mediterranean
- D. groenlandica** (Schiller,1933) Balech,1967
= *Phalacroma groenlandica* Schiller,1937
Baltic
- D. hastata** von Stein,1883
= *D. odiosa* ?,??
W Baltic, Öresund, Kattegatt, Oslofjord, Norway, North Sea, off N Scotland, NE & SW England, off NW Spain
- D. mucronata** (Kofoid & Skogsberg,1928) Sournia,1973
= *Phalacroma mucronata* Kofoid & Skogsberg,1928
= *Phalacroma paulseni* Schiller,1928
W Norway, off NW Britain
- D. nasutum** (von Stein,1883) Parke & Dixon,1968
= *Phalacroma nasutum* von Stein,1883
off NE & SW England, off Bretagne, Mediterranean
- D. norvegica** Claparède & Lachmann,1858/59
= *D. norvegica* var. *debilior* Paulsen,1907
* DSP-producing, toxic to mammals.
all Baltic, Öresund, Kattegatt, Skagerrak, Oslofjord, North Sea, N Scotland
- D. ovum** Schütt,1895
Baltic, N Scotland
- D. punctata** Jörgensen,1923
Plymouth, W France
- D. recurva** Kofoid & Skogsberg,1928
= *D. lenticula* Pavillard,1916, non of Daday,1888
Engl Channel & other areas around Britain, Mediterranean
- = *D. rotundata* Claparède & Lachmann,1859
= *D. whittingae* Balech,1971
= *Phalacroma rotundatum* (Claparède & Lachmann,1859) Kofoid & Michener,19??
all Baltic, Öresund, Kattegatt, Skagerrak, Oslofjord, North Sea, Norway, Greenland, Barents Sea, around Britain, Mediterranean
- D. sacculus** ??,??
* DSP-producing, toxic to mammals.
France
- D. skagii** Poulsen,1949
N Scotland
- Histioneis** von Stein,1883 (TM *H. remora* von Stein,1883)
= *Parahistioneis* Kofoid & Skogsberg,1928
- Ornithocercus** von Stein,1883 (TM *O. magnificus* von Stein,1883)
- Dinofurcula** Kofoid & Skogsberg,1928 (TM *D. ultima* (Kofoid) Kofoid & Skogsberg,1928)
- Histiophysis** Kofoid & Skogsberg,1928
* Regarded as a doubtful dinoflagellate by Sournia,1986.
* Not mentioned by Larsen & Sournia,1992, in Patterson & Larsen
- Heteroschisma** Kofoid & Skogsberg,1928 (TM *H. inaequale* Kofoid & Skogsberg,1928)
= *Latifascia* Loeblich Jr. & Loeblich III
= *Proheteroschisma* Tai & Skogsberg,19
- Metaphalacroma** Tai & Skogsberg,1934 (TM *M. skogsbergii* Tai, in Tai & Skogsberg,1934)
- Sinophysis** Nie & Wang,1944 (TM *S. microcephalus* Nie & Wang,1944)
- S. ebriolum** (Herdman,1924) Balech,1956

- * sand-living
Kattegatt, Isle of Man, Engl. Channel, Port Erin, Woods Hole
- Thaumatodinium*** Böhm,1933
* Regarded as a doubtful dinoflagellate by Sournia,1986.
* Not mentioned by Larsen & Sournia,1992, in Patterson & Larsen
- Amphisolenia*** von Stein,1883
- A. globifera*** von Stein,1883 (TM)
N Biscay, warm Atlantic, Mediterranean
- Triposolenia*** Kofoid,1906 (TM *T. truncata* Kofoid,1906)
- Oxyphysidae***
Oxyphytis Kofoid,1926 (TM *O. oxytoxoides* Kofoid,1926)
- ACTINISCALES**
- Actiniscaceae*** Kützing,1844
- Actiniscus*** Ehrenberg,1843
= *Diaster* Meunier,18
= *Gymnaster* Schütt,18
- A. pentasterias*** (Ehrenberg) Ehrenberg,1854 (TM)
N. Kattegatt, Skagerrak, Oslofjord
- Dicroerisma*** Taylor & Cattell,1969 (TM *D. psilonereiella* Taylor & Cattell,1969)
- Achradiana*** Lohmann,1903
- Amphitholus*** ??,1???
- DINOTHRICHALES** Pascher,1914
- Dinotrichidae*** = ***Dinothrichaceae*** Pascher,1914
- Dinothrix*** Pascher,1914
@ Marine plants
- D. paradoxa*** Pascher,1914 (TM)
?Mediterranean or ?North Sea
- Dinochlonium*** ??,1???
- DINAMOEBALES**
- Dinamoebaceae***
- Dinamoebidium*** Pascher,1916
- DINOCOCCALES** Pascher,1914
= **PHYTODINIALES**
- Collinella***?Cachon,1964?
* Correct suborder? Which family?
* Not mentioned by Larsen & Sournia,1992, in Patterson & Larsen
- C. ovoidea***?Cachon,1964?
@ *Acanthomitra pellucida*
Mediterranean
- Gloeodiniidae***
= ***Gloeodiniaceae*** Pascher, ex Schiller,1937
- Gloeodinium*** Klebs,1912 (TM *G. montanum* Klebs,1912 - fresh water)
=? *Hemidinium*
- Rufusiella*** Loeblich III,1967?
* Possibly a synonym of *Gloeodinium* Klebs, sensu Larsen & Sournia,1992
- Phytodiniaceae***
- Cystodinedria*** Pascher,1944
- Cystodinium*** ??,1???
- Dinastridium*** ??,1???
- = *Bourrallyella* ??,1???
- Hypnodinium*** ??,1???
- Manchudinium*** ??,1???
- * Not mentioned by Larsen & Sournia,1992, in Patterson & Larsen
- Paulsenella*** Chatton,1920
* Not mentioned by Larsen & Sournia,1992, in Patterson & Larsen
- P. chaetoceratis*** (Paulsen,1911)
@ *Chaetoceros decipiens* & *C. boreale*
E North Sea, E Greenland
- P. vonstoschii*** Drebes & Schnepf,1989
@ *Sreptothecca thamesis*, *Neostreptothecca subindica*, *Bellerochea malleus* & *Lithodesmium undulatum*
North Sea
- P. kornmanni*** Drebes & Schnepf,1989
@ *Eucampia zodiacus*
North Sea
- Phytodinedria*** ??,1????
* Not mentioned by Larsen & Sournia,1992, in Patterson & Larsen
- Phytodinium*** ??,1???
- Rhizodinium*** ??,1???
- * Not mentioned by Larsen & Sournia,1992, in Patterson & Larsen
- Spiniferodinium*** ??,1????
- Stylocodiaceae***
- Stylocodium*** Klebs,1912 (TM *S. globosum* Klebs,1912 - fresh water)
= *Dinopodiella* Pascher,19??
* only one marine species known
- S. gastrophilum*** Cachon,19??
@ in the cavities of the gastrozooids of the Siphonophores *Abylopsis tetragona*, *Chelophyses* sp. & *Sulculeolaria* sp.
Mediterranean
- Tetradinium*** ??,1????
- Dinococcus*** ??,1????
- Thoracosphaeraceae**
- Thoracosphaera*** Kamptner,1927 (TM *T. pelagica* Kamptner,1927)
- PYROCYSTALES**
- Pyrocystidae***
= ***Pyrocystaceae*** (Schütt) Lemmermann,1899
- Pyrocystis*** Murray, ex Haeckel,1890
= *Muracystis* Haeckel,1891
@ Marine invertebrate ectoparasites
- P. lunula*** (Schütt,1895) Schütt,1896
= *Gymnodinium lunula* Schütt,1895
S & W Norway
- P. noctiluca*** Murray, ex Haeckel,1890 (TM)
= *P. pseudonoctiluca* Wyville-Thomson, in Murray
= *Goniadoma concava* Gaarder,1954
oceanic warm water species occasionally found to the W of Britain
- P. hamulus*** Cleve,1900
Engl. Channel, Clyde Sea
- BRACHYDINIALES** Loeblich III, ex Sournia,1984
- Brachydiniaceae*** Sournia,1972
- Asterodinium*** Sournia,1972 (TM *A. gracile* Sournia,1972)
- Brachydinium*** Taylor,1963 (TM *B. capitatum* Taylor,1963)
- NOCTILUCALES** Haeckel,1894
- Kofoididiniaceae*** Taylor,1976
- Cymbodinium*** Cachon & Cachon,1967 (TM *C. elegans* Cachon & Cachon,1967 - Mediterranean)
- Kofoidinium*** Pavillard,1928
- K. yelleloides*** Pavillard,1928 (TM)
Britain north of 54° N
- Pomatodinium*** Cachon & Cachon-Enjumet,1966 (TM *P. impatiens* Cachon & Cachon-Enjumet,1966 - Mediterranean)
- Spatulodinium*** Cachon & Cachon,1968
- S. pseudonoctiluca*** (Pouchet,1885) Cachon & Cachon, ex Loeblich Jr. & Loeblich III,1969 (TM)
= *Gymnodinium pseudonoctiluca* Pouchet,1885
around NW England clockwise to NE England, Faeroes, Bretagne, Mediterranean
- Leptodiscaceae*** Kofoid,1905
- Cachonodinium*** Loeblich III,1980 (TM *C. caudatum* (Cachon & Cachon,1969) Loeblich III,1980 - Mediterranean)
= *Leptodinium* Cachon & Cachon,1969?
- Craspedotella*** Kofoid,1905 (TM *C. pileolus* Kofoid,1905 - Mediterranean & Tropical Pacific)
- Leptodiscus*** Hertwig,1877 (TM *L. medusoides* Hertwig,1877)
= *Pratjellia* Lohmann,19??
- Leptophyllus*** Cachon & Cachon-Enjumet,1964 (TM *L. dasypus* Cachon & Cachon-Enjumet,1964 - Mediterranean)
= *Abedinium* Loeblich Jr. & Loeblich III,19??
- Petalodinium*** Cachon & Cachon,1969 (TM *P. porcelio* Cachon & Cachon,1969 - Mediterranean)
- Scaphodinium*** Margalef,1963 (TM *S. mirabile* Margalef,1963 - Mediterranean)
= *Leptospathium* Cachon & Cachon-Enjumet,1964
- Noctilicidae** = **Noctilicaceae** Kent,1881
- Noctiluca*** Suriray, ex Lamarck,1816
- N. scintillans*** (Macartney,1810) Kofoid,1920
= *Medusa marina* Slabber,1771
= *Medusa scintillans* Macartney,1810
= *Noctiluca militaris* Suriray, ex Lamarck,1816 (TM)
* Toxic to fish.
W Baltic, Øresund, Kattegatt, Bohusl. Oslofjord, E North Sea, S & W Norway
- Noctilucales incertae sedis**
- Pronoctiluca*** Fabre-Domergue,1889
= *Pelagorhynchus* Pavillard,1917
- P. pelagica*** Fabre-Domergue,1889 (TM)
= *Pelagorhynchus marina* Pavillard,1917
= *Rhynchomonas marina* Lohmann,1902
= *Protodinium marinum* Kofoid & Swezy,1921
W Baltic, Isle of Man, Bretagne, N mid-Atlantic
- P. phaeocysticola*** ??,1???
- @ endoparasite in *Phaeocystis pouchetii*; ectoparasite on *Thalassiosira partheneia*?
P. sp.: Drebes,1974
@ *Coscinodiscus concinnus* & other centric diatoms
North Sea
- P. sp. cf. *phaeocysticola*
@ *Biddulphia sinensis*?
- Protodinifer*** Kofoid & Swezy,1921
* Correct order? Not mentioned by Larsen & Sournia,1992, in Patterson & Larsen
- GYMNODINIALES** Lemmermann,1910
- Gymnodiniidae*** Bergh,18?? = ***Gymnodiniaceae*** Lankester,1885
- Amphidinium*** Claparède & Lachmann,1859
= *Trochodinium* Conrad,19??
- A. operculatum*** Claparède & Lachmann,1859 (TM)
=? *A. steinii* (Lemmermann,1910) Kofoid & Swezy,1921

- = *A. klebsii* Kofoid & Swezy, 1921
 = *A. massartii* Biecheler, 1952
 = *A. rhynchocephalum* Anissimowa, 1926
 = *A. wistlouchii* Hulbert, 1957
 = *A. hofferti* Schiller & Diskus, 1955
 all Baltic, Skagerrak, North Sea, Norway, Belgium, Britain, Mediterranean
- A. acutissimum*** Schiller, 1933
 Kattegatt
- A. britannicum*** (Herdman, 1924) Lebour, 1925
 = *A. assymetricum* var. *britannicum* Herdman, 1924
 * sand-living.
 E North Sea, fr. E Scotland clockwise around Britain to SW Scotland
- A. britannicum*** (Herdman) Lebour, var. *compactum* Herdman, 1924
 = *A. assymetricum* Kofoid, var. *compactum* Herdman, 1924
 Tromsö, W Brittan
- A. carteri*** Hulbert, 1957
 = *A. klebsii* : Carter, 1937, non Kofoid & Swezy, 1921
 Belt Sea, Oslofjord, S Norway, around Britain
- A. chattonii*** (Hovasse, 1922) Taylor, 1971
 = *Endodinium chattonii* Hovasse, 1922 (TM of *Endodinium* Hovasse, 1922)
 = *Zooxanthella chattonii* : Hovasse, 1924
 @ intracellular in *Vetella veletta*
 Engl. Channel, Mediterranean, Florida
- A. longum*** Lohmann, 1908
 all Baltic but Bothnian Bay & Sea, Skagerrak, Oslofjord, Bartens Sea, Adriatic
- A. nutricola*** (Brandt, 1881)
 = *Zooxanthella nutricula* Brandt, 1881 (TM of *Zooxanthella* Brandt, 1881)
 @ in the radiolarians *Collozoum inerme* & *Sphaerozoum punctatum*?
 ?
- A. bipes*** Herdman, 1924
 * sand-living.
 E North Sea, Isle of Man, Isle of Wight, Sussex, Kent, Yorkshire, Galloway
- A. crassum*** Lohmann, 1908
 = *phaeocysticola* Lebour, 1925
 * sand-living.
 central & W Baltic, Kattegatt, Belgium, Britain, Adriatic
- A. flexum*** Herdman, 1923
 * sand-living.
 Belgium, British shores
- A. incoloratum*** Campbell, 1973
 Kattegatt
- A. latum*** Lebour, 1925
 ?E North Sea, Belgium, Britain
- A. manannini*** Herdman, 1924
 * sand-living.
 Belgium, Britain
- A. pellucidum*** Herdman, 1922
 * sand-living.
 Bothnian Sea, Gulf of Finland, N Baltic, E North Sea, Belgium, Britain
- A. scissum*** Kofoid & Swezy, 1921
 = *A. scissoides* Lebour, 1925
 * sand-living.
 E North Sea, Britain, California
- A. sphenoides*** Wulff, 1916
 = *Gymnodinium filum* Lebour, 1917
 * sand-living.
 @ *Chaetoceros eibenii*
 Kattegatt, Helgoland, E North Sea, Barents Sea, Plymouth, Woods Hole
- A. corputulentum*** Kofoid & Swezy, 1921
 * sand-living.
 Kent, California
- A. globosum*** Schröder, 1911
 Sussex, Adriatic
- A. pelagicum*** Lebour, 1925
 Britain
- A. vitreum*** Herdman, 1924
 Britain
- A. herdmaniae*** Kofoid & Swezy, 1921
 * sand-living. Is the spelling *herdmani*?
 E North Sea, Britain
- A. lacustre*** von Stein, 1883
 = *A. schroderi* Schiller, 1928
 = *A. lacustriforme* Schiller, 1928
 * fresh & brackish water species.
 Belgium, Britain, Adriatic
- A. ovoideum*** (Lemmermann, 1896) Lemmermann, 1900
 = *Prorocentrum ovoideum* Lemmermann, 1896
 W Baltic, Belgium, Britain
- A. ovum*** Herdman, 1924
 * sand-living.
 E North Sea, Britain
- A. testudo*** Herdman, 1924
 * Sand-living. May be an encysting form of *A. herdmaniae* Kofoid & Swezy.
 E North Sea, Isle of Man
- A. psittacus*** Larsen, 1985
 * ?Sand-living.
 E North Sea
- A. pectilochroum*** Larsen, 1985
 * ?Sand-living.
 E North Sea
- A. phaeocysticola*** ??, 19??
 @ *Thalassiosira partheneia*
 NW Africa
- Cochlodinium*** Schütt, 1896 (TM *C. strangulatum* (Schütt) Schütt, 1896)
- ?= *Schillingia* Schiller, 19??
- C. achromaticum*** Lebour, 1925
 Plymouth
- C. brandtii*** Wulff, 1916
 = *C. augustum* Kofoid & Swezy, 1921
 off Plymouth
- C. helicooides*** Lebour, 1925
 = *C. helix* Schütt, 1895 (p.p.), non (Pouchet, 1887)
 W Baltic, ?Kattegatt, off Plymouth, Irish Sea, Bretagne, Mediterranean
- C. helix*** (Pouchet, 1887) Lemmermann, 1899
 = *Gymnodinium helix* Pouchet, 1887
 Plymouth, Atlantic
- C. pellucidum*** Lohmann, 1908
 Kieler Bucht, Kattegatt, Skagerrak
- C. pulchellum*** Lebour, 1917
 off Plymouth
- C. pupa*** Lebour, 1925
 ?Kattegatt, Engl. Channel, Irish Sea
- C. schuetii*** Kofoid & Swezy, 1921
 = *Gymnodinium helix* : Schütt, 1895 (p.p.), non Pouchet, 1887
 Plymouth, W Scotland
- C. vincentum*** Kofoid & Swezy, 1921
 Engl. Channel, off California
- Gymnodinium*** von Stein, 1878 (TM *G. fuscum* (Ehrenberg) von Stein, 1878 - fresh water)
 = *Ceratodinium* Conrad, 19?? = *Balechina* Loeblich Jr. & Loeblich III, 19??
 = *Aureodinium* Dodge, 1967
- G. abbreviatum*** Kofoid & Swezy, 1921
 List, Helgoland, E North Sea, Engl. Channel, California
- G. achromaticum*** Lebour, 1917
 Plymouth
- G. albulum*** Lindemann, 1928
 * May be a synonyme of *G. simplex* (Lohmann) Kofoid & Swezy
 S Baltic
- G. arenicolus*** Dragesco, 1965
 = *G. variabile* : Dodge, 1982, non Herdman, 1924
 * sand-living.
 Bretagne
- G. catenatum*** Graham, 19??
 * PSP-producing, toxic to mammals, birds & fish
 Portugal, Spain, may be expected in Scandinavia, as resting spores have been found in Öresund
- G. conicum*** Kofoid & Swezy, 1921
 = *G. viridis* Lebour, 1917
 Plymouth
- G. elongatum*** Hope, 1954
 Kattegatt, Oslofjord
- G. endofasciculum*** Campbell, 1973
 Kattegatt
- G. galatheanum*** Braarud, 19??
 * Toxic to fish. Often occurring together with *Gyrodinium aureolum*
 Norway, North Sea area, SW Africa
- G. gracile*** Bergh, 1881
 W Baltic, Belt Sea, N Kattegatt
- G. heterostriatum*** Kofoid & Swezy, 1921
 = *G. spirale* var. *obtusum* Dogiel, 1906
 = *G. lucidum*
 = *G. hyalinum* Lebour, 1925
 = *G. rhomboides* Lebour, 1925
 @ *Thalassiosira partheneia*
 Baltic, Skagerrak, NE Britain, outer Hebrides, Engl. Channel, Mediterranean, NW Africa, California
- G. incertum*** Herdman, 1924
 * sand-living.
 Isle of Man
- G. lebourae*** Pavillard, 1921
 Kattegatt
- G. lohmannii*** Paulsen, 1908
 = *G. roseum* Lohmann, 1908
 W Baltic, Kattegatt, Skagerrak, E North Sea
- G. marinum*** Saville-Kent, 1880-82
 * from an infusion of hay & sea-water
 Jersey
- G. micrum*** (Leadbeater & Dodge, 1966) Loeblich III, 1970
 = *Woloszynska micra* Leadbeater & Dodge, 1966
 Kattegatt, Engl. Channel
- G. minus*** Lebour, 1917
 Plymouth
- G. ostenfeldii*** Schiller, 1928
 Kattegatt
- G. pigmentosum*** (Dodge, 1967) Loeblich, 1970
 = *Aureodinium pigmentosum* Dodge, 1967 (TM of *Aureodinium* Dodge, 1967)
 Kattegatt, W Engl. Channel
- G. pseudonotoiluca*** Pouchet, 1885
 Kattegatt
- G. pygmaeum*** Lebour, 1925
 * ?possibly identical with *Gyrodinium aureolum* Hulbert.
 Kattegatt, Engl. Channel
- G. rhomboides*** Schütt, 1895
 W Baltic, Skagerrak
- G. rubrocinctum*** Lebour, 1925
 Kattegatt

- G. rubrum* Kofoid & Swezy, 1921
Kattegatt
- G. sanguineum* Hirasaka, 1924
= *G. splendens* Lebour, 1925
Belt Sea, Kattegatt, Oslofjord, E North Sea, S & W Britain, E & W North America
- G. simplex* (Lohmann, 19??) Kofoid & Swezy, 1921
Kattegatt
- G. variabile* Herdman, 1924
non = *G. variabile* : Dodge, 1982 (see *G. areniculus*)
* sand-living. May be a species complex.
?E North Sea, Isle of Man, S & W Britain, Port Erin, Bretagne
- G. veneficum* Ballantine, 1956
* toxic to fish
Devon
- G. vestifictii* Schütt, 1895
W Baltic, N Kattegatt, Skagerrak
- G. vitilago* Ballantine, 1956
Plymouth, NW Scotland
- G. placidum* Herdman, 1922
E North Sea
- G. parasitica* (Poche, 1903)
= *Oxyrrhis parasitica* Poche, 1903
@ gastrovascular cavities of *Diphyes elongata*, *Monophyes gracilis* & *Nanomia bijuga*
Mediterranean
- Gyrodinium* Kofoid & Swezy, 1921
= *Spirodinum* Schütt, 1896 = *Plectodinium* Biecheler, 1934
= *Sclerodinium* Dodge, 1981
- G. spirale* (Bergh, 1881) Kofoid & Swezy, 1921 (TM)
= *Gymnodinium spirale* Bergh, 1881
Worldwide distribution, W & S Baltic, Belt Sea, Kattegatt, E North Sea, Britain, Atlantic, Mediterranean
- G. aureolum* Hulbert, 1957
?= *Gymnodinium nagasakiense* ??,??
* Warm water species. Toxic to fish & invertebrates. Originally described from N America. As the European form may be another species, its name is sometimes written *G. cf. aureolum*.
S Baltic, Öresund, Limfjorden, Kattegatt, Bohusl., Oslofjord, S & W Norway, E North Sea, Helgoland, Britain, Massachusetts
- G. estuariale* Hulbert, 1957
Oslofjord, S Norway, Trondheim, Massachusetts
- G. undulans* Hulbert, 1957
Kattegatt, Oslofjord, Massachusetts
- G. britannicum* Kofoid & Swezy, 1921
= *Spirodinum spirale* var. *acutum* Lebour, 1917
E North Sea, Plymouth
- = *Sclerodinium calyptroglyphae* Lebour, 1925
= *Sclerodinium calyptroglyphae* (Lebour, 1925) Dodge, 1981 (TM of *Sclerodinium* Dodge, 1981)
Kattegatt, E North Sea, around Britain, central Atlantic
- G. cochlea* Lebour, 1925
Plymouth
- G. crassum* (Pouchet, 1885) Kofoid & Swezy, 1921
= *Gymnodinium crassum* Pouchet, 1885
Kattegatt, Engl. Channel, N Atlantic, Mediterranean
- G. cuneatum* Kofoid & Swezy, 1921
Bretagne
- G. dominans* Hulbert, 1957
Kattegatt
- G. falcatum* Kofoid & Swezy, 1921
Plymouth
- G. fissum* (Levander, 1894) Kofoid & Swezy, 1921
= *Gymnodinium fissum* Levander, 1894
= *Spirodinum fissum* (Levander, 1894) Lemmermann, 1???
all Baltic, Engl. Channel, Adriatic, California
- G. fusiforme* Kofoid & Swezy, 1921
= *Spirodinum fusus* Meunier, 1910
Öresund, N Kattegatt, Skagerrak, Arctic, Britain
- G. grave* (Meunier) Kofoid & Swezy, 1921
Kattegatt
- G. herbaceum* Kofoid & Swezy, 1921
?Kattegatt
- G. lachryma* (Meunier, 1910) Kofoid & Swezy, 1921
= *Spirodinum lachryma* Meunier, 1910
North Sea, Arctic, Kara Sea
- G. lebouriae* Herdman, 1924
* sand-living.
Belgium, Isle of Man
- G. linguliferum* Lebour, 1925
Plymouth
- G. longum* (Lohmann, 1908) Kofoid & Swezy, 1921
= *Cochlodinium longum* Lohmann, 1908
Kieler Bucht
- G. obtusum* (Schütt, 1895) Kofoid & Swezy, 1921
= *Gymnodinium spirale* Bergh var. *obtusa* Schütt, 1895
Plymouth, Adriatic, California
- G. opimum* (Schütt, 1895) Lebour, 1925
= *Gymnodinium opimum* Schütt, 1895
Plymouth, Mediterranean, Atlantic
- G. pellucidum* (Wulff, 1916) Schiller, 1933
= *Gymnodinium pellucidum* Wulff, 1916
* Possibly the gamete stage of *Polykrikos kofoidii*, sensu Morey-Gaines & Ruse, 1980
Barents Sea
- G. pepo* (Schütt, 1895) Kofoid & Swezy, 1921
= *Gymnodinium spirale* var. *pepo* Schütt, 1895
- ?Kattegatt, Plymouth, Mediterranean, Atlantic
- G. pingue* (Schütt, 1895) Kofoid & Swezy, 1921
= *Gymnodinium spirale* var. *pinguis* Schütt, 1895
?Kattegatt, List, E North Sea, Barents Sea, Plymouth, Mediterranean, Atlantic
- G. prunus* (Wulff, 1920) Lebour, 1925
= *Spirodinum prunus* Wulff, 1920
Helgoland, E North Sea, Barents Sea, Adriatic
- G. resplendens* Hulbert, 1957
Kattegatt, Cornwall, Woods Hole
- G. striatum* (Dodge, 1981)
= *Sclerodinium striatum* Dodge, 1981
North Sea, off NW Africa
- G. uncatenatum* Hulbert, 1957
Hebrides, Woods Hole
- Katodinium* Fott, 1957
= *Massartia* Conrad, 1926, non Wildemann, 1897 (Zygomycetes)
- K. nieuportense* (Conrad, 1926) Loeblich Jr & Loeblich III, 1966 (TM)
= *Massartia nieuportensis* Conrad, 1926
Belgian brackish water
- K. rotundatum* (Lohmann, 1908) Fott, 1957
= *Amphidinium rotundatum* Lohmann, 1908
= *Massartia rotundata* (Lohmann, 1908) Schiller, 1933-37
= *Gymnodinium minutum* Lebour, 1925
= *Amphidinium pellucidum* Redcke, 1935
= *Amphidinium redekei* Conrad & Kuferath, 1954
All Baltic except Bothnia Sea & Bay, Kattegatt, Skagerrak, Oslofjord, E North Sea, Belgium, Tromsö, Barents Sea, White Sea, all around Britain, Adriatic
- K. asymmetricum* (Massart, 1900) Loeblich III, 1965
= *Gymnodinium asymmetricum* Massart, 1900
* in wet sands & salt marshes.
E North Sea, Belgium, S & W Britain, France
- K. fungiforme* (Anissimowa, 1926) Loeblich III, 1965
= *Gymnodinium fungiforme* Anissimowa, 1926
* in wet sands & salt marshes.
Engl. Channel, Galloway, Lewis, France
- K. glandulatum* (Herdman, 1924) Loeblich III, 1965
= *Gymnodinium glandula* Herdman, 1924
* in wet sand.
E North Sea, around most of Britain
- K. glaucum* (Lebour, 1917) Loeblich III, 1965
= *Gyrodinium glaucum* (Lebour, 1917) Kofoid & Swezy, 1921
= *Spirodinum glaucum* Lebour, 1917
= *Massartia glauca* : Schiller, 1933
Kattegatt, E North Sea, Plymouth, Adriatic
- Pavillardia* Kofoid & Swezy, in Kofoid, 1920 (TM *P. tentaculifera* Kofoid & Swezy, in Kofoid, 1920 - California)
- Protodinium* Lohmann, 1908
* Genus not recognized by Taylor, 1987, but by Sournia, 1986
- Protodinium simplex* Lohmann, 1908 (TM)
= *Gymnodinium simplex* (Lohmann, 1908) Kofoid & Swezy, 1921
Gulf of Finland, S, W & N Baltic, Belt Sea - Kattegatt area, E North Sea, NE & NW England, Engl. Channel, N Atlantic
- Pseliodinium* Sournia, 1972 (TM *P. vaubani* Sournia, 1972 - e.g. Mediterranean)
* Genus not recognized by Taylor, 1987.
- Symbiodinium* Freudenthal, 1962
@ Anthozoa
* Not mentioned by Larsen & Sournia, 1992, in Patterson & Larsen
- S. microadriaticum* Freudenthal, 1962
= *Zooxanthella microadriatica* (Freudenthal, 1962) Loeblich & Sherley, 1979
@ Coelenterata, e.g. *Anemona sulcata*
Britain (probably world-wide)
- Torodinium* Kofoid & Swezy, 1921
- T. teredo* (Pouchet) Kofoid & Swezy, 1921 (TM)
N Kattegatt, Skagerrak
- T. robustum* Kofoid & Swezy, 1921
Kattegatt, E North Sea, Essex, Plymouth
- Lophodinium* Lemmermann, 1910
* Not mentioned by Larsen & Sournia, 1992, in Patterson & Larsen
- Woloszynskia* Thompson, 1951 (TM *W. reticulata* Thompson, 1951 - fresh water)
* Not mentioned by Larsen & Sournia, 1992, in Patterson & Larsen
- W. danica* (Paulsen, 1907)
= *Glenodinium danicum* Paulsen, 1907
Arkona Sea, Belt Sea, Öresund, N Kattegatt, Skagerrak, E North Sea, Barents Sea, around Britain except Engl. Channel
- Polykrikaceae* Kofoid & Swezy, 1921
- Polykrikos* Bütschli, 1873
- P. schwartzii* Bütschli, 1873 (TM)
= *P. auricularia* Bergh, 1881
W Baltic, Öresund, Kattegatt, Skagerrak, Oslofjord, S Norway, around all Britain
- P. kofoidii* Chatton, 1914
off Plymouth
- P. lebourae* Herdman, 1924
* on wet sandy beaches.
around most of Britain
- Pheopolykrikos* Chatton, 1933
- P. beauchampi* Chatton, 1933 (TM)
= *Polykrikos beauchampi* (Chatton, 1933) Loeblich III, 1980
Newport, S France
- Ptychodiscaceae*
- Ptychodiscus* von Stein, 1883

- P. noctiluca* von Stein,1883 (TM)
 = *P. carinatus* Kofoid,1907
 = *P. inflatus* Pavillard,1916
 = *Diploctysis antarctica* Cleve,1900
 off N, W & SW Britain, Biscay, Mediterranean, Atlantic
- Warnowiaceae** Lindemann,1928
- Erythropsidinium* Silva,1960 (TM *E. agile* (Hertwig,1884) Silva,1960 - Mediterranean)
- Gruetodinium* Loeblich III,1980 (TM *G. cylindricum* (Gruet) Loeblich III,1968 - Mediterranean)
 = *Leucopsis* Gruet,1968?
- Nematodinium* Kofoid & Swezy,1921 (TM *N. partitum* Kofoid & Swezy,1921 - California)
 = *Nematopsisides* Gruet,1973
- N. armatum* (Dogiel,1906) Kofoid & Swezy,1921
 = *Pouchetia armata* Dogiel,1906
 Helgoland, E North Sea, Plymouth
- N. vigilans* (Marshall,1925)
 = *Nematopsisides vigilans* (Marshall,1925) Gruet,1973 (TM of *Nematopsisides* Gruet,1973)
 = *Proterothropsis vigilans* Marshall,1925
 Kattegatt, Helgoland, E North Sea, W Britain
- Warnowia** Lindemann,1928
 = *Pouchetia* Schütt,1895 = *Protopsis* Kofoid & Swezy,1921
 = *Proterothropsis* Kofoid & Swezy,1921 (TM *P. crassicaudata* Kofoid & Swezy,1921 - California)
- W. fusus* (Schütt,1895) Lindemann,1928 (TM)
 Britain
- W. nigra* (Pouchet,1887)
 = *Protopsis nigra* (Pouchet,1887) Kofoid & Swezy,1921 (TM of *Protopsis* Kofoid & Swezy,1921)
 = *Gymnodinium polyphemus* var. *nigrum* Pouchet,1887
 Concarneau
- W. parva* (Lohmann,1908) Lindemann,1928
 = *Pouchetia parva* Lohmann,1908
 W Baltic
- W. polyphemus* (Pouchet) Schiller,1933
 = *Pouchetia polyphemus* (Pouchet) Kofoid & Swezy,1921
 Engl. Channel, Atlantic
- W. rosea* (Pouchet,1897) Kofoid & Swezy,1921
 = *Gymnodinium polyphemus* var. *roseum* Pouchet,1897
 non = *Pouchetia rosea* Schütt,1895
 ?Kattegatt, Britain
- W. schuetti* (Kofoid & Swezy,1921) Schiller,1933?
 = *Pouchetia rosea* Schütt,1895
 Britain
- W. simplex* (Lebour,1925)
 = *Protopsis simplex* Lebour,1925
 Plymouth
- Gymnodiniales "incertae sedis"**
- Bernardinum* Chodat,1924 (TM *B. bernardinense* Chodat,1924 - fresh water)
- Filodinium* Cachon & Cachon,1968 (TM *F. hovassei* Cachon & Cachon,1968 - Mediterranean)
- Herdmania** Dodge,1981
- H. litoralis* Dodge,1981 (TM)
 = *Gymnodinium agile* Herdman,1922, non Kofoid & Swezy,1921
 * on sandy beaches.
 E North Sea, Yorkshire, Sussex, Kent, Dorset, Isle of Man, SW Scotland
- PERIDINIALES** Haeckel,1894
- Amphidomataceae**
- Amphidoma* von Stein,1883 (TM *A. nucula* von Stein,1883?)
- A. caudata* Halldal,1953
 = *Oxytoxum tonoli* Rønning,1969
 = *Oxytoxum margalefi* Rønning,1969
 Norway, off E Scotland, Galway, off S & W Britain, Portugal, Mediterranean
- Ceratiaceae** Kofoid,1907
- Ceratium* Schranck,1793 (TM *C. pleuroceras* Schrank,1793 - fresh water)
- C. (Archaeoceratium* Jørgensen,1911?) *gravidum* Gourret,1883
 off W Britain, tropical Atlantic
- C. (Ceratium) candelabrum* (Ehrenberg,1860) von Stein,1883
 = *Peridinium candelabrum* Ehrenberg,1860
 Skagerrak, off SW Britain, warm Atlantic
- C. (Ceratium) furca* (Ehrenberg,1836) Claparède & Lachmann,1858
 = *Peridinium furca* Ehrenberg,1836
 Cosmopolitan, W & S Baltic, Öresund, Kattegatt, Skagerrak, Bohusl, Oslofjord, North Sea, all around Britain
- C. (Ceratium) hirundinella* (O.F. Müller,1773) Schrank,1793
Bursaria hirundinella O.F. Müller,1773
 * Freshwater species, which does not tolerate the full salinity of the area.
 estuaries & bays around Finland
- C. (Ceratium) lineatum* (Ehrenberg,1854) Cleve,1899
 = *Peridinium lineatum* Ehrenberg,1854
 W Baltic, Öresund, Kattegatt, Skagerrak, Oslofjord, North Sea, all around Britain
- C. (Ceratium) minutum* Jørgensen,1920
 off N & W Britain
- C. (Ceratium) pentagonum* Gourret
 warm Atlantic (occasionally in Gulf Stream waters)
- C. (Ceratium) teres* Kofoid,19
 warm Atlantic (occasionally in Gulf Stream waters)
- C. (Ceratium) setaceum* Jørgensen,19
 warm Atlantic (occasionally in Gulf Stream waters)
- C. (Amphiceratium* Vanhöffen,1896) *fusus* (Ehrenberg,1834) Dujardin,1841
 = *Peridinium fusus* Ehrenberg,1834
 fairly cosmopolitan, W, S & central Baltic, Öresund, Kattegatt, Bohusl, Skagerrak, Oslofjord, North Sea, all around Britain
- C. (Amphiceratium** Vanhöffen,1896) *longirostrum* Gourret,
 warm Atlantic (occasionally in Gulf Stream waters)
- C. (Amphiceratium** Vanhöffen,1896) *inflatum* (Kofoid,19) Jørgensen,19
 warm Atlantic (occasionally in Gulf Stream waters)
- C. (Amphiceratium** Vanhöffen,1896) *extensum* (Gourret,18) Cleve
 warm Atlantic (occasionally in Gulf Stream waters)
- C. (Orthoceratium** Vanhöffen,1896) *azoricum* Cleve,1900
 off W & N Britain
- C. (Orthoceratium** Vanhöffen,1896) *petersii* Nielsen
 off W Ireland
- C. (Orthoceratium** Vanhöffen,1896) *breve* (Ostenfeld & Schmidt) Schröder
 warm Atlantic
- C. (Orthoceratium** Vanhöffen,1896) *compressum* Gran,1902
 off N & W Britain
- C. (Orthoceratium** Vanhöffen,1896) *platycorne* von Dayad,1888
 = *C. lamelliforme* Kofoid,1908
 warm Atlantic (occasionally in Gulf Stream waters)
- C. (Orthoceratium** Vanhöffen,1896) *tripos* (O.F. Müller,1777) Nitzsch,1817
 = *Cercaria tripes* O.F. Müller,1777
 Cosmopolitan, S & W Baltic, Öresund, Kattegatt, Skagerrak, Oslofjord, North Sea, all around Britain
- C. (Orthoceratium** Vanhöffen,1896) *tripos* (O.F. Müller,1777) Nitzsch,1817 var. *subsalsum*
 (Ostenfeld,1903) Paulsen,1907
 W-S Baltic
- C. (Orthoceratium** Vanhöffen,1896) *arietinum* Cleve,1900
 = *C. bucephalum* Cleve,1897
 * Why not use the older name?
 W Baltic, Belt Sea, Öresund, Kattegatt, Skagerrak, Oslofjord, North Sea, off N & W Britain
- C. (Orthoceratium** Vanhöffen,1896) *symmetricum* Pavillard,19
 E Atlantic
- C. (Orthoceratium** Vanhöffen,1896) *gibberum* Gourret
 ?
- C. (Orthoceratium** Vanhöffen,1896) *macroceros* (Ehrenberg,1840) Vanhöffen,1897
 = *Peridinium macroceros* Ehrenberg,1840
 Arcona Sea, W Baltic, Öresund, Kattegatt, Skagerrak, Oslofjord, North Sea, all around Britain
- C. (Orthoceratium** Vanhöffen,1896) *massiliense* (Gourret) Jørgensen,19
 E & Tropical Atlantic
- C. (Orthoceratium** Vanhöffen,1896) *trichoceros* (Ehrenberg,18) Kofoid,19
 off W Ireland, Tropical Atlantic
- C. (Orthoceratium** Vanhöffen,1896) *carriense* Gourret,
 off W Ireland, off Bretagne, Tropical Atlantic
- C. (Orthoceratium** Vanhöffen,1896) *hexacanthum* Gourret,1883
 = *C. reticulatum* (Pouchet,1883) Cleve,1903
 = *C. tripes* var. *reticulatum* Pouchet,1883
 off N Scotland, Engl. Channel, warm Atlantic
- C. (Orthoceratium** Vanhöffen,1896) *longipes* (Bailey,1855) Gran,1902
 = *Peridinium longipes* Bailey,1855
 W Baltic, Öresund, Kattegatt, Skagerrak, Oslofjord, North Sea, around most of Britain, cool Atlantic
- C. (Orthoceratium** Vanhöffen,1896) *longipes* (Bailey,1855) Gran,1902 var. *baltica* Ostenfeld,1903
 W Baltic (occasionally)
- C. (Orthoceratium** Vanhöffen,1896) *arcticum* (Ehrenberg,18) Cleve
 N Kattegatt, Skagerrak, North Sea, N Scotland, Arctic waters
- C. (Orthoceratium** Vanhöffen,1896) *horridum* (Cleve,1897) Gran,1902
 = *C. tripes* var. *horridum* Cleve,1897
 = *C. tenuie* (Ostenfeld & Schmidt) Jørgensen,1911
 = *C. intermedium* (Jørgensen) Jørgensen,1905
 = *C. buceros* Zacharias, emend. Bohn, in Schiller,1937
 = *C. batavum* Paulsen,1908
 Belt Sea, Kattegatt, Skagerrak, Oslofjord, North Sea, all around Britain
- Ceratocoryathaceae** Lindemann,1928
- Ceratocorys** von Stein,1883 (TM *C. horrida* von Stein,1883)
- Cladopyxiidae** Poche,1913
- Cladopyxis** von Stein,1883 (TM *C. brachiolata* von Stein,1883 (p.p.))
 = *Acanthodinium* Kofoid,19
- Palaeophalacroma** Schiller,1928 (TM *P. verrucosum* Schiller,1928)
 = *Epiperidinium* Gardner,1954
 = *Sinodinium* Nie,19
- P. unicinctum** Schiller,1928
 = *Heterodinium detoni* Rønning,1943
 = *Epiperidinium michaelarsi* Gaarder,1954
 N & S Atlantic, Adriatic
- Cryptocodiinae** Biecheler,1952
- Cryptocodium** Biecheler,1938
- C. cohnii* (Seligo, in Cohn,1887) Chatton, in Grassé,1952
 = *Glenodinium cohnii* Seligo, in Cohn,1887
 = *Gymnodinium furciforme* Küster,1908
 = *Gyrodinium cohnii* (Seligo,1887) Schiller,1933
 = *Cryptocodium setense* Biecheler,1938 (TM)
 * littoral, amongst decaying seaweeds.
 Baltic, Oslofjord, S Norway, France, Mediterranean
- Hemidinium** von Stein,1878
 * Genus not recognized by Taylor,1987, but placed in this fam. by Sournia,1986
- H. nasutum** von Stein,1878
 * May be the motile development stage of *Gloeodinium montanum* Klebs,1912
 European fresh, brackish & salt lake waters, e.g. Gulf of Finland, Riga Bay
- H. ochraceum** Levander,1900
 * Main occurrence in rock pools.

Gulf of Finland

Goniomitaceae Lindemann,1928 = Triadiiniaceae

Alexandrium Halim,1960 (TM A. minutum) Halim,1960 - E Mediterranean
= *Protogonyaulax* Taylor,1979

A. tamarens (Lebour,1925) Balech & Tangen,1985

= *Protogonyaulax tamarens* (Lebour,1925) Taylor,1979 (TM of *Protogonyaulax* Taylor,1979)

= *Gonyaulax tamarens* Lebour,1925

= *Gonyaulax excavata* (Braarud,1945) Balech,1971

= *Gonyaulax globosa* (Braarud,1945) Balech,1971

= *Gonyaulax trygvii* Parke,1976

= *Pyrodinium phoeniceus* Woloszynski & Conrad,1939

= *Gessnerium tamarens* Loeblich & Loeblich,1979

= *Protogonyaulax tamarens* : Taylor,1979

* PSP-producing, toxic to mammals, birds & fish.

Öresund, Kattegatt, Limfjorden, Oslofjord, Norway, E North Sea, Faeroes, around Britain, E USA

A. ostenfeldii (Paulsen,1904) Balech & Tangen,1985

= *Goniiodoma ostenfeldii* Paulsen,1904

= *Heteraulacus ostenfeldii* (Paulsen,1904) Loeblich III,1970

= *Gessnerium ostenfeldii* (Paulsen,1904) Loeblich III & Loeblich,1979

?= *Pyrodinium phoeniceus* Woloszynski & Conrad,1939

* PSP-producing, toxic to mammals, birds & fish.

Gdansk, Belt Sea, Öresund, N Kattegatt, Skagerrak, Limfjorden, North Sea

A. minutum Halim,1960

* Causing mussel poisoning in France.

France, Portugal, Spain, Italy, Egypt, E North America, Australia

A. catenella ??,1???

North America

A. fundyense ??,1???

E Canada

Gambierdiscus Adachi & Fukuyo,1979 (TM G. toxicus) Adachi & Fukuyo,1979

Gessnerium Halim, ex Halim,1969 (TM G. mochimaense) Halim, ex Halim,1969

Genus considered a synonym of Alexandrium Halim, by Taylor,1987

Goniiodoma von Stein,1883 (preoccupied - Lepidoptera)

= *Triadinium* Dodge,1981 (preoccupied - Ciliophora)

= *Heteraulacus* Diesing,1850 (p.p.)

G. acuminatum (Ehrenberg,1835) von Stein,1883 (TM)

= *Peridinium acuminatum* Ehrenberg,1835

= *Peridinium polyedricum* Pouchet,1883

= *Goniiodoma polyedricum* (Pouchet,1883) Jörgensen,1900

= *Heteraulacus polyedricus* (Pouchet,1883) Drugg & Loeblich jr.,1967

Kieler Bucht, Greifswald

G. polyedricum (Pouchet,1883) Jörgensen,1899

= *Goniiodoma acuminata* von Stein,1883 (p.p.), non *Peridinium acuminatum* Ehrenberg,1834, nec *Heteraulacus acuminatus* : Diesing,1850

= *Peridinium polyedricum* Pouchet,1883

= *Triadinium polyedricum* (Pouchet,1883) Dodge,1981

Britain, Mediterranean, Tropical seas

G. sphaericum Murray & Whitting,1899

= *Triadinium sphaericum* (Murray & Whitting,1899) Dodge,1981

Engl. Channel, Atlantic, Caribbean

Pyrodinium Plate,1906 (TM P. bahamense) Plate,1906

Gonyaulacaceae Lindemann,1928

Amylax Meunier,1910

A. triacantha (Jörgensen,1899) Sournia,1984

= *Gonyaulax triacantha* Jörgensen,1899

= *Amylax lata* Meunier,1910 (TM)

all Baltic except Bothnian Bay, Öresund, Kattegatt, Skagerrak, Oslofjord, Norway, Iceland, North Sea, N & W Britain

Gonyaulax Diesing,1866

G. spinifera (Claparède & Lachmann,1859) Diesing,1866 (TM)

= *Peridinium spiniferum* Claparède & Lachmann,1859

= *Gonyaulax levanderi* (Lemmermann,1900) Paulsen,1907

* 3 different cyst types : *Spiniferites mirabilis* (Rossignol) Sarjeant, *Nematospaeropsis labyrinthea* (Ostenfeld) Reid & *Tectatodinium peltatum* Wall have been germinated to give motile cells of *G. spinifera* type.

all Baltic except Bothnian Sea & Bay, Öresund, Kattegatt, Skagerrak, Oslofjord, E North Sea, all around Britain, most parts of the world

G. alaskensis Kofoid,1911

North Sea, Engl. Channel, Mediterranean

G. diegensis Kofoid,1911

E North Sea, N & W Britain, Pacific

G. digitale (Pouchet,1883) Kofoid,1911

= *Protoperidinium digitale* Pouchet,1883

* Cyst: *Spiniferites bentori* Ross.) Wall & Dale.

W Baltic, Kattegatt, Limfjorden, Oslofjord, E North Sea, Britain (mainly to the W & N)

G. helensis Woloszynska,1928

* Requires taxonomic reinvestigation.

Gdansk

G. polygramma von Stein,1883

= *G. schuetti* Lemmermann,1899

= *Protoperidinium pyrophyrum* Pouchet,1883

Skagerrak, Oslofjord, around Scotland, occasionally SW Britain

G. scrippssae Kofoid,1911

* Cyst: *Spiniferites bulbideus* (Deflandre & Cookson) Sarjeant

off Yorkshire, SW Britain

G. verior Sournia,1973

= *Amylax diacantha* Meunier,1919

= *Gonyaulax diacantha* (Meunier,1919) Schiller,1937, non = *Gonyaulax diacantha* Athanassopoulos,1931

= *Gonyaulax longispina* Lebour,1925

all Baltic except Bothnian Sea & Bay, Öresund, S Kattegatt, Oslofjord, Belgium, North Sea, N & W Britain, Mediterranean

Peridiniella Kofoid & Michener,1911 (TM P. sphaeroidea) Kofoid & Michener,1911

P. catena (Levander,1894) Balech,1977

= *Peridinium catenatum* Levander,1894

= *Gonyaulax catenata* (Levander,1894) Kofoid,1911

* Cold water & low salinity species.

All Baltic, Öresund, S Kattegatt, Limfjorden, Oslofjord

Protoceratum Bergh,1881 (TM P. aceros) Bergh,1881

P. reticulatum (Claparède & Lachmann,1859) Bütschi,1885

= *Peridinium reticulatum* Claparède & Lachmann,1859

= *Protoceratum aceros* Bergh,1881

= *Gonyaulax grindleyi* Reinecke,1967

non = *Gonyaulax reticulatum* Kofoid & Mitchener,1911

* Cyst: *Operculodinium centrocarpum* (Deflandre & Cookson) Wall

all Baltic, Öresund, Kattegatt, Skagerrak, Oslofjord, E North Sea, all around Britain, many parts of the world

Lingulodinium ??,1???

L. polyedra (von Stein,1883) Dodge,1899

= *Gonyaulax polyedra* von Stein,1883

* Cyst: *Lingulodinium machaerophorum* (Deflandre & Cookson) Wall

Arkona Sea, Belt Sea, Öresund, Kattegatt, Skagerrak, Oslofjord, N & W Britain (North Sea occasionally)

Shuettiella ??,1???

Heterodiniae Lindemann,1928

Dolichodinium Kofoid & Adamson,1933 (TM D. lineatum) (Kofoid & Michener) Kofoid & Adamson,1933 - tropical Atlantic & Pacific

Heterodinium Kofoid,1906 (TM H. scrippssae) Kofoid,1906

H. milneri (Murray & Whitting,1899) Kofoid,1906

= *Peridinium milneri* Murray & Whitting,1899

occasionally off SW Britain, central Atlantic

H. murrayi Kofoid,1906

= *Peridinium tripos* Murray & Whitting,1899

off Plymouth (one specimen), tropical Atlantic

Ostreopsis Schmidt,1901 (TM O. siamensis) Schmidt,1901

Oxytoxaceae Lindemann,1928

Coolia Meunier,1919

C. monotis Meunier,1919 (TM)

* littoral

Belgium, Britain, Bretagne, Mediterranean

Ostreopsis Schmidt,1901 (TM O. siamensis) Schmidt,1901

Oxytoxaceae Lindemann,1928

Amphidiniopsis Woloszynska,1929

= *Thecadinium* Kofoid & Skogsberg,1928

A. hirsutum (Balech,1956) Dodge,1982

= *Thecadinium hirsutum* Balech,1956

* In sandy beaches

Dorset, Bretagne

A. kofoidii Woloszynska,1929 (TM)

* In sandy beaches

Riga Bay, S Baltic, Sussex

A. petasatum (Herdman,1922)

= *Thecadinium petasatum* (Herdman,1922) Kofoid & Skogsberg,1928 (TM of *Thecadinium* Kofoid & Skogsberg,1928)

= *Amphidinium kofoidi* var. *petasatum* Herdman,1922

* sand-living

W Britain, Port Erin, Bretagne

A. semilunatum (Herdman,1923)

= *Thecadinium semilunatum* (Herdman,1923) Dodge,1982

= *Amphidinium semilunatum* Herdman,1923

= *Thecadinium inclinatum* Balech,1956

* sand-living

Gdansk, E North Sea, Isle of Man, Kent, Port Erin

A. swedmarkii (Balech,1956) Dodge,1982

= *Thecadinium swedmarki* Balech,1956

* In sandy beaches

Galloway, Dorset, Isle of Man, Bretagne

Centroldinium Kofoid,1907 (TM C. elongatum) Kofoid,1907

Corythodinium Loeblich Jr. & Loeblich III,1966 (TM C. tessellatum) (von Stein,1883?) Loeblich Jr. & Loeblich III,1966

= *Pyrgiadium* von Stein,1883

= *Murrayella* Kofoid,1907, non Schmitz

= *Pavillardinium* de Tonti,1936

C. diploconus (von Stein,1883) Taylor,1976

= *Oxytoxum diploconus* von Stein,1883

off Norway, off SW Ireland

C. michaelsarsi (Gaarder,1954) Taylor,1976

= *Oxytoxum michaelsarsi* Gaarder,1956

off Iberia, N Atlantic

C. reticulatum (von Stein,1883) Taylor,1976

= *Pyrgiadium reticulatum* von Stein,1883

= *Oxytoxum reticulatum* Bütschi,1885

?= *Oxytoxum brunellii* Rampa,1939

N Atlantic, Mediterranean

Oxytoxum von Stein,1883

O. scolopax von Stein,1883 (TM)

off W & SW Britain, warm Atlantic

O. laticeps Schiller,1937

off Norway, N Atlantic, Adriatic

O. milneri Murray & Whitting,1899

N Atlantic, Mediterranean

O. sphaeroideum von Stein,1883

off W Britain, S Atlantic

Planodinium Saunders & Dodge,1984

* Sand-living

Roscoffia Balech,1956

R. capitata Balech,1956 (TM)

* In sandy beaches

Roscoff

- Sabulodinium** Saunders & Dodge,1984
* Sand-living
* Genus not recognized by Sournia,1986 or Taylor,1987
- S. undulatum** Saunders & Dodge,19
* Sand-living.
E North Sea
- Thecadiniopsis** ??,19??
- Calcidinellaceae**
- Ensculifera** Balech,1967
- E. mexicana** Balech,1967 (TM)
Scrippsiella mexicana (Balech,1967) Indelicato & Loeblich,1986
Norway, Spain, tropical seas
- Scrippsiella** Balech,1959, ex Loeblich III (TM *S. sweeneyae* Balech,1959?)
= *Calcidinellum* ?,19??
* Lewis,1991 erected a number of new *Scrippsiella* species on basis of cyst morphology. Which of these may be found in north European waters?
- S. trochoidea** (von Stein,1883) Loeblich III,1976
= *Glenodinium trochoideum* von Stein,1883
= *Glenodinium acuminatum* Jørgensen,1899
= *Peridinium trochoideum* (von Stein,1883) Lemmermann,1910
= *Scrippsiella faeroense* : Dickensheets & Cox,1971, non (Paulsen,1905) Balech & Soares,1967
All Baltic except Bothnian Sea & Bay, Kattegatt, Limfjorden, Skagerrak, E North Sea, N Norway, all around Britain, Atlantic
- S. faeroense** (Paulsen,1905) Balech & Soares,1967
= *Peridinium faeroense* Paulsen,1905
Oresund, Oslofjord, E North Sea, Norway, Faeroes, all around Britain
- Peridiniidae = Peridiniaceae Ehrenberg,1828
- Boreadinum** Dodge & Hermes,1981 (TM *B. pisiforme* Dodge & Hermes,1981)
* Correct family?
- Glenodinium** Ehrenberg,1830, emend. von Stein,1878 (TM *G. cinctum* Ehrenberg,1837 - fresh water)
= *Sphaerodinium*
* Not mentioned by Larsen & Sournia,1992, in Patterson & Larsen
- G. danicum** Paulsen,1907
Kattegatt
- G. monensis** Herdman,1923
* Sand-living. Should probably be moved to another genus.
Isle of Wight, Isle of Man
- G. gymnodinum** Penard,1891
* Freshwater species, which does not tolerate the full salinity of the area.
all Baltic except Belt Sea
- G. paululum** Lindemann,1928
Gulf of Finland, Greifswald
- G. warmingii** Bergh,1882
Baltic Sea
- G. armatum** Levander,1900
Finland, Darsser area
- G. penardiforme** (Lindemann) Schiller
* The determination need confirmation.
Gulf of Finland
- Gotoiu** Abé,1981
* Correct family?
- G. eccentricum** (Nie,1943) Sournia,1984
= *Dissodium eccentricum* (Nie,1943) Loeblich III,1970
= *Diplopsalis eccentrica* Nie,1943
= *Gotoiu mutsuensis* Abé,1981 (TM)
* North Sea distribution doubtful, according to Sournia,1986
British North Sea coast, China Sea
- Cachonina** Loeblich III,1968
* Synonymized with *Heterocapsa* von Stein,1883, by Morrill & Loeblich,1981
- C. hallii** (Freudenthal & Lee,1963) Dodge,1982
= *Glenodinium hallii* Freudenthal & Lee,1963
= *Cachonina nisi* Loeblich III,1968 (TM)
= *Cachonina illdefinita* Herman & Sweeney,1976
SW Kattegatt, Limfjorden, E North Sea, off German North Sea coast
- Heterocapsa** von Stein,1883
- H. triquetra** (Ehrenberg,1840) von Stein,1883 (TM)
= *Glenodinium triquetrum* Ehrenberg,1840
= *Peridinium heterocapsa* Meunier,1919
= *Peridinium triquetrum* (Ehrenberg,1840) Lebour,1925
All Baltic except Riga Bay & Bothnian Sea & Bay, Öresund, Kattegatt, Skagerrak, Oslofjord, S, W & N Norway, E North Sea, all Britain
- H. minima** Pomroy,1989
Kattegatt
- Kansodinium** ??,19??
* Correct family?
- Kryptoperidinium** Lindemann,1924
= *Phyldinium* Conrad,1926
* Genus not recognized by Taylor,1987
* Correct family?
- K. foliaceum** (von Stein,1883) Lindemann,1924 (TM)
= *Glenodinium foliaceum* von Stein,1883
* brackish water species
all Baltic, S Kattegatt, E North Sea, Britain, Mediterranean
- Pentapharsodinium** Indelicato & Loeblich,1986
* This genus should perhaps be moved to Calcidinellaceae
- P. datei** Indelicato & Loeblich,1986 (TM)
= *Peridinium faeroense* : Dale,1977, non Paulsen,1905
= *Scrippsiella faeroense* : Dodge,1982, p. 153, fig 18N, non Paulsen,1905
?Kattegatt
- Peridiopsis** Lemmermann,1928?
* Correct family?
- P. balticum** (Levander,1894/95) Bourrelly,19??
= *Peridinium balticum* (Levander,1894/95) Lemmermann,1910
= *Glenodinium balticum* Levander,1894/95
all Baltic except Bothnian Sea & Bay and Belt Sea
- Peridinium** Ehrenberg,1832
= *Thompsodinium* ??,19??
* With no really marine species.
- P. hangoei** Schiller,1937
= *P. gracile* Lindemann,19??
Bothnian Sea, N Baltic, Gulf of Finland
- P. williei** Huitfeldt-Kaas,1900
= *P. volzii* Lemmermann,19??
* Freshwater species, which does not tolerate the full salinity of the area.
Finnish bays
- P. cinctum** (O.F. Müller,17??) Ehrenberg,1832 (TM)
* Freshwater species, which does not tolerate the full salinity of the area.
Finnish bays
- P. umbonatum** von Stein,1883 var. **umbonatum** von Stein,1883
= *P. inconspicuum* Lemmermann,1899
= *P. munuscum* Lindemann,19??
= *P. pusillum* (Penard,17??) Lemmermann,19??
* Freshwater species, which does not tolerate the full salinity of the area.
all Baltic
- Protoperidiinidae
- Diplopsalidoideae**
- Diplopsalis** Bergh,1882, non Sclater,1866 (Aves - sub-genus)
- D. lenticula** Bergh,1882 (TM)
= *Glenodinium lenticula* (Bergh,1882) Schiller,1937
= *Dissodium lenticulum* (Bergh,1882) Loeblich III,1970
Gdansk, Belt Sea, Öresund, Kattegatt, Skagerrak, Oslofjord, North Sea, all around Britain, Mediterranean
- Entzia** Lebour,1922, non von Daday,1883 (Foraminiferida)
= *Asteinia* Abé,19??
* This genus is probably a synonym of *Diplopsalis* Bergh,1882, sensu Larsen & Sournia,1992 in Patterson & Larsen
- E. acuta** (Apstein,1896) Lebour,1922
= *Glenodinium acutum* Apstein,1896
= *Diplopsalis acuta* (Apstein,1896) Entz,1904
= *Peridinium latum* Paulsen,1908
* Freshwater species, which does not tolerate the full salinity of the area.
Bothnian Bay, Gulf of Finland, Gdansk, Arcona Sea
- Diplopelta** von Stein, ex Jørgensen,1912
- D. bomba** von Stein, ex Jørgensen,1912
= *Dissodium asymmetricum* (Mangin,1911) Loeblich III,1970
= *Peridiopsis asymmetrica* Mangin,1911
= *Diplopelta symmetrica* Pavillard,1913
= *Diplopsalis asymmetrica* : Drobis & Elbrächter,1976
Baltic, Limfjorden, E North Sea, all around Britain, Mediterranean, Atlantic
- D. pusilla** Balech & Akselman,1988
Kattegatt
- Lebouria** Abé,1941 (TM *L. minuta* Abé,1941 - Japan)
* Genus not recognized by Taylor,1987, but by Sournia,1986.
- Diplopsalopsis** Meunier,1910
- D. orbicularis** (Paulsen,1907) Meunier,1910 (TM)
= *Peridinium orbiculare* Paulsen,1907
Öresund, Limfjorden, Kattegatt, Skagerrak, North Sea, Iceland, Engl. Channel
- Oblea** Balech, ex Loeblich Jr. & Loeblich III,1966
- O. baculifera** Balech, ex Loeblich Jr. & Loeblich III,1966 (TM)
Oslofjord
- O. rotunda** (Lebour,1922) Balech, ex Sournia,1973
= *Peridiopsis rotunda* Lebour,1922
= *Glenodinium rotundum* (Lebour,1922) Schiller,1937
= *Peridinium limnophilum* Lindemann,19??
* Warm water species.
All Baltic, Kattegatt, E North Sea, all around Britain
- Zygapikodinium** Loeblich jr. & Loeblich III,1970
= *Diplopeltopsis* Pavillard,1913
= *Dubridinium* Reid (cyst)
- Z. lenticula** Loeblich Jr. & Loeblich III,1970 (TM)
= *Diplopsalis lenticula* f. minor Paulsen,1907
= *Peridinium lenticulum* Mangin,1911
= *Glenodinium lenticula* f. minor (Paulsen,1907) Pavillard,1913
= *Diplopeltopsis minor* (Paulsen,1907) Pavillard,1913
All Baltic except Bothnian Bay, Öresund, SW Kattegatt, Limfjorden, E North Sea, around Britain
- Z. pseudoblea** Elbrächter,1975
List/Sylt, E North Sea
- Protoperidiinidae
- Protoperidinium** Bergh,1881
= *Peridinium* Ehrenberg,1831 (p.p.)
= *Archaeoperidinium* Jørgensen,1912
= *Congruentidium* Abé,19??
= *Minuscula* Lebour,1925
= *Propertidinium* Meunier,19
- P. (Archaeoperidinium) avellana** (Meunier,1919) Balech,1974
= *Propertidinium avellana* Meunier,1919
Bothnian Bay - N Baltic, Gulf of Finland, Belgium, Plymouth, Woods Hole
- P. (Archaeoperidinium) denticulatum** (Gran & Braarud,1935) Balech,1974
= *Peridinium denticulatum* Gran & Braarud,1935
= *Peridinium clavus* Abe,1936
Kattegatt, Norway, Helgoland, E North Sea, N Scotland
- P. (Archaeoperidinium) excentricum** (Paulsen,1907) Balech,1974
= *Peridinium excentricum* Paulsen,1907
Öresund, N Kattegatt, E North Sea, all around Britain, Mediterranean
- P. (Archaeoperidinium) laticeps** (Grøntved & Seidenfaden,1938)

Kattegatt, W Greenland

P. (Archaeoperidinium) Jörgensen,1912 minutum (Kofoid,1907) Loeblich III,1969
 = *Peridinium minutum* Kofoid,1907
 = *P. monospinum* Paulsen,1907
 SW Kattegatt, Limfjorden, Skagerrak, E North Sea, all around Britain, Atlantic

P. (Archaeoperidinium) Jörgensen,1912 thorianum (Paulsen,1905) Balech,1973
 = *Peridinium thorianum* (Paulsen,1905) Balech,1973
 Limfjorden, Kattegatt, Skagerrak, Oslofjord, Norway, E North Sea, Faeroes, Iceland, all around Britain

P. (Minusculum) Lebour,1925 bipes (Paulsen,1904) Balech,1974
 = *Glenodinium bipes* Paulsen,1904
 = *Peridinium minusculum* Pavillard,1905
 = *Minuscula bipes* (Paulsen,1904) Lebour,1925
 all Baltic except Bothnian Bay, Öresund, Kattegatt, Skagerrak, Oslofjord, E North Sea, all around Britain, Iceland, Greenland, Mediterranean

P. (Protoperidinium) depressum (Bailey,1855) Balech,1974
 = *Peridinium depressum* Bailey,1855
 Arkona Sea, W Baltic, Belt Sea, Öresund, Kattegatt, Skagerrak, Oslofjord, E North Sea, all around Britain, widespread fr. Arctic to Antarctic

P. (Protoperidinium) mariebelouriae (Paulsen,1930) Balech,1974
 = *Peridinium mariebelouriae* Paulsen,1930
 = *Peridinium obtusum* : (Karsten) Lebour,1925, non *P. obtusum* (Karsten) Faure-Fremiet,1980, non *P. divergens obtusum* Karsten,1906
 Limfjorden, Oslofjord, E North Sea, all around Britain

P. (Protoperidinium) oceanicum (Vanhöffen,1897) Balech,1974
 = *Peridinium oceanicum* Vanhöffen,1897
 Belt Sea, N Kattegatt, Skagerrak, Oslofjord, off N, NW & SW Britain, off Iceland - Labrador, Mediterranean

P. (Protoperidinium) oblongum (Aurivillius,1898) Parke & Dodge, in Parke & Dixon,1976
 = *P. divergens* Ehrenberg, var. *oblongum* Aurivillius,1898
 Öresund, Kattegatt, Skagerrak, E North Sea, all around Britain

P. (Protoperidinium) saltans (Meunier,1910) Balech,1973
 = *Peridinium saltans* Meunier,1910
 North Sea, Barents Sea, off N & W Britain, Atlantic

P. (Protoperidinium) claudicans (Paulsen,1907) Balech,1974
 = *Peridinium claudicans* Paulsen,1907
 off S Fyn, Öresund, Limfjorden, Skagerrak, E North Sea, all around Britain, coastal waters of Europe & N America

P. (Protoperidinium) punctulatum (Paulsen,1907) Balech,1974
 = *Peridinium punctulatum* Paulsen,1907
 = *Peridinium submersum* Paulsen, var. *punctulatum* (Paulsen) Schiller,1937
 Öresund, N Kattegatt, Skagerrak, S & E North Sea, all around Britain

P. (Protoperidinium) achromaticum (Levander,1902) Balech,1974
 = *Peridinium achromaticum* Levander,1902
 all Baltic, E North Sea, all around Britain

P. (Protoperidinium) conicoides (Paulsen,1905) Balech,1973
 = *Peridinium conicoides* Paulsen,1905
 W Baltic, Öresund, Limfjorden, Kattegatt, Skagerrak, Oslofjord, Norway, E North Sea, Iceland, Greenland, Arctic, all around Britain

P. (Protoperidinium) conicum (Gran,1900) Balech,1974
 = *Peridinium divergens* var. *conicum* Gran,1900
 Belt Sea, Öresund, Kattegatt, Skagerrak, Oslofjord, E North Sea, all around Britain, Mediterranean

P. (Protoperidinium) leonis (Pavillard,1916) Balech,1974
 = *Peridinium leonis* Pavillard,1916
 off S Fyn, E North Sea, all around Britain, Atlantic, Mediterranean

P. (Protoperidinium) obtusum (Karsten,1906) Parke & Dodge, in Parke & Dixon,1976
 = *Peridinium divergens obtusum* Karsten,1906, non *P. obtusum* Faure-Fremiet,1980
 E North Sea, Plymouth

P. (Protoperidinium) pentagonum (Gran,1902) Balech,1974
 = *Peridinium pentagonum* Gran,1902
 = *Peridinium sinuosum* Lemmermann,1905
 Kieler Bucht, Öresund, Limfjorden, Kattegatt, Skagerrak, E North Sea, all around Britain

P. (Protoperidinium) subinerme (Paulsen,1904) Loeblich III,1969
 = *Peridinium subinermis* Paulsen,1904
 N Kattegatt, Skagerrak, Oslofjord, E North Sea, Barents Sea, Kara Sea, Iceland, Greenland, all around Britain

P. (Protoperidinium) brevipes (Paulsen,1908) Balech,1974
 = *Peridinium brevipes* Paulsen,1908
 = *Peridinium varicans* Paulsen,1911
 = *Peridinium incurvum* Lindemann,1924
 all Baltic except Bothnian Bay, Öresund, Kattegatt, Skagerrak, Oslofjord, off Norway, E North Sea, all around Britain

P. (Protoperidinium) cerasus (Paulsen,1907) Balech,1973
 = *Peridinium cerasus* Paulsen,1907
 = *Peridinium quarnnerense* Schiller,1937 (p.p.)
 Öresund, Limfjorden N Kattegatt, Skagerrak, all around Britain

P. (Protoperidinium) curtipes (Jörgensen,1912) Balech,1974
 = *Peridinium curtipes* Jörgensen,1912
 = *Peridinium crassipes* Paulsen,1907 (p.p.), non : Paulsen,1930, nec : Schiller,1937
 Kattegatt, all around Britain

P. (Protoperidinium) crassipes (Kofoid) Balech,1974
 = *Peridinium crassipes* Kofoid,1974
 Öresund, Kattegatt, Limfjorden, Skagerrak, Oslofjord

P. (Protoperidinium) decipiens (Jörgensen,1899) Parke & Dodge, in Perke & Dixon,1976
 = *Peridinium decipiens* Jörgensen,1899
 W Belt Sea, Öresund, Skagerrak, W Norway, Arctic Seas, Britain

P. (Protoperidinium) divergens (Ehrenberg,1840) Balech,1974
 = *Peridinium divergens* Ehrenberg,1840
 Arkona Sea, W & 'S Baltic, Öresund, Kattegatt, Skagerrak, Oslofjord, E North Sea, N Scotland, SW England

P. (Protoperidinium) globulum (von Stein,1883) Balech,1974
 = *Peridinium globulus* von Stein,1883
 North Sea, Engl. Channel, Atlantic, Mediterranean

P. (Protoperidinium) ovatum Pouchet,1883
 = *Peridinium ovatum* (Pouchet,1883) Schütt,1???

W Belt Sea, Öresund, Kattegatt, Skagerrak, Oslofjord, E North Sea, all around Britain, Atlantic, Mediterranean

P. (Protoperidinium) subcurvipes (Lebour,1923) Balech,1974
 = *Peridinium subcurvipes* Lebour,1923
 off NE England, Engl. Channel, Spitzbergen

P. (Protoperidinium) granii (Ostenfeld,1906) Balech,1974

= *Peridinium granii* Ostenfeld,1906

= *Peridinium finlandicum* Paulsen,1907

= *Peridinium divergens* var. *levanderi* Lemmermann,19??

* Cold water species.

all Baltic, Öresund, Kattegatt, Skagerrak, Oslofjord, North Sea, Engl. Channel, W Britain, off Iceland

P. (Protoperidinium) mite (Pavillard,1916) Balech,1974

= *Peridinium mite* Pavillard,1916

all around Britain, Atlantic, Mediterranean

P. (Protoperidinium) pyriforme (Paulsen,1905) Balech,1974

= *Peridinium steini* var. *pyriformis* Paulsen,1905

Öresund, S Kattegatt, Limfjorden, Skagerrak, Oslofjord, Norway, E Greenland, all around Britain

P. (Protoperidinium) pyriforme (Paulsen,1905) Balech,1974 ssp. *breve* (Paulsen,1905) Balech,1971

W & 'S Baltic, Öresund

P. (Protoperidinium) steini (Jörgensen,1900) Balech,1974

= *Peridinium steini* Jörgensen,1900

= *Peridinium michaelis* von Stein,1883 (p.p.)

Arkona Sea, W Baltic, Öresund, Kattegatt, Skagerrak, Oslofjord, E North Sea, all around Britain, Atlantic

P. (Protoperidinium) curvipes (Ostenfeld,1906) Balech,1974

= *Peridinium curvipes* Ostenfeld,1906

Gdansk, W Baltic, Limfjorden, Skagerrak, E North Sea, all around Britain, Greenland, Florida

P. (Protoperidinium) diabolum (Cleve,1900) Balech,1974

= *Peridinium diabolus* Cleve,1900

Oslofjord, all around Britain, Atlantic, Mediterranean

P. (Protoperidinium) islandicum (Paulsen,1904) Balech,1973

= *Peridinium islandicum* Paulsen,1904

Skagerrak, off N Britain, Iceland, Spitzbergen, Greenland

P. (Protoperidinium) pallidum (Ostenfeld,1899) Balech,1973

= *Peridinium pallidum* Ostenfeld,1899

W Belt Sea, Öresund, Kattegatt, Skagerrak, Oslofjord, E North Sea, all around Britain, Arctic Seas, Atlantic, Mediterranean

P. (Protoperidinium) pellucidum Berg,1881 (TM)

= *Peridinium pellucidum* (Berg,1881) Schütt,1???

all Baltic, Öresund, Kattegatt, Skagerrak, North Sea, Oslofjord, all around Britain, off Greenland, Atlantic, Mediterranean

P. (Protoperidinium) thulese (Balech,1958) Balech,1973

= *Peridinium thulese* Balech,1958

= *Peridinium conicum* f. *istandica* Braarud,1935

= *Peridinium symphyla* Hermosilla & Balech,1969

off NE Scotland, Engl. Channel, off Greenland

P. (Protoperidinium) deficiens (Meunier,1919) Balech,1974

= *Peridinium deficiens* Meunier,1919

W & S Baltic, E North Sea

P. (Protoperidinium) longispinum (Kofoid,1907) Balech,1974

= *Peridinium longispinum* Kofoid,1907

= *Peridinium michaelis* Ehrenberg,1840 (p.p.)

Greifswald, W Baltic, Belt Sea

P. (Protoperidinium) quarnerense (Schröder)

= *Peridinium globulus* var. *quarnerense* Schröder

Oslofjord, E North Sea

P. (Protoperidinium) brochii (Kofoid & Swezy,19??) Balech,19???

?Kattegatt

P. (Protoperidinium) sp. A : Thomsen & al., 1992

= *Peridinium paulsenii* Abé,1981, non Pavillard,1905

Kattegatt

P. (Protoperidinium) sp. B : Thomsen & al., 1992

= *P. pallidum* : Dodge,1985, p. 61, non (Ostenfeld,1899)

Kattegatt

P. (Protoperidinium) sp. C : Thomsen & al., 1992

Kattegatt

P. bispinum (Schüller,1???) Balech,1???

N Scotland

P. tuba (Schüller,1???) Balech,1???

N Scotland

P. roseum (Paulsen) Balech,1974?

= *Peridinium roseum* Paulsen,1???

Öresund, N Kattegatt, Skagerrak

Chalubinskia ??,1???

* Genus not recognized by Sournia,1986.

* Not mentioned by Larsen & Sournia,1992, in Patterson & Larsen

* Correct family?

Podolampadaceae Lindemann,1928

Blepharocysta Ehrenberg,1873

B. splendormaris (Ehrenberg,1860) Ehrenberg,1873 (TM)

= *Peridinium splendor-maris* Ehrenberg,1860

off Plymouth, warm Atlantic

B. paulsenii Schiller,1937

off SW Britain, Atlantic, Mediterranean

Podolampas von Stein,1883 (TM *P. bipes* von Stein,1883?)

= *Parrocella* Gourret

P. palmipes von Stein,1883

Skagerrak, occasionally N Atlantic, warm Atlantic

Dissodium Abe,1941 (TM *D. parvum* Abe,1941)

* Genus not recognized by Sournia,1986, but by Taylor,1987.

* Not mentioned by Larsen & Sournia,1992, in Patterson & Larsen

D. assymetricum (Mangin,1???) Loeblich,19???

N Scotland

Thoracosphaeraceae

* Not mentioned by Larsen & Sournia,1992, in Patterson & Larsen

Thoracosphaera ??,1???

* Not mentioned by Larsen & Sournia,1992, in Patterson & Larsen

Pyrophacaceae Lindemann,1928

- Fragilidium** Balech, ex Loeblich III, 1965 (TM) *F. heterolobum* Balech, ex Loeblich III, 1965
= *Helgolandinium* von Stosch, 1969
- F. subglobosum** (von Stosch, 1969) Loeblich, 19??
= *Helgolandinium subglobosum* von Stosch, 1969 (TM of *Helgolandinium* von Stosch, 1969)
Limfjorden, Kattegatt, E North Sea, around Britain
- Pyrophacus** von Stein, 1883
= *Tuberculodinium* Wall (cyst)
- P. horologicum** von Stein, 1883 (TM)
Gulf of Finland, Belt Sea, Kattegatt, Skagerrak, Oslofjord, E North Sea, around Britain
- Peridiniales "incertae sedis"**
- Acanthogonyaulax** (Kofoid) Graham, 1942 (TM) *A. spinifera* (Murray & Whitting) Graham, 1942 - tropical seas
* Genus not recognized by Taylor, 1987, but by Sournia, 1986.
- Adenoides** Balech, 1956
* Genus not recognized by Taylor, 1987, but by Sournia, 1986.
- A. elutens** (Herdman, 1922) Balech, 1956 (TM)
= *Amphidinium elutens* Herdman, 1922
* Sand dweller.
Isle of Man, Norfolk, Bretagne, River Conway estuary
- A. kofoidi** (Herdman, 1922) Dodge, 1982
= *Amphidinium kofoidi* Herdman, 1922
= *Phalacroma kofoidi* : Herdman, 1924
non = *Amphidinium kofoidi* var. *petasatum* Herdman, 1924
* Sand dweller.
Isle of Man
- Ceratoperidinium** Margalef, ex Loeblich III, 1980 (TM) *C. yeye* Margalef, ex Loeblich III, 1980 - Mediterranean
* Genus not recognized by Taylor, 1987, but by Sournia, 1986.
- Dinospaera** ??,1???,
* Genus not recognized by Sournia, 1986.
- Glenodiniopsis** ??,1???
- Halophilodinium** Loeblich Jr. & Loeblich III, 1968
- Hollandella** Cachon, 1964
* Correct suborder? Which family?
* Not mentioned by Larsen & Sournia, 1992, in Patterson & Larsen
- H. mycetoides** ?Cachon, 1964?
@ the radiolarian *Spongospaera* sp.
Mediterranean
- H. lobata** ?Cachon, 1964?
@ the radiolarian *Plegnospaera coronata*
Mediterranean
- H. piriformis** ?Cachon, 1964?
@ the radiolarian *Actinosphaera* sp.
Mediterranean
- Micracanthodinium** Deflandre, 1937
- M. setiferum** (Lohmann, 1902) Deflandre, 1937 (TM)
= *Cladopyxis setifera* Lohmann, 1902
S Baltic, Belt Sea, N Atlantic, Adriatic
- M. claytonii** (Holmes, 1956) Dodge, 1982
= *Cladopyxis claytonii* Holmes, 1956
Arkona Sea, Oslofjord, off Norway, Labrador
- Microceratium** Sournia, 1972 (TM) *M. orstomi* Sournia, 1972 - Mozambique Canal
* Genus not recognized by Taylor, 1987, but by Sournia, 1986.
- Spiraulax** Kofoid, 1911 (TM) *S. jollifei* (Murray & Whitting) Kofoid, 1911 - tropical seas
= *Spiraulaxina* Loeblich III, 1970
- KOLKWITZIELLALES**
* Not mentioned by Larsen & Sournia, 1992, in Patterson & Larsen
- Kolkwitziellaceae**
* Not mentioned by Larsen & Sournia, 1992, in Patterson & Larsen
- Berghiella** Kofoid & Michener, 1911
* Regarded as a doubtful dinoflagellate by Sournia, 1986.
* Not mentioned by Larsen & Sournia, 1992, in Patterson & Larsen
- Kolkwitziella** ??,1???,
* Not mentioned by Larsen & Sournia, 1992, in Patterson & Larsen
?
- BLASTODINIALES** Schiller, 1937
* Not mentioned by Larsen & Sournia, 1992, in Patterson & Larsen
- Apodiniaceae** = **Apodinidae**
- Apodinium** Chatton, 1907 (TM) *A. mycetoides* Chatton, 1907 - on gill interstitia of *Fritillaria pellucida* - Mediterranean
= *Parapodinium* Chatton, 1920
@ Appendicularians
* one other sp., *A. rhizophorum* Chatton, 1912 - on the tail of *Oikopleura cophocerca* -, is also known from the Mediterranean
- A. chattoni** Cachon & Cachon, 19??
= *Parapodinium chattoni* & *Oikopleura fusiformis*
@ Appendicularia sicula
Mediterranean
- A. rhizophorum** Cachon & Cachon, 19??
= *Parapodinium catenaria*
@ Appendicularia sicula & *Fritillaria cophocerca*
?
- A. sp.**
= *Parapodinium stylipes* Chatton, 1920? (TM of *Parapodinium* Chatton, 1920)
@ *Oikopleura formica* & *O. dioica*
S France
- Blastodiniaceae**
= **Blastodinidae**
- Blastodinium** Chatton, 1906
- * in alimentary channel of copepods
- B. hyalinum** Chatton, 1911
= *B. contortum* *hyalinum* Chatton, 1911
@ *Paracalanus parvus* Claus, *Centropages typicus* Krøyer, *Calanus finmarchicus* (Gunnerus), *Acartia clausi* Giesbrecht, *Pseudocalanus elongatus* & *Clausocalanus* sp.
* the only species without colour
Kattegatt, North Sea, Roscoff, Mediterranean
- B. spinulosum** Chatton, 1908
@ *Paracalanus* spp., *Clausocalanus* spp., *Acrocalanus gracilis* Giesbrecht
S France
- B. crassum** Chatton, 1908
@ *Paracalanus parvus* Claus & *Clausocalanus* spp.
S France
- B. pruvoti** Chatton, 1906
@ *Paracalanus parvus* Claus, *Acartia clausi* Giesbrecht & *Clausocalanus* spp., etc.
S France
- B. contortum** Chatton, 1908
@ *Paracalanus parvus* Claus, *Acartia clausi* Giesbrecht & *Clausocalanus* spp., etc.
S France
- B. oviforme** Chatton, 1912
@ *Oithona* spp., *Corycaeus* spp.
?
- B. inornatum** Chatton, 1920
@ *Clausocalanus arcuicornis* (Dana), *C. furcatus* (Brady), *Paracalanus parvus* (Claus), *P. aculeatus* Giesbrecht
?
- B. mangini** Chatton, 1908
@ *Corycelia gibbulus* Giesbrecht, *C. rostrata* Farran, *Oncaea conferta* Giesbrecht, *O. media* Giesbrecht, *O. venusta* Philippi
?
- B. mangini oncae** Chatton, 1912
@ *Oncaea* spp. & *Corycelia rostrata* Claus
?
- B. navicula** Chatton, 1912
@ *Corycelia venustus* Dana
?
- B. elongatum** Chatton, 1912
@ *Scolecithrix bradyi* Giesbrecht
?
- B. apsteini** Sewell, 1951
@ *Clausocalanus arcuicornis* (Dana), *C. furcatus* (Brady), *Paracalanus aculeatus* Giesbrecht
?
- B. chattoni** Sewell, 1951
@ *Clausocalanus arcuicornis* (Dana), *C. furcatus* (Brady), *Nannocalanus minor* (Claus), *Paracalanus aculeatus* Giesbrecht, *P. denudatus* Sewell, *P. parvus* (Claus), *Cosmocalanus darwini* (Lubbock)
?
- Diplomorphidae**
= *Cachonellaceae*
= *Cachonellidae*
- Actinodinium** Chatton & Hovasse, 1938
- A. apsteinii** Chatton & Hovasse, 1938 (TM)
@ *Acartia clausi* Giesbrecht
Roscoff
- Cachonella** ??,1???,
= *Diplomorpha* Rose & Cachon, 1951, *non* Ancey, 1884 (Mollusca)
- C. paradoxa** (Rose & Cachon, 1951)
= *Diplomorpha paradoxa* Rose & Cachon, 1951 (TM of *Diplomorpha* Rose & Cachon, 1951)
@ *Abylopsis tetragona*, *Forskaalia contorta*, *Chelophys* sp. & other Siphonophores
W Mediterranean
- Haplozoaceae** = **Haplozoonidae**
- Haplozoon** Dogiel, 1907
@ Polychaeta
- H. armatum** Dogiel, 1907 (TM - Murmansk)
@ *Travisia forbesi*
Bergen, Murmansk
- H. lineare** Dogiel, 1907
@ *Nicomache lumbicalis* (O. Fabricius)
Murmansk
- H. delicatulum** Dogiel, 1910
@ a maldanid species
Trondheim
- H. tuberculatum** Dogiel & Mikelsson, 1923
@ a maldanid species
Trondheim
- H. ariciae** Dogiel, 1910
@ *Phylo norvegica* (M. Sars)
Trondheim
- H. villosum** Dogiel & Mikelsson, 1923
@ *Polypisia crassa*
Trondheim
- H. obscurum** Dogiel, 1910
@ *Terebellides stroemi*
Trondheim
- H. macrostylum** Dogiel, 1910
@ a maldanid species
Trondheim
- H. dogieli** Shumway, 1924
@ *Leiochone clypeata*
Roscoff
- H. clymenellae** (Calkins, 1915)
@ *Clymenella torquata*
Woods Hole
- H. clymenidis** Dogiel & Mikelsson, 1923

- @ Euclymene palermiana*
Gulf of Naples
- H. inerme* Cachon,1964
@ Appendicularia sicula
?
- Oodiniaceae* Chatton,1920
= *Oodinidae*
- Oordinum* Chatton,1912
= *Salpicola* Bargoni,1894
@ ectoparasites on Appendicularians, Salps, Annelids & Molluscs
- O. pouchetii* (Lemmermann,1899) Chatton,1912 (TM)
= *Gymnodinium pulvisculum* Pouchet,1883
= *Salpicola amyloacea* Bargoni,1894
@ e.g. *Oikopleura dioica* Fol & *Salpa democratica* Forskål
Concarneau, Banyuls-sur-mer
- O. fritillariae* Chatton,1912
@ *Fritillaria pellucida* & ? *Fritillaria borealis*
?Davis Strait, Mediterranean
- O. acanthometrae* ?Cachon,1964?
@ *Acanthometra pellucida*
Mediterranean
- Amyloodinium* Brown & Hovasse,1946
@ fish
- A. ocellatum* (Brown,1931)
= *Oodinium ocellatum* Brown,1931 (TM)
@ causing "velvet disease" in various fish species
Virginia, Eilat, New York, Bermudas, West Indies, Danmarks Akvarium
- Crepidoodinium* Lom & Lawler,1981
= *Oodinium cyprinodontum* Lawler,1967 (TM)
@ different cyprinodontid fish species
Virginia
- Caryotoma* ??,19??
- C. bernardi* ??
@ endoplasm of radiolarian *Thalassicolla spumida*
?
- Protoodiniaceae*
= *Protoodinidae*
- Crepidoodinium*
- Piscinoodinium*
* with 2 known species from fresh water fishes.
- Protoodinium* Hovasse,1935
- P. chattoni* Hovasse,1935 (TM)
@ external parasite of *Lizzia blondina* ; occasionally on *Sarsia* sp. & *Podocoryne* spp.
Mediterranean, Bosporus
- P. hovassei* Cachon & Cachon,1964?
@ Siphonophores
?Mediterranean
- CHYTRIODINIALES**
- Chytriodiniaceae*
= *Chytriodinidae*
- Chytriodinium* Chatton,1912
@ eggs of pelagic copepods, euphausiids & shrimps
* 3 species known 1952 from not determined hosts
- C. roseum* (Dogiel,1906)
@ *Meganyctiphanes norvegica* (Sars)
?
- C. affine* (Dogiel,1906)
@ copepods
?
- C. parasiticum* (Dogiel,1906)
@ ?penaeids
?
- Syltordinum* Drebes,1989
- S. listii* Drebes,1989
@ eggs of *Temora*, *Acartia*, *Pseudocalanus* & *Synchaeta*
North Sea
- Dissodinium* Klebs, in Pascher,1916, emend. Elbrächter & Drebes,1978 (TM) *D. lunula* (Schütt) Klebs, in Pascher,1916
= *Diplodinium* Klebs,1912, non Fiorentini,1890 (Protostista)
= *Sporodinium* Gönnert,1936 (?) *Sporodinium* according to Neave
@ Copepod eggs
- D. pseudolunula* Swift, ex Elbrächter & Drebes,1978
= *Pyrocystis lunula* : Auctt., non (Schütt),1895
@ eggs of *Temora longicornis* (Müller), *Acartia* spp., rotifer cysts
Gdansk, W Baltic, Øresund, N Kattegatt, Skagerrak, Oslofjord, E North Sea
- D. pseudocalani* (Gonnert,1936) Drebes, ex Drebes & Elbrächter,1978
= *Sporodinium pseudocalani* Gonnert,1936
@ eggs of *Pseudocalanus elongatus* (Boeck)
Helgoland, N Scotland
- Myxodinium* Cachon, Cachon & Bouquahieux,1969
- M. pipiens* Cachon, Cachon & Bouquahieux,1969 (TM)
@ *Halosphaera viridis* & *Halosphaera minor*
Mediterranean
- Sporodinium*
- Schizochytriodinium* Elbrächter,1988
- S. calani* Elbrächter,1988 (TM)
- © eggs of *Calanus glacialis* Jashnov & *C. hyperboreus* Krøyer
Greenland Sea
- PARADINIOPHYCIDAE**
* Systematic placement rather uncertain.
- Paradinium* Chatton,1910
* genus incertae sedis. Systematic placement very tentative
- P. pouchetti* Chatton,1910
@ hemocoel of *Acartia* spp. & ? *Oithona plumifera*
?Kattegatt, Roscoff, Concarneau, Banyuls-sur-mer
- P. caulleryi* Chatton & Soyer,1973
@ *Oncaea media* Giesbrecht
?
- P. mesnilii* Chatton & Soyer,1973
@ *Oncaea conifera* Giesbrecht
?
- ELLOBIOPHYCIDAE**
- THALASSOMYCETALES**
- Thalassomyctaceae*
= *Ellobiopsidae* Coutière,1911
- Ellobiocystis* Coutière,1911
- E. catenatus* Coutière,1911
@ epibionts on 3rd maxillipeds of *Pasiphaea multidentata* Esmark & on *Acanthephyra purpurea* Milne-Edwards Oslofjord, Faeroes, S & W Ireland, NE Atlantic
- E. caridarum* Coutière,1911
@ *Acanthephyra purpurea* Milne-Edwards, *A. eximia* Smith, *Antarctomyces maxima* Hansen, *Pasiphaea pacifica* Rathbun, *P. semispinosa* Holthuis, *Sergestes* spp., *Systellaspis debilis* (Milne-Edwards) (and @ mouthparts of copopods (*Pseudaeidetus armatus* (Boeck), *Gaidius tenuispinus* G.O. Sars, *Gaidius brevispinus* G.O. Sars, *Gaetanus kruppii* Giesbrecht, *Gaetanus pileatus* Farran, *Gaetanus latifrons* G.O. Sars & *Pseudochirella notacantha* (G.O. Sars)) * Should probably refer to another species). NE Atlantic
- E. tenuis* Coutière,1911
@ *Pasiphaea sivado* (Risso)
?
- E. tuberosus* Coutière,1911
@ *Acanthephyra purpurea* Milne-Edwards
?
- Ellobiopsis* Caulfield,1910
- E. chattoni* Caulfield,1910 (TM)
@ *Calanus helgolandicus* Claus, *Calanus finmarchicus* (Gunnerus), *Pseudocalanus minutus* (Krøyer) & *Acartia clausii* Giesbrecht, *Clausocalanus arcuicornis* (Dana), *Ctenocalanus vanus* Giesbrecht, *Euchaeta marina* (Prestandrea), *E. wolfendeni* Scott, *Metridia longa* (Lubbock), *Pleuronema gracilis* (Claus), *P. borealis* (Dahl), *Pseudocalanus elongatus* (Boeck), *Cosmocalanus vulgaris* (Sewell), zoeae of *Portunus pelagicus* (L.) Baltic, Kattegatt, Bohusl, N North Sea, Clyde Sea area, Barents Sea
- E. elongata* Steuer, 1932
@ *Clausocalanus arcuicornis* (Dana), *Ctenocalanus vanus* (Giesbrecht), *Cosmocalanus darwini*, *Euchaeta marina* (Prestandrea), *Gaetanus antarcticus* Wolfenden, *G. curvicornis* Sars, postlarvae of *Penaeopsis retacuta* (Bate) ?
- E. fagei* Hovasse, 1951
@ pelagic copepods, *Clausocalanus arcuicornis* (Dana)
?
- Thalassomyces* Niezabitowski,1913
= *Staphylocystis* Coutière,1911, non Villot,1877 (Cestoda)
= *Amallocystis* Fage,1936
- T. fagei* (Boschma,1948)
= *Staphylocystis racemosus* : Macdonald,1927, non Coutière,1911
= *Amallocystis fagei* Boschma,1948
@ in centre of dorsal surface of carapace of Euphausiace (e.g. *Meganyctiphanes norvegica* (M. Sars), *Thysanoessa raschii* (M. Sars), *Thysanoessa inermis* (Krøyer), *Euphausia* spp. and several other genera) N North Sea, Clyde Sea area, Faeroe - Iceland area, NE Atlantic
- T. capillous* (Fage,1938)
= *Amallocystis capillous* Fage,1938
@ on rostrum of *Pasiphaea tarda* Krøyer, *P. multidentata* Esmark, *P. acutifrons* Bate & *P. pacifica* Rathbun Skagerrak, Faeroe - Shetland area, S & W Ireland
- T. racemosus* (Coutière,1911)
= *Staphylocystis racemosus* Coutière,1911
= *Thalassomyces batei* Niezabitowski,1913
@ on ventral surface of proximal half of abdomen of Caridea (*Pasiphaea tarda* Krøyer, *P. cristata* Bate, *Hymenodora glacialis* Buchholz, *Acanthephyra pelagica* (Risso), *A. sexspinosa* Kemp) Faeroe - Iceland area, NE Atlantic, Mediterranean
- T. fasciatus* (Fage,1936)
@ *Gnathophausia* spp., e.g. *G. zoea* Willemoes-Suhm
?
- T. albatrossi* Wing,1975
@ *Stiomysis major* Tattersall
?
- T. boschmai* (Nouvel,1954)
@ *Acanthomyysis macropsis* (Tattersall), *A. nephrophthalma* Banner, *A. pseudomacropsis* (Tattersall), *Gastromyzon lobatus* Nouvel, *Leptomyysis gracilis* (Sars), *Meterythrops robusta* Smith, *Neomysis kadiakensis* Ottmann
?
- T. marsupii* Kane,1964
@ in amphipods: *Cyphocaris challengerii* Stebbing, *Cystisoma* sp., *Eusirus leptocarpus* Sars, *E. longipes* Boeck, *Parathemisto abyssorum* (Boeck), *P. gaudichaudii* (Guerin), *P. gracilipes* Norman, *P. libellula* (Lichtenstein), *P. pacifica* Stebbing, *Rhachotropis aculeata* (Lepechin), *R. helleri* Boeck, *R. macropus* Sars
?
- T. niezabitowski* Hoenigman,1960
@ *Leptomyysis megalops* Zimmer
?
- T. nouveli* (Hoenigman,1954)
@ *Anchialina agilis* (Sars)
?
- T. spiczakovi* Niezabitowski,1913
* May possibly be a synonym of *T. racemosus* (Coutière,1911)
@ *P. sivado* (Risso)
?

- T. umbellatus** (Boschma,1949)
@ Oplophoridae: *Oplophorus gracilirostris* Milne-Edwards, *O. grimaldii* Coutière, *O. novaezealandiae* de Man?
- T. californiensis** Collard,1966
@ *Pasiphaea emarginata* Rathbun California
- T. sp.**
@ *Siriella clausi* Sars?
?
- Parallobiopsis** Collin,1913
@ pelagic Crustacea
- P. coutieri** Collin,1913 (TM)
@ *Nebalia bipes* Fabricius?
- Rhizellobiopsis** Hovasse,1926
- R. eupraxiae** (Zachs,1923)
@ on parapodes of *Nephthys ciliata* Black Sea
- SYNDINIOPHYCIDAE**
- COCCIDINIALES**
- Amoebophryaceae** = Amoebophryidae
- Amoebophrya** Koeppen,1894
= *Hyalsoccus* Koeppen,1903
@ Dinoflagellata, apostome & tintinnid ciliates, Acantharia & Radiolaria (*Sticholonche*)
- A. acanthometrae**? Koeppen,1894?
@ several species of Acantharian family Acanthometridae Mediterranean
- A. sticholonchae**? Koeppen,1894?
@ Colonial radiolarian *Sticholonche zanklea* Hertwig,1877?
- A. ceratii** (Koeppen,1903)
= *Hyalsoccus ceratii* Koeppen,1903
@ Dinoflagellata e.g. *Ceratium* spp. & *Polykrikos schwartzii* & other Amoebophrya species, i.e. *A. acanthometrae* Helgoland, Mediterranean
- A. grassei**? Cachon,1964?
@ cytoplasm of *Oodinium poucheti* from *Oikopleura dioica* Mediterranean
- A. rosei**? Cachon,1964?
@ Foettingeriidae from the Siphonophor *Abylopsis tetragona* Mediterranean
- A. tintinni**? Cachon,1964?
@ *Xystonella lohmanni* Mediterranean
- Coccidiiniaceae**
- Coccidinium** Chatton & Biecheler,1934
@ Nuclear & cytoplasm parasites of other dinoflagellates
- C. duboscqui** Chatton & Biecheler,1934 (TM - Sète)
@ *Peridinium balticum* Lemmermann French Mediterranean
- C. legeri** Chatton & Biecheler,1934?
@ *Peridinium sociale* (Henneguy) French Mediterranean
- C. punctatum** Chatton & Biecheler,1934?
@ *Ostreopopsis mononis* (Meunier) French Mediterranean
- C. mesnilii** Chatton & Biecheler,1934?
@ *Cryptoperidinium foliaceum* von Stein French Mediterranean
- Dubosquellaceae** = Dubosquelliidae
- Dubosquella** Chatton,1920
@ Especially marine tintinnids, but also Dinoflagellata (e.g. *Noctiluca* & *Leptodiscus*)
- D. tintinnicola** (Lohmann,1908)
@ tintinnids, e.g. *Cytaroclysis*, *Tintinnopsis* & *Favella* Kieler Bucht, Bergen, Mediterranean
- D. anisospora** Grassé,19??
@ *Cytaroclysis*?
- D. aspida** Cachon,1964?
@ *Favella ehrenbergi*, *Coxiella lacinosa*, *Codonella campanula* & *Tintinnus fraknoi* Mediterranean
- D. cnemata** Cachon,1964?
@ *Favella ehrenbergi* Mediterranean
- D. melo** Cachon,1964?
@ perinuclear cytoplasm of *Noctiluca scintillans* Mediterranean
- D. nucleocola** Cachon,1964?
@ nucleus of *Leptodiscus medusoides* Mediterranean
- D. caryophaga** Cachon,1964?
@ *Strombidium*, *Strombilidium* & *Prorodon* Mediterranean
- Dubosquodinium** Grassé,1952?
@ Ciliata
- D. collini** Grassé, in Grassé,1952 (TM)
@ *Tintinnus fraknoi*?
- D. kofoidi** Grassé, in Grassé,1952
@ *Codonella campanula*?
?
- Dogelodinium**
@ Acantharia
- Keppenodinium**
= *Hollandella*
@ Radiolaria
- Spiromonadaceae** = Spiromonadidae Hollande, in Grassé,1952
- Spiromonas** Perty,1852 (TM *S. volubilis* Perty,1852 - not a flagellate, perhaps not even a prototist, according to Patterson & Zöllfel,1992)
- Sphaeriparaceae** = Sphaeriparidae
- Sphaeripara**
= *Neresheimeria*
@ Appendicularia
- Atlanticeclodinium** Cachon & Cachon-Enjumet,1965
@ Phaeodarea
- SYNDINIALES**
- Syndiniaceae** = Syndinidae
- Syndinium** Chatton,1912
@ Hemocoel of marine copepods & cavities of other invertebrates (appendicularians)
* 8 species known 1952
- S. turbo** Chatton,1912 (TM)
@ *Paracalanus parvus* Claus & ?*Calanus finmarchicus* (Gunnerus)
* The species in *Calanus* may be another.
?North Sea, Mediterranean
- S. minutum** Chatton,19??
@ *Paracalanus parvus* Claus?
?
- S. corycae** Chatton in Grasse,1952
@ *Corycaeus geisbrechti* Dahl?
?
- S. gammari** Manier, Fize & Grizel,1971
@ *Gammarus locusta* (L.)?
- S. borgerti**? Hollande, Enjumet & Manciet,1953?
@ *Aulacantha scolymantha*?
- Synhemidinium** Chatton,19??
- S. rostratum**? 19??
@ *Paracalanus parvus* Claus Mediterranean
- Actinodinium** Chatton & Hovasse,1938
- A. apsteini** Chatton & Hovasse,1938 (TM)
@ *Acartia clausa* Giesbrecht?
- Merodinium** Chatton,1923
@ in radiolarian cytoplasm
- M. brandti** Hovasse,1923 (TM)
@ intracapsular in cytoplasm of *Collozoum inerme* Müller Mediterranean
- M. vernale** Hovasse,19??
@ extracapsular in cytoplasm of *Collozoum inerme* Müller?
- M. insidiosum** Chatton,19??
@ cytoplasm of *Collozoum pelagicum* Haeckel?
- M. mendax** Chatton,19??
@ cytoplasm of *Myxosphaera caerulea* Haeckel?
- M. dolosum** Chatton,19??
@ cytoplasm of *Sphaerozoum punctatum* Müller?
- M. astatum** Chatton,19??
@ cytoplasm of *Sphaerozoum acuferum* Müller?
- M. belari**? 19??
@ cytoplasm of *Collozoum fulvum*?
- Hematodinium** Chatton & Poisson,1931
@ hemolymph of crabs
- H. perezii** Chatton & Poisson,1931
@ *Carcinus maenas*, *Lioxarcinus depurator*, *Cancer pagurus*, etc. Roscoff, Arcachon, Mediterranean
- H. australis** Hudson & Shields,1994
@ *Portunus pelagicus* (L.), *Scylla* sp., etc. Indopacific
- H.?** sp.
@ *Pandalus* spp., e.g. *P. borealis* Krøyer?
- Ichthyodinium** Hollande & Cachon,1952
- I. chabardi** Hollande & Cachon,1952
@ ova & freshly hatched larvae of *Sardina pilchardus* Mediterranean
- Solenodinium** Chatton,1913
@ in radiolarian nuclei
- S. fallax** Chatton,1913? (TM)
@ intranuclear in *Thalassicolla spinula* (Haeckel)

| | |
|---|--|
| Gulf of Naples | <i>B. gallica</i> Bourrelly,19 Parke & Dixon,1976 area |
| <i>Trypanodinium</i> Chatton,1912 @ Copepod eggs | <i>B. hooglandii</i> Nicolai & Baas-Becking,1935 (TM) Parke & Dixon,1976 area |
| <i>T. ovicola</i> Chatton,1912 (TM) @ intracytoplasmic in eggs of <i>Oncae media</i> Giesbrecht, <i>Oithona</i> spp., <i>Clytemnestra</i> spp. S France | <i>Chrysobiotriella</i> Strand,1926 |
| <i>Cochlosyndinium</i> Chatton,19?? @ <i>Coryceus</i> | <i>C. spondylorum</i> (Conrad,1926) Strand,1926 (TM) Belgium, Britain |
| DOUBTFUL DINOFAGELLATES (sensu Sournia,1986) | <i>Entodesmis</i> Borzi,1892 * Not mentioned in Chrétiennot-Dinet,1990 |
| <i>Archaeosphaerodiniopsis</i> Rampi,1943 | <i>E. litoralis</i> (Anand,1937) Parke & Green, in Parke & Dixon,1976 = <i>Gloeochrysis litoralis</i> Anand,1937 Parke & Dixon,1976 area |
| <i>Dinoasteromonas</i> Bursa,1971 | <i>E. maritima</i> (Anand,1937) Parke & Green, in Parke & Dixon,1976 = <i>Gloeochrysis maritima</i> Anand,1937 Parke & Dixon,1976 area |
| <i>Dinoceras</i> Schiller,1931 | <i>Monas</i> O.F. Müller,1773 = <i>Spumella</i> Cienkowski,1870 (Suggestion about proposal for name conservation from Preisig, Vørs & Hälfors,1992, in Patterson & Larsen) (TM) <i>Spumella vulgaris</i> Cienkowski,1870 - fresh water) |
| <i>Goniodinium</i> Dangeard,1927 | <i>M. uniguttata</i> Skuja,1939 Finland |
| <i>Melanodinium</i> Schiller,1935 | <i>M. dangeardii</i> Lemmermann,19?? Plymouth |
| <i>Nephrodinium</i> Meunier,1910 | <i>M. sociabile</i> Meyer,1??? |
| <i>Pachydinium</i> Pavillard,1915 | <i>Chromulinaceae</i> Engler, in Engler & Prantl,1900 |
| <i>Pseudoactiniscus</i> Bursa,1969 | <i>Chromulina</i> Cienkowski,1870 (TM) <i>C. nebulosa</i> Cienkowski,1870 - fresh water) |
| <i>Thaurilens</i> Pavillard,1917 | <i>C. pleiades</i> Parke,1949 Isle of Man |
| <i>Atelodinium</i> Chatton,1920 | <i>C. lunaris</i> Carter,19 Parke & Dixon,1976 area |
| <i>A. microsporum</i> Chatton,1920 @ coelom parasite of <i>Paracalanus</i> Banyuls-sur-mer | <i>C. ovalis</i> Klebs,18 Parke & Dixon,1976 area |
| <i>A. parasiticum</i> Chatton,1920 @ coelom parasite of <i>Paracalanus</i> Banyuls-sur-mer | <i>C. pallida</i> Perty Parke & Dixon,1976 area |
| <u>CHRYSONADIDA</u> = <u>CHRYSOPHYCEAE</u> Christensen,1962, non Pascher,1914 * All Pankow,1990 included & all Parke & Green, in Parke & Dixon,1976 included. This taxon belongs to the informal group Straminopiles Patterson,1989. | <i>C. parvula</i> Conrad,19 Parke & Dixon,1976 area |
| <i>Bordnamonas</i> Larsen & Patterson,1990 * Correct order? Which suborder & family? | <i>Oikomonas</i> Kent,1880 (TM) <i>O. mutabilis</i> Kent,1880-82? - fresh water) * Colourless |
| <i>B. tropicana</i> Larsen & Patterson,1990 (TM) Tåvråmine, Denmark, Wadden Sea, England, Ireland, N Atlantic, Belize, Panama, Brazil, Fiji, Australia | <i>O. termo</i> (O.F. Müller,1773) Kent,1880-82 = <i>Monas termo</i> O.F. Müller,1773 * Fresh, Brackish & Marine waters Plymouth |
| Palisporomonadidae de Saedeleer,1946 * Correct order? Which suborder? | <i>Sphaleromantis</i> Pascher,1910 (TM) <i>S. ochracea</i> Pascher,1910 |
| <i>Palisporomonas</i> de Saedeleer,1946 | <i>S. subsalsus</i> Conrad,19 * Systematic position uncertain. Parke & Dixon,1976 area |
| <i>P. apodium</i> de Saedeleer,1946 @ the diatoms <i>Actinocyclus subtilis</i> , <i>Striatella</i> sp. & <i>Licmophora</i> sp. Mediterranean | <i>S. marina</i> Pienaar,19 E North Sea |
| <u>CHROMULINALES</u> = <u>OCHROMONADALES</u> Pascher,1910 | <i>Chrysoplasma</i> Pascher,1910 (TM) <i>C. fenestrata</i> Pascher,1910 - fresh water) = <i>Pascherella</i> Conrad,1926 |
| <u>Ochromonadaceae</u> Lemmermann,1899 | <i>C. yserensis</i> (Conrad,1926)? = <i>Pascherella yserensis</i> Conrad,1926? Parke & Dixon,1976 area |
| <i>Dendromonas</i> Stein,18?? = <i>Monadodendron</i> Pascher,1942 | <i>Dinobryonaceae</i> Ehrenberg,1834 |
| <i>D. latypes</i> (Valkanov,19??) Preisig, Vørs & Hälfors, in Patterson & Larsen,1992 = <i>Monadodendron latipes</i> Valkanov,1970 * Brackish water | <i>Codonoea</i> James-Clark,1867 |
| <i>Ochromonas</i> Vysotskii,1887 (TM) <i>O. triangulata</i> Vysotskii,1887 - fresh water) = <i>Chromophyton</i> Woronin | <i>C. costata</i> James-Clark,1867 (TM) * Marine species ? |
| <i>O. olivacea</i> Büttner,1911 Kieler Bucht | <i>Dinobryon</i> Ehrenberg,1835 |
| <i>O. minima</i> Thronsdæn,1969 Oslofjord, Bergen, Tromsö | <i>D. petiolatum</i> Willén,1963 * Warm water species. All Baltic, Öresund, Kattegatt, Parke & Dixon,1976 area |
| <i>O. bourrellyi</i> Magne,19?? Parke & Dixon,1976 area | <i>D. sueicum</i> Lemmermann,1904 * Freshwater species, which does not tolerate the full salinity of the area. Gulf of Finland, Bothnian Bay, S Norway |
| <i>O. cosmopolita</i> Ruinen,19?? Parke & Dixon,1976 area | <i>D. bavaricum</i> Imhof,1890 * Freshwater species, which does not tolerate the full salinity of the area. Gulf of Finland, Bothnian Bay & Sea |
| <i>O. crenata</i> Klebs,1??? Parke & Dixon,1976 area | <i>D. divergens</i> Imhof,1890 * Freshwater species, which does not tolerate the full salinity of the area. Bothnian Bay & Sea, Gulf of Finland, German Baltic, Öresund, N Kattegatt |
| <i>O. minuscula</i> Conrad,19?? Parke & Dixon,1976 area | <i>D. sociale</i> Ehrenberg,1832 * Freshwater species, which does not tolerate the full salinity of the area. Gulf of Finland, German Baltic |
| <i>O. oblonga</i> Carter,19?? Parke & Dixon,1976 area | <i>D. sertularia</i> Ehrenberg,1835 (TM) * Freshwater species, which does not tolerate the full salinity of the area. Gulf of Finland, Lübeck |
| <i>Uraglena</i> Ehrenberg,1835 = <i>Uroglonopsis</i> Lemmermann | <i>D. balticum</i> (Schütt,1896) Lemmermann,1900 = <i>Dinodendron balticum</i> Schütt,1896 = <i>Dinobryon pellucidum</i> Levander,1??? All Baltic, Öresund, Kattegatt, Skagerrak, Parke & Dixon,1976 area |
| <i>U. americana</i> Calkins,1892 * Warm water species. Bothnian Bay & Sea, Gulf of Finland, N Baltic | <i>D. cylindricum</i> Imhof,1890 * Freshwater species, which does not tolerate the full salinity of the area. Bothnian Bay |
| <i>U. volvox</i> Ehrenberg,1835 (TM) * Freshwater species, which does not tolerate the full salinity of the area. Gulf of Finland | <i>Ollicola</i> Vørs,1992 = <i>Calycomonas</i> : Auctt., non Lohmann,1908 |
| <i>U. marina</i> Büttner,1911 * Possibly identical with <i>U. volvox</i> Ehrenberg, sensu Huber-Pestalozzi,1941. Kieler Bucht | |
| <i>Anthophysa</i> Bory de St. Vincent,1823 * Not mentioned in Chrétiennot-Dinet,1990 | |
| <i>A. vegetans</i> (O.F. Müller,1786) von Stein,1878 = <i>Volvox vegetans</i> O.F. Müller,1786 Gulf of Finland, German Baltic | |
| <i>Boekloewia</i> Nicolai & Baas-Becking,1935 | |

- O. vangorii* (Conrad,1938) Vørs,1992
 = *Codonomonas vangorii* (Conrad,1938) Lund,1960
 = *Codonomonas Van Goorii* Conrad,1938 (TM)
 = *Calycomonas wulffii* Conrad & Kufferath,1954
 = *Calycomonas gracilis* : Lohmann, in Wulff,1916, non Lohmann,1908
 Bothnian Sea, Gulf of Finland, N Baltic, Arcona Sea, Kieler Bucht, Kattegatt, Oslofjord, North Sea, S, W & N Norway, Barents Sea, Belgium
- O. cylindrica* (Conrad & Kufferath,1954) Vørs,1992
 = *Codonomonas cylindrica* Conrad & Kufferath,1954
 = *Calycomonas cylindrica* (Conrad & Kufferath,1954) Lund,1960
 Belgium
- O. dilatata* (Conrad & Kufferath,1954) Vørs,1992
 = *Codonomonas dilatata* Conrad & Kufferath,1954
 = *Calycomonas dilatata* (Conrad & Kufferath,1954) Lund,1960
 Belgium
- O. pascheri* (Van Goor,1925) Vørs,1992
 = *Codonomonas pascheri* Van Goor,1925
 = *Calycomonas pascheri* (Van Goor,1925) Lund,1960
 ?Netherlands
- Pseudokephyrion* Pascher,1913 (TM *P. undulatum* (Klebs) Pascher,1913 - fresh water)
- P. formosissimum* Conrad,19
 Parke & Dixon,1976 area
- Chrysococcaceae* Lemmerman,1889
- Chrysococcus* Klebs,1892
- C. rufescens* Klebs,1892 (TM)
 * Freshwater species, which does not tolerate the full salinity of the area.
 Gulf of Finland, Parke & Dixon,1976 area
- C. biporus* Skuja,1939
 * Freshwater species, which does not tolerate the full salinity of the area.
 Gulf of Finland
- C. minutus* (Fritsch,1918) Nygaard,1929
 * Freshwater species, which does not tolerate the full salinity of the area.
 = *Trachelomonas volvocina* f. *minuta* Fritsch,1916
 Bothnian Sea
- C. pascheri* Lohmann,19
 Parke & Dixon,1976 area
- C. cinctus* Lackey,19??
 Plymouth
- Kephryion* Pascher,1911 (TM *K. sitta* Pascher,1911)
- K. hemispaericum* (Lackey,1938) Conrad,1939
 = *Chrysococcus hemispaericum* Lackey,1938
 Gulf of Finland
- K. cinetum* (Lackey) Bourrelly,19
 Parke & Dixon,1976 area
- K. ovum* Pascher,19
 Plymouth
- K. petasatum* Conrad,19
 Parke & Dixon,1976 area
- Calycomonas* Lohmann,1908 (TM *Calycomonas gracilis* Lohmann,1908)
 = *Codonomonas* van Goor,1925
 * This genus is now a synonym of *Paulinella* Lauterborn,1895 (FILOSEA)
- C. globosa* Lohmann,19
 * Either a synonym of another species or should be transposed to another genus.
 Parke & Dixon,1976 area
- Angulochrysis* Lackey,1940 (TM *A. erratica* Lackey,1940)
- Paraphysomonadaceae* Preisig & Hibberd,1983
- Paraphysomonas* de Saedeleer,1929
 = *Lepidochromonas* Kristiansen,19
- P. antarctica* Takahashi,1987
 Tärminne, Öresund, Kattegatt, Limfjorden, Norway, E North Sea, England, Mediterranean, N midAtlantic, widespread
- P. circumvallata* Thomsen,19??
 * A fresh- & brackish water species
 ?
- P. corbidifera* Pennick & Clarke,19
 Parke & Dixon,1976 area
- P. cribrosa* Lucas,1968
 S Kattegatt, Bergen, E North Sea, Plymouth
- P. cylicophora* Leadbeater,1972
 Tärminne, S Kattegatt, Norway
- P. favoletata* Rees, Leedale & Cmiecik,19??
 Parke & Dixon,1976 area
- P. foraminifera* Lucas,1967
 Tärminne, W Baltic, Öresund, Kattegatt, Limfjorden, Bergen, Engl. Channel, Greenland, Italy
- P. imperforata* Lucas,1967
 Tärminne, Öresund, Limfjorden, Kattegatt, Britain, Greenland, N midAtlantic, cosmopolitan
- P. oligocycta* Takahashi,1987
 Tärminne, Arctic Canada, Antarctica
- P. siderophora* Thomsen,19??
 S Kattegatt, Parke & Dixon,1976 area
- P. subrotacea* Thomsen,19??
 * A fresh & brackish water species
 ?
- P. takahashii* Cronberg & Kristiansen,1980
- * Normally widespread in fresh waters of Northern hemisphere
 Tärminne
- P. vestita* (Stokes,1885) de Saedeleer,1929 (TM)
 = *Physomonas vestita* Stokes,1885
 * Freshwater species, which does not tolerate the full salinity of the area.
 Tärminne, Öresund, S Kattegatt, ?Bergen, cosmopolitan in fresh & brackish water
- Polylepidomonas* Preisig & Hibberd,1983
- P. vacuolata* (Thomsen) Preisig & Hibberd,1983 (TM)
 E North Sea
- Spiniferomonas* Takahashi,19??
- S. bourrellyi* Takahashi,1973
 = *Paraphysomonas bourrellyi* (Takahashi,1973) Hibberd & Preisig,1982
 Gulf of Finland
- Chromophysomonas* Preisig & Hibberd,1982
- C. tririalis* (Takahashi,1973) Preisig & Hibberd,1982
 = *Spiniferomonas tririalis* Takahashi,1973
 Gulf of Finland
- Mallomonadaceae* Diesing,1866
- Mallomonas* Perty,1852 (TM *M. ploesslii* Perty,1852 - fresh water)
- M. akrokomos* Ruttner, in Pascher,1913, emend. Bourrelly,1957
 * Freshwater species, which does not tolerate the full salinity of the area.
 Tärminne
- M. punctifera* Korschikov,1941
 = *M. reginae* Teiling,1946
 = *M. pulchella* (Kisselev,19??) Cronberg & Kristiansen,19??
 * Freshwater species, which does not tolerate the full salinity of the area.
 Tärminne
- M. tonsurata* Teiling,1912, emend. Krieger,1930
 * Freshwater species, which does not tolerate the full salinity of the area.
 Tärminne
- M. producta* Iwanoff,1899
 * Freshwater species, which does not tolerate the full salinity of the area.
 Gulf of Finland
- M. caudata* Iwanoff,1899
 * Freshwater species, which does not tolerate the full salinity of the area.
 Gulf of Finland
- M. epithalattia* Drost,19
 Parke & Dixon,1976 area
- M. subsalina* Conrad & Kufferath,1954?
 Parke & Dixon,1976 area
- PARMALES* Booth & Marchant,1987
- Triparmaeae* Booth & Marchant,1987
- Triparma* Booth & Marchant,1987 (TM *T. columacea* Booth & Marchant,1987)
- Tetraparma* Booth, in Booth & Marchant,1987 (TM *T. pelagica* Booth & Marchant,1987)
- Pentalaminiaceae* Merchant, in Booth & Marchant,1987
- Pentalamina* Merchant, in Booth & Marchant,1987 (TM *P. corona* Merchant, in Booth & Marchant,1987)
- CHRYSAMOEBALES* Matvienko,1965
- Chrysamoebaceae* Bourrelly,1957
- Chrysamoeba* Klebs,1892 (TM *C. radians* Klebs,1892 - fresh water)
- Rhizochrysinida* Hibberd & Chrétiennot-Dinet,1979 (TM *R. marina* Hibberd & Chrétiennot-Dinet,1979 - Mediterranean)
- Rhizochrysinidae* Pascher, ex Reichenow,1928
- Leukapsis* Pascher,1940 (TM *L. vorax* Pascher,1940)
- Stylococcaceae* Lemmermann,1899
- Chrysopodocystis* Billard,1978
- C. socialis* Billard,1978 (TM)
 Engl. Channel, Mediterranean
- Heliopsis* Pascher,1940 (TM *H. mutabilis* Pascher,1940 - fresh water)
- SARCINOCHRYSIDALES* Gayral & Billard,1977
- Sarcinochrysidaeae* Gayral & Billard,1977
- Sarcinochrysis* Geitler,1930
- S. marina* Geitler,1930 (TM)
 Parke & Dixon,1976 area
- Ankylochrysis* Billard, in O'Kelly & Billard,1991?
- A. lutea* (Van der Veer,1970) Billard, in O'Kelly & Billard,1991? (TM)
 = *Ankyloneton luteum* van der Veer,1970
 Parke & Dixon,1976 area
- ?Family?
- Actinomonas* Kent,1880
- A. mirabilis* Kent,1880 (TM)
 = *A. marina* Kufferath,19?? = *A. pusilla* Kent,18??
 = *A. radiosa* Roskin,19?? = *Pteridomonas scherffelii* Lemmermann,19??
 * Impossible to discriminate from *Pteridomonas danica* without EM thin sectioning
 ?Tärminne, Wadden Sea, Oslofjord, Bergen, England, France, ?midAtlantic, New Zealand, Brazil, Fiji, Hawaii
- Pteridomonas* Pénard,1890 (TM *P. pulex* Pénard,1890 - fresh water)
 * Like *Pedinella*, but without chromatophores

- P. danica* Patterson & Fenchel,1985
?Tvärminne, Denmark, Ireland, N midAtlantic, Brazil, Fiji, Hawaii, Australia
- P. salina* Bourrelly,19??
* Brackish & ?marine waters
?
- Ciliophryidae**
- Ciliophysa* Cienkowski,1875
- C. infusionum* Cienkowski,1876 (TM)
= *C. marina* Caulery,1910
Tvärminne, Denmark, Bergen, Germany, Engl. Channel, Mediterranean, SE North America, Fiji, Hawaii, Australia, European fresh water
- Pedinellaceae* Pascher,1910
- Pedinella* Wyssotzki,1887
- P. hexacostata* Wyssotzki,1887 (TM)
?Baltic, Plymouth
- Parapedinella* Pedersen & Thomsen, in Pedersen, Beech & Thomsen,1986
- P. reticulata* Pedersen & Thomsen, in Pedersen, Beech & Thomsen,1986 (TM)
Belt Sea, S & central Kattegatt
- Pseudopedinella* Carter,1937
- P. elastica* Skuja,1948
Gulf of Finland
- P. pyriforme* N. Carter,1937 (TM)
?Finland, S Kattegatt, Oslofjord, S Norway, Bergen, Tromsö, Belgium, Engl. Channel, Massachusetts
- P. tricostata* (Rouchijainen,1966) Thomsen,1986
= *Pedinella tricostata* Rouchijainen,1966
Gulf of Finland
- Apedinella* Thronsdsen,1971
- A. spinifera* (Thronsdsen,1969) Thronsdsen,1971 (TM)
= *Pseudopedinella spinifera* Thronsdsen,1969
Gulf of Finland, W Baltic, Öresund, Limfjorden, Kattegatt, Bohusl., Oslofjord, S Norway, Bergen, Arctic Seas, Mediterranean
- A. mediterranea* (Lohmann,1903)
= *Meringosphaera radians* Lohmann,1903
* Dubious species, sensu Ettl,1978. ?A synonyme of *Apedinella spinifera* (Thronsdsen)?
* Värs,1992 keeps it in *Meringosphaera* and think that it is related to *Choanocystis*
Kieler Bucht, N Kattegatt
- CHRYSMONADIDA "Incertae sedis" or doubtful**
- Aureococcus* Hargrave & Sieburth, in Sieburth & al.,1988 (TM) *A. anophagefferens* Hargrave & Sieburth, in Sieburth & al.,1988)
- Aurospphaera* Schiller,1925 (TM) *A. ovalis* Schiller,1925
- Meringosphaera* Lohmann,1902, emend. Pascher,1932
- M. mediterranea* Lohmann,1902 (TM)
?Finland, Kieler Bucht, N Kattegatt, Oslofjord, Parke & Dixon,1976 area
- M. serrata* Lohmann,1908
* Dubious species, sensu Ettl,1978
Kieler Bucht, N Kattegatt
- M. aculeata* Pascher,19
Parke & Dixon,1976 area
- M. brevispina* Pascher,19
Parke & Dixon,1976 area
- Microglena* Ehrenberg,1831 (TM) *M. monadina* Ehrenberg,1831
- M. multipunctata* Ehrenberg,1838
Kiel
- M. arenicola* Droop,19
Parke & Dixon,1976 area
- Pelagococcus* Norris, in Lewin & al.,1977 (TM) *P. subviridis* Norris, in Lewin & al.,1977
- Synrypta* Ehrenberg,1835 (TM) *S. volvox* Ehrenberg,1835
= *Synochromonas* Korchikov,19
= *Synopsis* Schiller,19
= *Volvochrysis* Schiller,19
= *Pseudosynura* Kissilev,19
- S. glomerifera* Clarke & Pennik,19
Parke & Dixon,1976 area
- S. spongiarum* Bidder,19??
@ *Granita compressa* & *Sycon ciliatum*
Parke & Dixon,1976 area
- Chrysosphaerella* Lauterborn, emend. Korschikov,1941
- C. brevispina* Korschikov,1942, emend. Harris & Bradley,1958
* Freshwater species, which does not tolerate the full salinity of the area.
Tvärminne
- * Taxa below are not mentioned in Chrétiennet-Dinet,1990
- Stichogloea* Chotat,1897
* STICHOGLOEALES Borrelly,1957, *Stichogloeaceae* Lemmermann,1899, according to Pankow,1990
- S. doederleinii* (Schmidle,1902) Wille,1911
= *Oodesmus doederleinii* Schmidle,1902
* Freshwater species, which does not tolerate the full salinity of the area.
Gulf of Finland
- Nematochrysis* Pascher,1925
* PHAEOTHAMNIALES Bourrelly,1957, *Phaeothamniaceae* (Lagerheim,1884) De Toni,1888, according to Pankow,1990
- N. hieroglyphica* Waern,1952
@ *Cladophora* & *Phragmites*
Öreggrund, Norfolk
- N. sessilis* Pascher, var. *rectensis* Carter,
- Parke & Dixon,1976 area
- Chrysomeris* Carter,1937
* PHAEOTHAMNIALES Bourrelly,1957, *Phaeosaccionaceae*, according to Parke & Green, in Parke & Dixon,1976
- C. articulata* Carter,19
Parke & Dixon,1976 area
- C. ramosa* Carter,19
SW Kattegatt, Parke & Dixon,1976 area
- Phaeosaccion* Farlow,1882
* PHAEOTHAMNIALES Bourrelly,1957, *Phaeosaccionaceae*, according to Parke & Green, in Parke & Dixon,1976
- P. collinsi* Farlow,
Parke & Dixon,1976 area
- Bicosoeca* James-Clark,1868
= *Bicoeca* von Stein,1878 = *Stylobryon* Fromentel,1874
= *Cadomona* Lackey,1939
= *Domatomona* Lackey,1939
* STRAMINOPILES Patterson,1989 (informal group name), BICOSOCALES Grasse,1926, *Bicosoecaceae* J. Clark,1868, according Värs,1992
- B. ainikkiae* Järnefelt,19??
= *B. tubiformis* Skuja,19??
* Freshwater species, which does not tolerate the full salinity of the area.
Bothnian Bay, Gulf of Finland
- B. conica* Lemmermann,1914
* Freshwater & marine
midAtlantic
- B. gracilipes* James-Clark,18?? (TM)
* Marine
Parke & Dixon,1976 area
- B. lacustris* James-Clark,1868
* Freshwater species, which does not tolerate the full salinity of the area.
Gulf of Finland, Parke & Dixon,1976 area
- B. maris* Picken,1941?
= *B. griessmannii* (Griessmann) Bourrelly,19??
S Kattegatt, Parke & Dixon,1976 area
- B. mediterranea* Pavillard,19??
Parke & Dixon,1976 area
- B. mignotii* Moestrup, Thomsen & Hibberd, in Värs,1992
* Freshwater species, which does not tolerate the full salinity of the area.
Tvärminne, Denmark, England
- B. ovata* Lemmermann,1913
* Freshwater species.
Estonia, German Baltic
- B. petiolata* (von Stein,18??) E.G: Pringsheim,19??
* Freshwater species, which does not tolerate the full salinity of the area.
Gulf of Finland
- B. planctonica* Kissilew,1931
= *B. multianulata* Skuja,1948
= *B. annulata* Stolc,1887
* Freshwater species, which does not tolerate the full salinity of the area.
Bothnian Bay, Gulf of Finland
- B. pocillum* Kent,18??
* Not mentioned by Preisig, Värs & Hällfors,1992, in Patterson & Larsen
Parke & Dixon,1976 area
- B. pontica* Valkanov,19??
* Brackish water species
?
- B. tenuis* Kent,18??
* Not mentioned by Preisig, Värs & Hällfors,1992, in Patterson & Larsen
Parke & Dixon,1976 area
- Cafeteria* Fenchel & Patterson,1988
* STRAMINOPILES Patterson,1989 (informal group name), BICOSOCALES Grasse,1926, *Bicosoecaceae* J. Clark,1868, according Värs,1992
- C. ligulifera* Larsen & Patterson,1990
?
- C. marsupialis* Larsen & Patterson,1990
?
- C. minuta* (Ruinen,19??) Larsen & Patterson,1990
?
- C. roenbergensis* Fenchel & Patterson,1988 (TM)
Tvärminne, Limfjorden, Greenland, Canaries, midAtlantic, Belize, Australia, Antarctic
- Pseudobodo* Griessmann,1913, non Hollande,1942
* STRAMINOPILES Patterson,1989 (informal group name), BICOSOCALES Grasse,1926, *Bicosoecaceae* J. Clark,1868, according Värs,1992
- P. minimus* Ruinen,19??
* Marine
?
- P. tremulans* Griessmann,1913 (TM)
* Marine
Denmark, N midAtlantic
- Hedraeophysa* Kent,1881
* CHROMULINALES Pascher,1913, *Bicosoecaceae* J. Clark,1868, according to Parke & Green, in Parke & Dixon,1976
- H. bulla* Kent,18
Parke & Dixon,1976 area
- Thallochrysis* Conrad,1914
* THALLOCHRYSIDALES Thallochrysidae, according to Parke & Green, in Parke & Dixon,1976
- T. littoralis* Anand,
Parke & Dixon,1976 area
- T. pascheri* Conrad,19
Parke & Dixon,1976 area

- Synura** Ehrenberg,1835
 * **OCHROMONADALES** Pascher,1910, Synuraceae Lemmermann,1899 according to Pankow,1990
- S. petersonii** Korschikov,1929
 * Freshwater species, which does not tolerate the full salinity of the area.
 Gulf of Finland
- S. sphagnicola** (Korschikov,1927) Korschikov,1929
 = *Skadovskielia sphagnicola* Korschikov,1927
 * Freshwater species, which does not tolerate the full salinity of the area.
 Gulf of Finland
- S. spinosa** Korschikov,1929
 * Freshwater species, which does not tolerate the full salinity of the area.
 Gulf of Finland
- S. uvella** Ehrenberg,1835, emend. Korschikov,1929
 * Freshwater species, which does not tolerate the full salinity of the area.
 Gulf of Finland, German Baltic
- HETEROMONADIDA** Leedale,1983
 = **TRIOPHYCEAE** Hibberd,1981 (p.p.)
 = **XANTHOPHYCEAE** Allorge,1930 (p.p.)
 * Doubtful order.
- Reticulosphaera** Grell,1989
 * Which suborder & family?
- R. japonensis** Grell,1990
 Japan
- R. socialis** Grell,1989 (TM)
 ?
- TRIBONEMATALES**
- Tribonemataceae**
- Tribonema** Derbès & Solier,1856
 = *Conferva affinis* G.S. West,1903
 * Freshwater species, which does not tolerate the full salinity of the area.
 * Chiefly littoral
 Gulf of Finland
- T. bombycinum** Derbès & Solier,1856
 Parke & Dixon,1976 area
- T. endozooticum** (Wille) Magne,19
 Parke & Dixon,1976 area
- T. viride** Pascher,1925
 * Freshwater species, which does not tolerate the full salinity of the area.
 * Chiefly littoral.
 Gulf of Finland
- Brachynema** Ålvik,1934
- B. bacillare** Ålvik,19
 Parke & Dixon,1976 area
- CHLORAMOEBALES** Silva,1962
 = *Chloramoebaceae* Luther,1899 = *Heterochloridae* Pascher,1926
- Chloromeson** Pascher,1930
- C. agile** Pascher,1930 (TM)
 Parke & Dixon,1976 area
- C. luteoviride** Conrad,19
 Parke & Dixon,1976 area
- C. parvum** Carter,1937
 Oslofjord, Belgium, Engl. Channel
- Heterochloris** Pascher,1914
- H. mutabilis** Pascher,1914 (TM)
 Parke & Dixon,1976 area
- Anotropis** Pascher,1939
- A. subsalina** Pascher,1939 (TM)
 Parke & Dixon,1976 area
- Ankylonoton** Pascher,1932
- A. pyreniger** Pascher,1932 (TM)
 Parke & Dixon,1976 area
- Chlorokardion** Pascher,1930 (TM *C. pleurochloron* Pascher,1930)
- C. subsalsum** Conrad,19
 Parke & Dixon,1976 area
- Phacomonas** Lohmann,1903
- P. pelagica** Lohmann,1903 (TM)
 Parke & Dixon,1976 area
- Beckingia** Ruinen,1938
- B. salina** Ruinen,1938 (TM)
 ?
- Bothrochloris** Pascher,1932
- B. longeciliata** Pascher,1932 (TM)
 ?
- RHIZOCHLORIDALES** Pascher,1925
 = *Rhizochloridaceae* Pascher,1925
- Rhizochloris** Pascher,1917
- R. mirabilis** Pascher,1917 (TM)
 Parke & Dixon,1976 area
- R. arachnoides** Carter,19
- Parke & Dixon,1976 area
- R. illoensis** Conrad,19
 Parke & Dixon,1976 area
- R. nodulariae** Bursa,1953
 @ *Nodularia spumigena*
 Gdansk
- R. solitaria** Bursa,1968
 @ *Nodularia spumigena*
 Gdansk
- Rhizolekanidae** Deflandre, in Grassé,1952
- Rhizolekane** Pascher,1932
- R. sessilis** Pascher,1932?
 ?
- TRIOPHYCEAE "incertae sedis"**
- Monodus** Chodat,1913 (TM *M. acuminata* (Germbeck) Chodat,1913)
- M. amicimei** Pascher,19
 Parke & Dixon,1976 area
- M. subsalsus** Conrad,19
 Parke & Dixon,1976 area
- Ophiocyathium** Nägeli,1??
 * Is this a Triophyceae?
- O. elongatum** W. & G.S. West,1??
 * Freshwater species, which does not tolerate the full salinity of the area.
 Gulf of Finland
- Schilleriella** Pascher,19??
 * Is this a Triophyceae?
- S. anuraea** Pascher,19??
 * Requires taxonomic reinvestigation. Systematic position uncertain.
 * May be a reduced form of *Chaetoceros subtilis*
 Gulf of Finland
- RAPHIDOMONADIDA** Heywood & Leedale,1983
 = **RAPHIDOPHYCEAE** Chadefaud, ex Silva,1980 = **RAPHIDOMONADALES** Chadefaud,1960
 = **CHLOROMONADIDA** Klebs,1892 = **CHLOROMONADOPHYCEAE**
- Merotricha** ?Mereschkowsky?,1878?
 * Marine? Which family?
- Vacuariaceae** Luther,1899 = *Chloromonadidae* Hollande, in Grassé,1952
- Olisthodiscus** N. Carter,1937
- O. luteus** N. Carter,1937 (TM)
 * Benthic, interstitial.
 Oslofjord, Belgium, Engl. Channel, E coast of USA
- Chattonella** Biecheler,1936
 = *Hemiautropicia* Hada,1974
 = *Hornellia* Subrahmanyam,1954
 = *Entostigma* Schiller,19 (p.p.)
 = *Heterostigma* Hada,19 (p.p.)
- C. subsalsa** Biecheler,1936 (TM)
 ?= *Hornellia marina* (Subrahmanyam,1954)
 Mediterranean
- Vacuolaria** Cienkowski,1870
- V. virescens** Cienkowski,1870?
 ?
- Fibrocapsa** Toriumi & Takano,1973 (TM *F. japonica* Toriumi & Takano,1973 - Japan)
- Gonyostomum** Diesing,1866 (TM *G. semen* Diesing,1866)
- G. sp. : Edler, Hälfors & Niemi,1984**
 * Freshwater species, which does not tolerate the full salinity of the area.
 Gulf of Finland
- Heterosigma** Hada,1968
- H. akashiwo** (Hada, ex Hara & Chihara,1967) Hada,1968 (TM - Japan)
 = flagellate X
 = *Olisthodiscus luteus* : Auctt., non N. Carter,1937
 * Toxic to fish. Pelagic.
 Limfjorden, Faeroes, Iceland, Scotland, Ireland, Canada, Japan
- PRYMNESIIDA**
 = **HAPTOPHYCEAE** Christensen,1962
 = **PRYMNESIOPHYCEAE** Hibberd,1976
 * Sub-classification may need revision.* All Pankow,1990 & Parke & Green, in Parke & Dixon,1976 included.
- PAVLOVINA**
 = **PAVLOVALES** Green,1976
- Pavlovaceae** Green,1976
- Pavlove** Butcher,1952
 = *Nephrochloris* Geitler & Gemisi,1925 (p.p.)
 = *Monochrysis* Skuja (p.p.)
 = *Corcontochrysis* Kalina,1970
 = *Chrysocapsa* Pascher (p.p.)
 = *Exanthemachrysis* Lepaileur (p.p.)
- P. gyrans** Butcher,1952 (TM), emend. Green & Manton,1970
 Britain
- P. lutheri** (Droop,1953) Green,1976
 = *Monochrysis lutheri* Droop,1953
 Gulf of Finland
- P. salina** (Carter,1937) Green,1976
 = *Nephrochloris salina* Carter,1937
 = *P. mesolychnon* van der Veer,1969
 Oslofjord, S, W & N Norway, Belgium, Engl. Channel, Massachusetts
- P. helicata** van der Veer,1972
 SW North Sea

- P. calceolata* van der Veer,1976
Tamar estuary
- P. virescens* Billard,1976
Parke & Dixon,1976 area
- P. ennoea* van der Veer & Leeuwis,1977
England
- Diacronema* Prauser,1958, emend. Green & Hibberd,1977
- D. vltianum* Prauser,1958, emend. Green & Hibberd,1977 (TM)
Isle of Wight
- Exanthemachrysis* Lepailleur,1970
= *Pavlova* Butcher (p.p.)
- E. gayralae* Lepailleur,1970 (TM)
Parke & Dixon,1976 area
- PRYMNESINA**
= **PRYMNESIALES** (Fritsch) Papenfuss,1955
- Those families previously placed in the order ISOCHRYSIDALES Pascher,1910 by Parke & Green (in Parke & Dixon,1976)
- Isochrysidaeae* Parke,1949
- Isochrysis* Parke,1949
- I. galbana* Parke,1949 (TM), emend. Green & Pienaar,1977
Isle of Man fish pond
- I. litoralis* Billard & Gayral,1972
Parke & Dixon,1976 area
- Chrysotila* Anand,1937, emend. Green & Parke,1975
* Benthic genus
- C. lamellosa* Anand,1937 emend. Green & Parke,1975
= *Isochrysis maritima* Billard & Gayral,1972
Parke & Dixon,1976 area
- C. stipitata* Anand,1937 emend. Green & Parke,1975
Parke & Dixon,1976 area
- I. mantonii* Reynolds,1974
- I. rotunda* Reynolds,1974 (TM), emend. Green & Pienaar,1977
S Kattegatt, Parke & Dixon,1976 area
- Dicerateria* Parke,1949
- D. gilva* Parke,1949
Oslofjord, Plymouth
- D. inornata* Parke,1949 (TM), emend. Green & Pienaar,1977
Oslofjord, S, W & N Norway, Bjørnøya, Plymouth, Wales
- Apistonema* Pascher,1925
* Benthic genus. Correct family? According to Jordan & Green,1994 does *Pleurochrysis* spp. have "Apistonema"-stages in their life cycle, so the taxonomic position of this genus is in doubt and no longer used within the classification of extant Haptophyta.
- A. pyrenigerum* Pascher,1931
@ *Phragmutes* & *Cladophora*
Øregrend, SW Kattegatt, E North Sea
- A. aestuarii* Magne,19
Parke & Dixon,1976 area
- A. carterae* Anand,1937
Parke & Dixon,1976 area
- A. submarinum* Dang.,19
Parke & Dixon,1976 area
- Chrysonema* Anand,1937
* Not included in Chriennet-Dinet,1990. Correct family? Genus of uncertain affinity according to Jordan & Green,1994
- C. littorale* Anand,1937
Parke & Dixon,1976 area
- Noelaenhabdaceae* Jerkovic,1970
- Emiliania* Hay & Mohler, in Hay, Mohler, Roth, Schmidt & Boudreaux.,1967
- E. huxleyi* (Lohmann,1902) Hay & Mohler, in Hay, Mohler, Roth, Schmidt & Boudreaux.,1967 (TM)
= *Pontosphaera huxleyi* Lohmann,1902
= *Coccolithus huxleyi* (Lohmann,1902) Kampfner,1943
= *Gephyrocapsa huxleyi* (Lohmann,1902) Reinhartd,1972
Tvråminne, Kieler Bucht, Øresund, Kattegatt, Skagerrak, all Norway, Plymouth, all oceans except Polar & Antarctic seas
- Gephyrocapsa* Kampfner,1943
- G. oceanica* Kampfner,1943 (TM)
Parke & Dixon,1976 area
- G. kampfneri* Deflandre & Fert,1954
* Not mentioned in Jordan & Green,1994
Faeroe-Shetland area, Parke & Dixon,1976 area
- Reticulofenestra* Hay, Mohler, Roth, Schmidt & Boudreaux,1967, emend. Gallagher,1989
= *Crenalithus* Roth,1973
- Those families previously placed in the order PRYMNESIALES by Parke & Greene (in Parke & Dixon,1976)
- Phaeocystaceae* Lagerheim,1896
- Phaeocystis* Lagerheim,1893
- P. pouchetii* (Hariot, in Pouchet,1892) Lagerheim,1893 (TM)
= *Tetraspora pouchetii* Hariot, in Pouchet,1892
?= *P. amoeboides* Büttner,1910 ?= *P. sphaeroides* Büttner,1910
* The two species described by Büttner from Kieler Bucht have not been recorded since their description. Central & W Baltic, Belt Sea, Øresund, Kattegatt, Limfjorden, Skagerrak, Oslofjord, Bergen, North Sea
- P. globosa* Scherffel,1899
?
- Prymnesiaceae* Conrad,1926 = *Prymnesiidae* Conrad,1926
- * Zooxanthellae of three acantharians, *Amphilochete elongata*, *Lithopelta muelleri* & *Acanthometra pellucida* have been identified as belonging to this family (Febvre & Febvre-Chevalier,1979).
- Prymnium* Massart,1920
- P. annuliferum* Billard,1983
?
- P. gladiociliatum* (Büttner,1910) Jordan & Green,1994
= *Wysotskia gladiociliata* Büttner,1910
* In need of detailed re-examination according to Jordan & Green,1994
Kieler Bucht
- P. parvum* Carter,1937
* Toxic to fish.
S Baltic, Belt Sea, Limfjorden, S & central Kattegatt, Oslofjord, S & N Norway, E North Sea, Netherlands, Engl. Channel, Massachusetts
- P. patelliferum* Green, Hibberd & Pienaar,1982
* Probably toxic to fish. Not distinguishable from *P. parvum* in light microscope (differs by scale morphology). Baltic, cosmopolitan
- P. saltans* Massart, ex Conrad,1926 (TM)
Parke & Dixon,1976 area
- P. minutum* N. Carter,1937
Parke & Dixon,1976 area
- P. zebrinum* Billard,1983
?
- Chrysocromulina* Lackey,1939 (TM) *C parva* Lackey, emend. Parke & al.,1972 - fresh water
= *Chrysocampanda* Fournier,1971
- C. acantha* Leadbeater & Manton,1971
Kattegatt, Bergen, E North Sea, Plymouth
- C. alifera* Parke & Manton, in Parke, Manton & Clarke,1956
N Kattegatt, Bergen, Plymouth
- C. aphelis* Moestrup & Tangen,1986
S Kattegatt
- C. bergenensis* Leadbeater,1972
N Kattegatt, Bergen, E North Sea
- C. birgeri* Hallfors & Niemi,1974
Gulf of Finland
- C. brachycylindra* Hallfors & Thomsen,1986
Finland, S Kattegatt, S Norway
- C. brevifilum* Parke & Manton, in Parke, Manton & Clarke,1956
N Kattegatt, Bergen, off SW Britain
- C. camella* Leadbeater & Manton,1969
Parke & Dixon,1976 area
- C. campanulifera* Manton & Leadbeater,1974
N Kattegatt, E North Sea
- C. chiton* Parke & Manton, in Parke, Manton & Clarke,1958
N Kattegatt, Bergen, Plymouth
- C. cyathophora* Thomsen,1979
* Warm water species.
W Baltic, Belt Sea, SW Kattegatt
- C. cymbium* Leadbeater & Manton,1969
N Kattegatt, Bergen, Plymouth
- C. elegans* Estep, Davis, Hargraves & Sieburth,1984
N Atlantic
- C. ephippium* Parke & Manton, in Parke, Manton & Clarke,1956
N Kattegatt, Oslofjord, Bergen, E North Sea, Plymouth, New Zealand
- C. ericina* Parke & Manton, in Parke, Manton & Clarke,1956
Finland, W Baltic, Øresund, Belt Sea, Kattegatt, Limfjorden, Oslofjord, S & W Norway, Bjørnøya, Plymouth, Mediterranean
- C. fragilis* Leadbeater,1972
?N Kattegatt, Bergen, Parke & Dixon,1976 area
- C. herdmani* Leadbeater,1972
N Kattegatt, E North Sea, Bergen
- C. hirta* Manton,1978
Finland, W Baltic, Øresund, Belt Sea, Kattegatt, S Norway, Greenland
- C. kappa* Parke & Manton, in Parke, Manton & Clarke,1955
Oslofjord, Bergen, Bjørnøya, Isle of Man, Irish Sea, off S England, New Zealand
- C. leadbeateri* Estep, Davis, Hargraves & Sieburth,1984
Norwegian W coast
- C. macrura* Manton,1972
= *C. IIIb* : Thronsdene,1969
Denmark, S, W & N Norway, Britain, Mediterranean
- C. mantoniæ* Leadbeater,1972
Gulf of Finland, W Baltic, Øresund, Belt Sea, Kattegatt, Bergen, Greenland, Mediterranean
- C. megacylindra* Leadbeater,1972
N Kattegatt, Bergen
- C. microcylinatra* Leadbeater,1972
N Kattegatt, Bergen
- C. minor* Parke & Manton, in Parke, Manton & Clarke,1955
Baltic (>4 ppt S), SW Kattegatt, Oslofjord, Bergen, off SW England
- C. pachycylindra* Manton, Oates & Course,1981
Engl. Channel
- C. parkeae* Green & Leadbeater,1972
W Baltic, Belt Sea, Kattegatt, Limfjorden, Bergen, Plymouth
- C. pelagica* Estep, Davis, Hargraves & Sieburth,1984
N Atlantic
- C. polyplepis* Manton & Parke,1962
* Toxic to fish & invertebrates.

- Baltic (>4 ppt S), Kattegatt, Bohusl., S & W Norway, Plymouth
- C. pringsheimii* Parke & Manton, 1962
W Baltic, Belt Sea, Øresund, Limfjorden, Kattegatt, Oslofjord, S Norway, Bergen, Plymouth, Biscay, Mediterranean
- C. pyramidosa* Thomsen, 1977
Belt Sea, S Kattegatt
- C. simplex* Estep, Davis, Hargraves & Sieburth, 1984
N Atlantic
- C. spinifera* (Fournier, 1971) Pienaar & Norris, 1979
= *Chrysocampanula spinifera* Fournier, 1971
Finland, E Kattegatt, Limfjorden, S Norway, Britain, Greenland
- C. strobilus* Parke & Manton, in Parke, Manton & Clarke, 1959
N Kattegatt, Bergen, Bjørnøya, Plymouth
- C. tenuisquama* Estep, Davis, Hargraves & Sieburth, 1984
N Atlantic
- Corymbellus* Green, 1976
- C. aureus* Green, 1976 (TM)
Engl. Channel
- Platychrysis* Geitler, 1930, emend. Gayral & Fresnel, 1983
- P. nigra* Geitler, 1930 (TM)
Parke & Dixon, 1976 area
- P. pienaarii* Gayral & Fresnel, 1983
?
- P. simplex* Gayral & Fresnel, 1983
?
- Those families previously placed in the order COCCOSPHAERALES (= COCCOLITHOPHORALES Schiller, 1926) by Parke & Green (Parke & Dixon, 1976)
- Braarudosphaeraceae** Deflandre, 1947
- Braarudosphaera** Deflandre, 1947
- B. bigelovii* (Gran & Braarud) Deflandre, 1947 (TM)
S Kattegatt, Engl. Channel
- B. magnei* Lefort, 1972
Parke & Dixon, 1976 area
- Calciosoleniaceae** Kamptner, 1937
- Calciosolenia** Gran, in Murray & Hjort, 1912
= *Acanthosolenia* Bernard, 1939 (TM) *A. mediterranea* Bernard, 1939 - Mediterranean
- C. murrayi* Gran, in Murray & Hjort, 1912 (TM)
Parke & Dixon, 1976 area
- Anoplosolenia* Deflandre, 1952
- A. brasiliensis* (Lohmann) Deflandre, 1952 (TM)
= *Cylindrotheca brasiliensis* Lohmann
Faeroe-Shetland area, Parke & Dixon, 1976 area
- Navisolenia* Lecal, 1965, emend. Leadbeater & Morton, 1973 (TM) *N. aprilei* Lecal, 1965 - E Mediterranean
- Ceratolithaceae** Norris, 1965
- Ceratolithus** Kamptner, 1950 (TM) *C. cristatus* Kamptner, 1950
- Coccophoraceae** Kamptner, 1928
- Coccophora** Lohmann, 1902
- Coccolithus** Schwartz, 1894
= *Crystallolithus* Garder & Markali, 1956 (p.p.)
= *Coccophora* Lohmann, 1902
= *Coccospheara* Walllich, 1877
- C. pelagicus* (Walllich, 1877) Schiller, 1930
= *Crystallolithus hyalinus* Gaarder & Markali, 1956 (its holococcolith-bearing mobile phase)
= *Crystallolithus braarudii* Garder, 1962 (its holococcolith-bearing mobile phase)
Parke & Dixon, 1976 area
- Calcidiscus* Kamptner, 1950
- C. leptotorus* (Murray & Blackman, 1898) Loeblich Jr. & Tappan, 1978 (TM)
= *Cyclococcolithus leptotorus* (Murray & Blackman, 1898) Kamptner, 19
= *Crystallolithus rigidus* Gaarder, in Heimdal & Gaarder, 1980 (its holococcolith-bearing mobile phase)
Parke & Dixon, 1976 area
- Crenalithus* Roth, 1973 (TM) *C. doronicoidea* Roth, 1973 - fossil
- Cruciplacolithus** Hay & Mohler, in Hay, Mohler, Roth, Schmidt & Boudreux, 1967 (TM) *C. tenuis* (Stradner) Hay & Mohler, in Hay, Mohler, Roth, Schmidt & Boudreux, 1967 - fossil
- Cyclolithus** Kamptner ex Deflandre, 1962
= *Cyclolithella* Loeblich Jr. & Tappan, 1963
- Hayaster* Bukry, 1973 (TM) *H. perplexus* (Bramlette & Riedel) Bukry, 1973 - W Pacific
- Neosphaera** Lecal-Schlauder, 1950 (TM) *N. coccilithomorpha* Lecal-Schlauder, 1950 - N Africa
- Oolithotus** Reinhardt, in Cohen & Reinhardt, 1968
= *Cyclococcolithina* Wilcoxon, 1970
- O. fragilis* (Lohmann) Reinhardt, 1972 (TM)
= *Cyclococcolithus fragilis* (Lohmann) Deflandre, in Deflandre & Fert, 1954
Parke & Dixon, 1976 area
- Umbilicosphaera* Lohmann, 1902 (TM) *U. sibogae* (Weber Van Bosse) Gaarder, 1970
= *Cycloplacolithus* Kamptner, 1963
- Helicosphaeraceae** Black, 1971, emend. Jafar & Martini, 1975
- Helicosphaera** Kamptner, 1954
= *Coccophora* Lohmann, 1902 (p.p.)
= *Coccospheara* Walllich, 1877 (p.p.)
= *Helicopontosphaera* Hay & Mohler, in Hay, Mohler, Roth, Schmidt & Boudreux, 1967
- H. carteri* (Walllich, 1877) Kamptner, 1954 (TM)
Parke & Dixon, 1976 area
- Hymenomonadaceae** Senn, in Engler & Prantl, 1900, emend. Gayral, 1989
- Hymenomonas** von Stein, 1878
- H. roseola* von Stein, 1878 (TM)
= *H. coccolithophora* Massart & Conrad, in Conrad, 1926
= *H. danubiensis* Kamptner, 1928
= *H. flava* Stokes, 1888
= *H. scherffelii* Conrad, 1928
* Fresh water species.
Finland (brackish water), Parke & Dixon, 1976 area
- H. globosa* (Magne, 1954) Gayral & Fresnel, 1976
= *Pontosphaera globosa* Magne, 1954
Engl. Channel
- H. lacuna* Pienaar, 1976
Normandie
- H. coronata* Mills, 1975
Texas coast
- Ochrosphaera** Schussnig, 1930
- O. neapolitana* Schussnig, 1930 (TM)
= *O. verrucosa* Schussnig, 1930
Engl. Channel, W France, Mediterranean, Parke & Dixon, 1976 area
- Papposphaeraceae** Jordan & Young, 1990
= *Deflandriaceae* Black, emend. Norris, 1983
- Papposphaera* Tangen, 1972
= *Turrisphaera* Manton, Sutherland & Oates, 1976 (p.p.) (holococcolith-bearing phase)
- P. borealis* Manton, Sutherland & Oates, 1976
= *Turrisphaera borealis* Manton, Sutherland & Oates, 1976 (TM) of *Turrisphaera* Manton, Sutherland & Oates, 1976
S Norway, W Greenland
- P. arctica* Manton, Sutherland & Oates, 1976
= *Turrisphaera arctica* Manton, Sutherland & Oates, 1976
W Baltic, Belt Sea, SW & central Kattegatt, W Greenland
- P. lepida* Tangen, 1972 (TM)
Øresund, SW Kattegatt, Parke & Dixon, 1976 area
- P. polybotrys* (Thomsen, 1980) Thomsen, Østergaard & Hansen, 1991
= *Turrisphaera polybotrys* Thomsen, 1980
W Greenland
- P. pomponas* Manton & Oates, 1975
= *Turrisphaera* Manton, Sutherland & Oates, 1976 (p.p.) (holococcolith-bearing phase)
- P. virgulosa* Manton & Sutherland, 1975
Gulf of Finland, W Baltic, Belt Sea, Kattegatt, S Norway, E North Sea, W Greenland
- P. flabellifera* Manton & Oates, 1975 (TM)
W Baltic, Belt Sea, Kattegatt, S Norway, E North Sea, Greenland
- Trigonaspis* Thomsen, 1980
- T. minutissima* Thomsen, 1980
Belt Sea, SW, N & E Kattegatt, S Norway, W Greenland
- T. discoensis* Thomsen, 1980
SW & E Kattegatt, W Greenland
- Pleurochrysidae** Gayral, 1989
- Pleurochrysis** Pringsheim, 1955, emend. Gayral & Fresnel, 1983
= *Cricospheara* Braarud, 1960 (p.p.)
- P. scherffelii* Pringsheim, 1955 (TM)
= *Hymenomonas pringsheimii* Parke & Green, in Parke & Dixon, 1976
non = *Hymenomonas scherffelii* Conrad, 1928
Parke & Dixon, 1976 area
- P. carterae* (Braarud & Fagerland, 1946) Christensen, 1978
= *Syracospheara carterae* Braarud & Fagerland, 1946
= *Cricosphaera carterae* (Braarud & Fagerland, 1946) Braarud, 1960
= *Hymenomonas carterae* (Braarud & Fagerland, 1946) Braarud, 1954
= *Syracospheara brandtii* Carter, 1937
Gulf of Finland, S & W Baltic, Øresund, Belt Sea, Kattegatt, Oslofjord, S, W & N Norway, Engl. Channel
- P. gayraliae* (Beaufort, 1978) Jordan, Kleijne & Heimdal, 1993
= *Cricospheara gayraliae* Beaufort, 1978
Isle of Man
- P. pseudoroscoffensis* Gayral & Fresnel, 1983
Calvados
- P. placolithoides* Gayral, 1989
Belle-Ile-en-Mer (Atlantic France)
- P. roscoffensis* (Dangeard, 1934) Gayral, 1989
= *Syracospheara roscoffensis* Dangeard, 1934
= *Pontosphaera roscoffensis* Chadeauf & Feldmann, 1949
= *Syracolithus roscoffensis* (Dangeard, 1934) Magne, 1954
= *Cricosphaera roscoffensis* (Dangeard, 1934) Gayral & Fresnel, 1976
= *Cricosphaera roscoffensis* var. *haptonemofera* Inouye & Chihara, 1979
= *Cricosphaera haptonemofera* (Inouye & Chihara, 1979) Gayral & Fresnel, 1983
= *Hymenomonas roscoffensis* (Chadeauf & Feldmann, 1949) Lecal, in Parke & Dixon, 1976
* occasionally forming brown coatings over fine sand. Cells may attach to *Cladophora* Roscoff
- P. elongata* (Droop, 1955) Jordan, Kleijne & Heimdal, 1993
= *Cricospheara elongata* (Droop) Braarud, 1960 (TM)
= *Syracospheara elongata* Droop, 1955
= *Hymenomonas elongata* (Droop) Parke & Green, in Parke & Dixon, 1976
Parke & Dixon, 1976 area
- P. sp.*
= *Cricospheara* sp.: Gayral, 1989
= *Coccolithus pelagicus*: Auctt., non (Wallich)
?
- Pontosphaeraceae** Lemmermann, in Brandt & Apstein, 1909
- Pontosphaera** Lohmann, emend. Burns, 1973
= *Discolithina* Loeblich Jr. & Tappan, 1963 (TM) *D. vigintifera* (Kamptner, ex Deflandre) Loeblich Jr. & Tappan, 1963
- * Only 3 species recognized by Chrétiennet-Dinet, 1990. Which?

- P. syracusana* Lohmann,1902 (TM)
Parke & Dixon,1976 area
- P. borealis* Ostenfeld
* Not mentioned in Jordan & Green,1994
Parke & Dixon,1976 area
- P. haekelii* Lohmann,19
* Not mentioned in Jordan & Green,1994
Parke & Dixon,1976 area
- P. nana* Kamptner,19
* Not mentioned in Jordan & Green,1994
Parke & Dixon,1976 area
- P. pellucida* Lohmann,19
* Not mentioned in Jordan & Green,1994
Parke & Dixon,1976 area
- P. variabilis* Halldal & Markali,1955
* Not mentioned in Jordan & Green,1994
Northermost part of Parke & Dixon,1976 area
- P. discopora* Schiller,1925
Adriatic
- P. turgida* Müller, in Müller, Blanc-Vernet, Chamley & Froget,1974
Mediterranean
- Lohmannsphaera* Schiller,1913 (TM *L. adriatica* Schiller,1913)
Genus of uncertain affinity, possibly =*Scyphosphaera* Lohmann,1902, although one species was transferred tentatively to the prasinophyte genus *Pterosperma*.
- Scyphosphaera* Lohmann,1902
- S. apsteinii* Lohmann,1902 (TM)
off Bergen, Parke & Dixon,1976 area
- Rhabdosphaeraceae Haeckel,1894
- Rhabdosphaera* Haeckel,1894
= *Aspidorhabdus* Hay & Towe,1962
- R. clavigera* Murray & Blackman,1898 (TM)
Parke & Dixon,1976 area
- R. clavigera* var. *stylifera* (Lohmann,19??) Kleijne & Jordan,1990
Parke & Dixon,1976 area
- R. hispida* Lohmann,19
* Not mentioned in Jordan & Green,1994
Parke & Dixon,1976 area
- R. longistylis* Schiller,19
* Not mentioned in Jordan & Green,1994
Parke & Dixon,1976 area
- Palusphaera* Lecal,1965, emend. Norris,1984 (TM *P. vandellii* Lecal,1965, emend. Norris,1984 - Indian Oc.)
- Discosphaera* Haeckel,1894 (TM *D. thomsoni* Ostenfeld,1900)
- D. tubifera* (Murray & Blackman,1898) Ostenfeld,1900
= *Rhabdosphaera tubifera* Murray & Blackman,1898
off Poland, Parke & Dixon,1976 area
- Acanthoica* Lohmann,1903, emend. Schiller,1913, emend. Kleijne,1992 (TM *A. coronata* Lohmann,1902)
- A. acanthifera* Lohmann ex Lohmann,1913
Bergen, Parke & Dixon,1976 area, Gulf Stream, Sargasso Sea, Bermuda, Adriatic
- A. biscayensis* Kleijne,1992
Biscay
- A. janchenii* Schiller,1925
Adriatic
- A. maxima* Heimdal, in Heimdal & Gaarder,1981
N Atlantic
- A. quattrospina* Lohmann,1903
Belt Sea, Kattegatt, Limfjorden, Oslofjord
- Algiosphaera* Schlauder,1945 emend. Norris,1984
- A. oryzia* Schlauder,1945 (TM)
S Mediterranean
- Anacanthoica* Deflandre,1952 (TM *A. acanthos* (Schiller) Deflandre,1952)
- A. cidaris* (Schlauder) Kleijne,1992
Mediterranean
- Cyrtosphaera* Kleijne,1992
- C. aculeata* (Kamptner) Kleijne,1992
= *Acanthoica aculeata* Kamptner,19??
Parke & Dixon,1976 area
- C. cucullata* (Lecal-Schlauder) Kleijne,1992
Mediterranean
- Anthosphaera* Kamptner,1936
* The species in this genus should be transferred to either *Calyptrolithina* Heimdal,1982 or *Algiosphaera* Schlauder, according to Chrétiennot-Dinet,1990
- A. quadricornu* (Schiller,1925) Halldal & Markali,1955
= *Syracosphaera quadricornu* Schiller,1925
Bergen, Parke & Dixon,1976 area, Adriatic, Gulf Stream
- A. robusta* (Lohmann) Kamptner,19
Parke & Dixon,1976 area
- Syracosphaeraceae (Lohmann) Lemmermann,1903
= *Deutschlandiaceae* Kamptner,1928
= *Halopappaceae* Kamptner,1928
- Alisphaera* Heimdal,1973 emend. Jordan & Chamberlain,1993 (TM *A. ordinata* (Kamptner) Heimdal,1973)
- A. capulata* Heimdal, in Heimdal & Gaarder,1981
NE Atlantic
- Alveosphaera* Jordan & Young,1990 (TM *A. bimurata* (Okada & McIntyre) Jordan & Young,1990)
- Calciopappus* Gaarder & Ramsfjell,1954, emend. Manton & Oates,1985
- C. caudatus* Gaarder & Ramsfjell,1954 (TM)
W Baltic, Belt Sea, SW Kattegatt
- C. rigidus* Heimdal, in Heimdal & Gaarder,1981
NE Atlantic
- Canistrolithus* Jordan & Chamberlain,1993 (TM *C. valliformis* Jordan & Chamberlain,1993)
- Coronosphaera* Gaarder, in Gaarder & Heimdal,1977
- C. mediterranea* (Lohmann) Gaarder, in Gaarder & Heimdal,1977 (TM)
= *Syracosphaera mediterranea* Lohmann,T9
= *Syracosphaera pulchroides*
= *Syracosphaera tuberculata*
N Kattegatt, Oslofjord, Parke & Dixon,1976 area
- Gaarderia* Kleijne,1993 (TM *G. corolla* (Lecal) Kleijne,1993)
- Halopappus* Lohmann,1912 (TM *H. valselii* Lohmann,1912)
* Not refound since description and possibly a synonym of *Michaelsarsia* Gran according to Jordan & Green,1994
- Michaelsarsia* Gran, in Murray & Hjort,1912, emend. Manton, Bremer & Oates,1984
- M. adriaticus* (Schiller) Manton, Bremer & Oates,1984
= *Halopappus adriaticus* Schiller,19??
Adriatic
- M. elegans* Gran, in Murray & Hjort,1912 (TM) emend. Manton, Bremer & Oates,1984
Parke & Dixon,1976 area
- M. falklandica* Lohmann
* Not mentioned in Jordan & Green,1994
Northermost part of Parke & Dixon,1976 area
- Ophiaster* Gran, emend. Manton & Oates,1983
= *Bernardosphaera* Lecal-Schlauder,1951
= *Lacrymasphaera* Lecal,1966
- O. hydroides* (Lohmann,1903) Lohmann,1913 (TM), emend. Manton & Oates,1983
= *Meringosphaera hydroides* Lohmann,1903
Oslofjord, S Norway, N Atlantic, Mediterranean
- O. formosus* Gran, in Murray & Hjorth,1912 sensu Gaarder, emend. Manton & Oates,1983
?
- Najadea* Schiller,1913 (*N. gloriosa* Schiller,1913 - Adriatic)
* Genus of uncertain affinity according to Jordan & Green,1994
- Syracosphaera* Lohmann, emend. Gaarder, in Gaarder & Heimdal,1977
= *Deutschlandia* Lohmann,1912, emend. Gaarder, in Heimdal & Gaarder,1981
= *Syracosphaera* Lecal & Bernheim,1960
= *Canosphaera* Gaarder, in Gaarder & Heimdal,1977
- S. anthos* (Lohmann,1912) Janin,1987
= *Deutschlandia anthos* Lohmann, emend. Gaarder, in Heimdal & Gaarder,1981 (TM)
Faeroe-Shetland area, Parke & Dixon,1976 area
- S. halldalii* Gaarder ex Jordan & Green,1994
Norw. Sea
- S. histrica* Kamptner,1941
NE Mediterranean
- S. molischii* Schiller,1925
Parke & Dixon,1976 area
- S. nodosa* Kamptner,1941
Parke & Dixon,1976 area, NE Mediterranean
- S. pulchra* Lohmann,1902 (TM)
Parke & Dixon,1976 area
- S. binodata* (Kamptner) Kamptner,19
* Not mentioned in Jordan & Green,1994
Oslofjord, Faeroe-Shetland area, Parke & Dixon,1976 area
- S. dentata* Lohmann,19
* Not mentioned in Jordan & Green,1994
Parke & Dixon,1976 area
- S. spinosa* Lohmann,19
* Not mentioned in Jordan & Green,1994
Parke & Dixon,1976 area
- S. subsalsa* (Conrad) Kamptner,19
* Not mentioned in Jordan & Green,1994
Parke & Dixon,1976 area
- S. tenuis* Lohmann,19
* Not mentioned in Jordan & Green,1994
Parke & Dixon,1976 area
- Tetralithoides* Theodoridis,1984, emend. Jordan, Kleijne & Heimdal,1993
- Umbellophaera* Paasche, in Markali & Paasche,1955, emend. Gaarder, in Heimdal & Gaarder,1981 (TMU).
tenuis (Kamptner) Paasche, in Markali & Paasche,1955
- Calyptrösphaeraceae Boudreux & Hay,1969
- Anthosphaera* Kamptner, emend Kleijne,1991 (TM *A. fragaria* Kamptner, emend. Kleijne,1991)
- Calicospaera* Kleijne,1991
- Calyptrolithina* Heimdal,1982
= *Cavosphaera* Lecal & Bernheim,1960
= *Corisphaera* Kamptner,1936 (p.p.)
- C. divergens* (Halldal & Markali,1955) Heimdal,1982 (TM)
= *Zygospaera divergens* Halldal & Markali,1955
Parke & Dixon,1976 area
- C. wettsteinii* (Kamptner) Kleijne,1991
= *Homozygospaera wettsteinii* (Kamptner) Halldal & Markali,1955
Faeroe-Shetland area, Parke & Dixon,1976 area
- Calyptrolithophora* Heimdal, in Heimdal & Gaarder,1980
= *Corisphaera* Kamptner,1936 (p.p.)

- C. gracillima* (Kamptner) Heimdal, in Heimdal & Gaarder, 1980
NE Atlantic
- C. hastiana* (Gaarder) Heimdal, in Heimdal & Gaarder, 1980
NE Atlantic
- C. papillifera* (Halldal) Heimdal, in Heimdal & Gaarder, 1980 (TM)
= *Sphaerocalyptra papillifera* (Halldal) Halldal
Parke & Dixon, 1976 area
- Calyptrisphaera* Lohmann, 1902 (TM *C. globosa* Lohmann, 1902)
- C. cialdii* Borsetti & Cati, 1976
Mediterranean
- C. oblonga* Lohmann, 1902
Parke & Dixon, 1976 area
- C. sphaeroidea* Schiller, 1913
Parke & Dixon, 1976 area
- C. sp.* : Edler, Hälfors & Niemi, 1984
Kattegatt - Belt Sea area
- Corisphaera* Kamptner, 1937
- C. gracilis* Kamptner, 1937 (TM)
Mediterranean
- C. strigalis* Gaarder, 1962
South part of Parke & Dixon, 1976 area
- C. tyrrheniensis* Kleijne, 1991
Mediterranean
- Daktylethora* Gartner, in Gartner & Bükry, 1969 (TM *D. punctulata* Gartner, in Gartner & Bükry, 1969 - fossil)
- Flosculosphaera* Jordan & Kleijne, in Kleijne, Jordan & Chamberlain, 1991
- F. calceolariopsis* Jordan & Kleijne, in Kleijne, Jordan & Chamberlain, 1991 (TM)
NE Atlantic
- F. sacculus* Kleijne & Jordan, in Kleijne, Jordan & Chamberlain, 1991
NE Atlantic
- Gliscolithus* Norris, 1985 (TM *G. amitakareniae* Norris, 1985 - Indian Ocean)
- Helladosphaera* Kamptner, 1937
= *Corisphaera* Kamptner, 1936 (p.p.)
- H. cornifera* (Schiller) Kamptner, 1937 (TM)
Parke & Dixon, 1976 area
- Homozygosphaera* Deflandre, 1952 (TM *H. spinosa* (Kamptner) Deflandre, 1952)
= *Corisphaera* Kamptner, 1936 (p.p.)
- H. triarcha* Halldal & Markali, 1955
?N Atlantic
- H. vavilovii* Borsetti & Cati, 1976
Mediterranean
- H. versellii* Borsetti & Cati, 1979
Mediterranean
- Periphyllophora* Kamptner, 1937 (TM *P. mirabilis* (Schiller) Kamptner, 1937)
- Poricalyptra* Kleijne, 1991
- P. gaarderae* (Borsetti & Cati) Kleijne, 1991
Mediterranean
- P. isselii* (Borsetti & Cati) Kleijne, 1991
Mediterranean
- P. magnaghii* (Borsetti & Cati) Kleijne, 1991
Mediterranean
- Poritectolithus* Kleijne, 1991
- Sphaerocalyptra* Deflandre, in Grasse, 1952 (TM *S. quadridentata* (Schiller) Deflandre, 1952)
= *Corisphaera* Kamptner, 1936 (p.p.)
- Syracolithus* (Kamptner) Deflandre, 1952 (TM *S. dalmaticus* (Kamptner) Loeblich Jr. & Tappan, 1963)
- S. corolla* Lecal, 19
- * Not mentioned in Jordan & Green, 1994
Faeroe-Shetland area, Parke & Dixon, 1976 area
- S. quadriperforata* (Kamptner) Gaarder, in Heimdal & Gaarder, 1980
= *Homozygosphaera quadriperforata* (Kamptner) Gaarder
Parke & Dixon, 1976 area
- Zygospaera* Kamptner, 1937, emend. Heimdal, 1982
- Z. amoena* Kamptner, 1937
Mediterranean
- Z. hancockii* (Borsetti & Cati) Heimdal, 1982
Mediterranean
- Z. hellenica* Kamptner, 1937 (TM)
Mediterranean
- Z. marsili* (Borsetti & Cati) Heimdal, 1982
Mediterranean
- Quaternariella* Thomsen, 1980
* Genus of unknown taxonomic affiliation according to Jordan & Green, 1994
- Q. obscura* Thomsen, 1980 (TM)
W Greenland
- Balaniger* Thomsen & Oates, 1978
* Genus of unknown taxonomic affiliation according to Jordan & Green, 1994
- B. balticus* Thomsen & Oates, 1978 (TM)
Gulf of Finland, Tvärminne, W Baltic, Belt Sea, Öresund, S & E Kattegatt, Greenland
- COCCOLITHOPHORALES "incertae sedis"**
- Calciarcus* Manton, Sutherland & Oates, 1977 (TM *C. alaskensis* Manton, Sutherland & Oates, 1977 - S Alaska)
- Florisphaera* Okada & Honjo, 1973 (TM *F. profunda* Okada & Honjo, 1973)
- Gladiolithus* Jordan & Chamberlain, 1993 (TM *G. flabellatus* (Halldal & Markali, 1955) Jordan & Chamberlain, 1993)
- Jomonolithus* Inouye & Chihara, 1983 (TM *J. littoralis* Inouye & Chihara, 1983 - NW Pacific)
- Polycretater* Manton & Oates, 1980 (TM *P. galapagensis* Manton & Oates, 1980 - Galapagos)
- Thorosphaera* Ostendorf, 1910 (TM *T. elegans* Ostendorf, 1910)
* According to Ostendorf & Green, 1994 is this a genus of uncertain affinity, possibly *Scyphosphaera*?
Lohmann, 1902.
- Turrilithus* Jordan, Knapperbusch, Simpson & Chamberlain, 1991 (TM *T. latericioides* Jordan, Knapperbusch, Simpson & Chamberlain, 1991)
- Vexillarius* Jordan & Chamberlain, 1993 (TM *V. cancellifer* Jordan & Chamberlain, 1993)
- Wigwamma* Manton, Sutherland & Oates, 1977
- W. arctica* Manton, Sutherland & Oates, 1977 (TM)
E & N Kattegatt, W Greenland
- W. scenozonion* Thomsen, 1980
N Kattegatt
- R. regalis** (Gaarder) Gaarder, 19
Parke & Dixon, 1976 area
- VOLVOCIDA** Francé, 1894
= **VOLVOCALES** Oltmans, 1904
* All Pankow, 1990 included.
- ?SUBORDER?**
- Planktococcus* Korschikov, 1???
- P. sphaerocystiformis* Korschikov, 1???
- * Freshwater species, which does not tolerate the full salinity of the area.
Gulf of Finland
- Planktosphaeria* G.M. Smith, 1???
- P. gelatinosa* G.M. Smith, 1???
- * Freshwater species, which does not tolerate the full salinity of the area.
Gulf of Finland
- CHLOROCOCCIDINA**
- Coccomyxaceae**
- Gloeocystis* Nägeli, 1???
- G. plantonica* (W. & G.S. West, 18??) Lemmermann, 19??
* Freshwater species, which does not tolerate the full salinity of the area.
Gulf of Finland
- Sphaerocystis* Chodat, 1???, emend. Korschikov, 1???
- non = *Sphaerocystis* Léger, 1892 (Apicomplexa)
- S. schroeteri* Chodat, 1???, emend. Korschikov, 1???
- * Freshwater species, which does not tolerate the full salinity of the area.
Gulf of Finland, Bothnian Bay, Bothnian Sea
- POLYBLEPHARIDINA**
- Polyblepharidae** G.S. West, 1916
- Dunaliella* Teodoresco, 1905
- D. salina* (Dunal, 1838)
= *Haematococcus salinus* Dunal, 1838 (TM)
S France saline water
- D. marina* ??, 1???
- p. resund
- Ulochloris* Pascher, 1916
- U. oscillans* Pascher, 1916 (TM)
?
- CHLAMYDOMONADINA**
- = **CHLAMYDOMONADACEAE** G.M. Smith, 1920
- Haematococcaceae**
- Haematococcus* Agardh, 18??
* 5 sp., which may occur in fresh water or in small saline rockpools
- H. droebakiensis* ??, 1???
- p. oslofjord
- Chlamydomonadaceae**
- Chlamydomonas* Ehrenberg, 1837
- C. reginae* Ettl & Green, 1973
* sand-living.
Bretagne
- C. marina* Cohn, 1???
- p. cold water species. Requires taxonomic reinvestigation.
-
- Riga Bay
- C. sp.* : Edler, Hälfors & Niemi, 1984
Gulf of Finland, Archipelago Sea, ?Kattegatt
- Brachiomonas* Bohlin, 1897
- B. submarina* Bohlin, 1897
Gulf of Finland, Kattegatt, Oslofjord, S Norway, Bergen, Ålesund, Trondheim, Tromsö, Briain & Europe
brackish water
- B. simplex* Hazen, 1922
Oslofjord (rock pool), Ålesund, Britain brackish water
- Scherffelia* Pascher

*Carteriaceae**Carteria* Diesing,1866, emend. Francé,1896*C. marina* Wulff,1916
Baltic*Phacotaceae**Pteromonas* Seligo,
SPONDYLOMORINA
* No marine species?VOLVOCINA
= VOLVOCACEAE (Ehrenberg,1838) Kirchner,1878Volvocidae Pavillard, in Grassé,1952
= Volvocaceae*Volvox* Linnaeus,1758, emend. Ehrenberg,1830*V. aureus* Ehrenberg,1830
Baltic*V. tertius* Meyer,1896
German Baltic*Gonium* O.F. Müller,1773*G. sociale* (Dujardin,1841) Warming,1876= *Tetramonas socialis* Dujardin,1841
* Freshwater species, which does not tolerate the full salinity of the area.
Gulf of Finland, Gdańsk, German Baltic*G. pectorale* O.F. Müller,1773 (TM)* Freshwater species, which does not tolerate the full salinity of the area.
Gdansk, German Baltic*Pandorina* Bory,1824, emend. Ehrenberg,1830*P. morum* (O.F. Müller,1773) Bory,1824= *Volvox morum* O.F. Müller,1773 (TM)
* Freshwater species, which does not tolerate the full salinity of the area.
Gulf of Finland, Gdańsk*Eudorina* Ehrenberg,1832*E. elegans* Ehrenberg,1832 (TM)* Freshwater species, which does not tolerate the full salinity of the area.
Gulf of Finland, Gdańsk*E. unicocca* G.M. Smith,1931* Freshwater species, which does not tolerate the full salinity of the area.
Gulf of Finland*Pleodorina* Shaw,1894*P. californica* Shaw,1894 (TM)
Banyuls-sur-mer, California*Volvulinidae*
* Marine?*Astrophomenaceae*
* Marine?*Pyrobotryvaceae*

* Marine?

EUSTIGMATOMONADIDA= *EUSTIGMATALES* Hibberd,1981*Monopsidaceae* Hibberd,1981*Nannochloropsis* Hibberd,1981 (TM) *N. oculata* (Droop) Hibberd,1981

N. sp. : Edler, Hällfors & Niemi,1984

Gulf of Finland

PRASINOMONADIDA= *PRASINOPHYCEAE* Christensen,1962, sensu Moestrup & Thronsdæn,1988 + *LOXOPHYCEAE*

Christensen,1962

* All Pankow,1990 inkludert

PEDINOMONADIDA Ettl,1980* This taxon is only recognized as "Chlorophytes "incertae sedis"" by Chrétiennot-Dinet,1990. Moestrup,1991 described the new class *PEDINOPHYCEAE* for species without scales, thus giving these species the same rank as the scale-bearing *PRASINOPHYCEAE*.*Pedinomonadaceae* Korschikov,1938*Micromonas* Manton & Parke,1960, non Borrel,1902 (Flagellata incertae sedis)*M. pusilla* (Butcher,1952) Manton & Parke,1960 (TM)= *Chromulina pusilla* Butcher,1952
Gulf of Finland, S Baltic, Öresund, Belt Sea, Kattegatt, Oslofjord, S, W & N Norway, North Sea, Britain, Biscay*Pedinomonas* Korschikov,1923 (TM) *P. minor* Korschikov,1923*P. minutissima* Skuja,1956

Oslofjord, Fresh water

P. sp. : Edler, Hällfors & Niemi,1984

Gulf of Finland

Resultor Moestrup,1991*R. mikron* (Thronsdæn,1969) Moestrup,1991= *Pedinomonas mikron* Thronsdæn,1969
Baltic (at >6 ppt S), Kattegatt, Oslofjord, Japan, Australia*MAMIELLALES* Moestrup,1984*Mamiellaceae* Moestrup,1984*Mantoniella* Desikachary,1972*M. squamata* (Manton & Parke,1960) Desikachary,1972 (TM)= *Micromonas squamata* Manton & Parke,1960

Gulf of Finland, central, S & W Baltic, Öresund, Belt Sea, Kattegatt, E North Sea, Oslofjord, Bergen, British estuaries, more or less cosmopolitan

Mamiella Moestrup,1984*M. gilva* (Parke & Rayns,1964) Moestrup,1984
= *Nephroselminis gilva* Parke & Rayns,1964
Kattegatt, Oslofjord, Bergen, Plymouth, Thailand*Dolichomastix* Manton,1977*D. lepidota* Manton,1977
Kattegatt, Mediterranean, South Africa*D. nummulifera* Manton,1977 (TM)
Isefjorden, S Kattegatt, Frederikshavn, Arctic Canada, Alaska, South Africa
CHLORODENDRALES Fritsch,1918, emend. Moestrup & Thronsdæn,1988
= PYRAMIMONADALES Chadeauf,1950
= HALOSPHAERALES Christensen,1962 (n. nud.)
= TETRASELMIDALES Ettr,1982 (n. nud.)*Halosphaeraceae* Haeckel,1894, emend. Moestrup & Thronsdæn,1988*Pachysphaera* Ostenfeld, in Knudsen & Ostenfeld,1899*P. pelagica* Ostenfeld, in Knudsen & Ostenfeld,1899 (TM)
Gdansk*P.* sp. : Edler, Hällfors & Niemi,1984
Gulf of Finland*Halosphaera* Schmitz,1879*H. viridis* Schmitz,1879 (TM - Mediterranean)
=? *Cryptomonas pelagica* Lohmann,1903 (motile phase)
Central & W Baltic, Öresund, Kattegatt, Bohuslän, ?Oslofjord, Plymouth, Biscay, Mediterranean*H. minor* Ostenfeld,1899
?W Baltic, Engl. Channel, Biscay, Mediterranean*H. russelli* Parke, in Parke & den Hartog-Adams,1965
SW Britain, Biscay*Collodictyon* Carter,1865
* Genus incertae sedis. Classification very tentative.*C. triciliatum* Carter,1865?
Plymouth*Cymbomonas* Schiller,1913 (TM) *C. tetramitiformis* Schiller,1913 - Mediterranean)
= *Schilleriomonas* Iyengar & Desikachary,1976*C. tetramitiformis* Schiller,1913
Isefjorden, Frederikshavn, Mediterranean, Japan, Australia*Pterosperma* Pouchet,1893 (TM) *P. rotundum* Pouchet,1893
= *Coccoperma* Silva,19
= *Cysta* Cleve,
= *Cymatosphaera* O. Wetzel,
= *Pterococcus* Lohmann,19
= *Pterocystis* Lohmann,19
= *Pterospermella* Eisenach,
= *Pterospermopsis* W. Wetzel,
= *Pterosphaera* (Jörgensen) Lohmann,19*P. cristatum* Schiller,1926
Gulf of Finland, Belt Sea, S Kattegatt, Limfjorden, Skagerrak, Oslofjord, E North Sea*P. cuboides* Gaarder,1954
Oslofjord*P. moebii* (Jörgensen) Ostenfeld,19
Skagerrak, Oslofjord*P. vanhoeffennii* (Jörgensen) Ostenfeld, in Knudsen & Ostenfeld,1899?
N Kattegatt, Skagerrak, Oslofjord, E North Sea*P. dictyon* (Jörgensen) Ostenfeld, in Ostenfeld & Schmidt,19
Skagerrak*Pyramimonas* Schmarda,1850 (TM) *P. tetrarhynchus* Schmarda,1849 - fresh water)
= *Pyramidomonas* von Stein,18*P. (Trichocystis* ??,1???) *grossii* Parke,1949
W Baltic, Kattegatt, Limfjorden, Oslofjord, S Norway, Bergen, Tromsø, Bjørnøya, E North Sea, Plymouth, Mediterranean, Arctic, Japan, Australia*P. (Trichocystis* ??,1???) *cirralae* Pennick,1982
Kattegatt, Norfolk, Norw. Sea, Greenland, Australia*P. (Trichocystis* ??,1???) *nansenii* Braarud,1935
Isefjorden, E Greenland, Arctic Canada*P. (Vestigifera* ??,1???) *orientalis* Butcher,1959
Bothnian Sea, Gulf of Finland, N & W Baltic, Öresund, Kattegatt, ?Oslofjord, S, W & N Norway, E North Sea, Mediterranean, W North America, Greenland, Australia*P. (Vestigifera* ??,1???) *mantonae* Moestrup & Hill,1992
Isefjorden, Australia, South Africa*P. (Vestigifera* ??,1???) *norrisii* Sym & Pienaar,1991
Isefjorden, Kattegatt, South Africa*P. (Vestigifera* ??,1???) *disomata* Butcher,1959
S Kattegatt, Oslofjord, S Norway, Bergen, Tromsø, Essex, Norfolk, Scotland, W North America, Japan, Australasia*P. plurioculata* Butcher,1959
* Probably identical with *P. disomata* Butcher.
Oslofjord, Tromsø, Plymouth*P. (Vestigifera* ??,1???) *cordata* McFadden, Hill & Weatherbee,1986
S Kattegatt, South Africa, Australia*P. (Tetrarhynchus* ??,1???) *longicauda* van Meel,1969
= *Pyramimonas ostendensis* van Meel,19??
Kattegatt, Belgium, Japan, Australia*P. (Tetrarhynchus* ??,1???) *octopus* Moestrup & Aa. Kristiansen,19??
Öresund, Kattegatt, Vadehavet*P. (Tetrarhynchus* ??,1???) *propulsa* Moestrup & Hill,1991

- Limfjorden, England, France, Italy, South Africa, Australia, Japan
P. (Virginica) ??,??? *virginica* Pennik,1977
 N Baltic, ?Isefjorden, Virginia, W Greenland, Australia, New Zealand
- P. obovata* Carter,19
 Kattegatt, Limfjorden
- P. octociliata* N. Carter,19??
 Plymouth
- Chlorodendraceae Oltmanns,1904, emend. Moestrup & Thronsen,1988
 = Tetraselmiidae Christensen,1962 (n. nud.)
 = Prasinocladaceae Ettl,1966
- Tetraselmis* von Stein,1878
 = *Prasinocladus* Kuckuck,1894
 = *Platymonas* G.S. West,1916?
 = *Aulacochlamys* Margalef,19
 = *Pyramischlamys* H. & O. Ettl,19 (p.p.)
- T. marina* (Cienkowski,1881) Norris, Hori & Chihara,1980
 = *Chlorangium marinum* Cienkowski,1881
 = *Prasinocladus marinus* (Cienkowski,1881) Waern,1952
 = *Prasinocladus lubricus*
 * This may be the only marine species, belong in *Prasinocladus* Kuckuck, if that genus will be reestablished by Moestrup & Thronsen (in prep.)
 Baltic S of Öresund
- T. cordiformis* (Carter,1859) von Stein,1878 (TM)
 = *Cryptoglena cordiformis* Carter,1859
 = *Carteria cordiformis* (Carter,1859) Diehl,1896
 = *Platymonas tetrathalos* G.S. West,1916?
 Gdańsk, ?N Kattegatt, ?Bergen
- T. suecica* (Kylin,1935) Butcher,1959
 = *Platymonas suecica* Kylin,1935
 Belt Sea
- T. gracilis* (Kylin,1935)
 = *Platymonas gracilis* Kylin,1935
 W Sweden, Oslofjord, Engl. Channel
- T. rubens* Butcher,1959
 Oslofjord, Norfolk
- T. convolutae* (Parke & Manton,1967) Norris, Hori & Chihara,1980
 = *Platymonas convolutae* Parke & Manton,1967
 @ *Convoluta roscoffensis* Graff
 NW France, ?S Wales
- T. carteriiformis* Butcher,1959
 Britain
- T. chui* Butcher,1959
 Britain
- T. roscoffensis* (Dangeard,1931)
 = *Platymonas roscoffensis* Dangeard,1931
 Roscoff
- T. contracta* (N. Carter,1937) Butcher,19??
 = *Platymonas contracta* N. Carter,1937
 Limfjorden, Isle of Wight, Essex
- Pseudoscourfieldia* Manton,1975
- P. marina* (Thronsen,1969) Manton,1975 (TM)
 = *Scourfieldia marina* Thronsen,1969
 Gulf of Finland, Belt Sea, Kattegatt, Oslofjord, S Norway, Bergen, South Africa, Australia
- Nephroselmis* von Stein,1878
 = *Heteromastix* Korschikov,1923
 = *Anisomonas* Butcher,19
 = *Bipedinomonas* Carter,19
 = *Sennia* Pascher,19 (p.p.)
- N. olivacea* von Stein,1878 , emend. Moestrup & Ettl,19?? (TM)
 = *Heteromastix angulata* Korschikov,1923 (TM of *Heteromastix* Korschikov,1923)
 = *Nephroselmis angulata* (Korschikov,1923) Skuja,19??
 * Freshwater species, which does not tolerate the full salinity of the area.
 Gulf of Finland
- N. pyriformis* (N. Carter,1937) Ettl,19??
 = *Bipedinomonas pyriformis* Carter,1937
 = *Heteromastix pyriformis* (Carter,1937) Manton, in Parke & Dixon,1964
 = *Nephroselmis longifilis* (Butcher,1959) Norris,19??
 = *Heteromastix longifilis* (Butcher,1959) Rayns,19??
 Baltic, Kattegatt, Oslofjord, S W & N Norway, Engl. Channel, Massachusetts
- N. rotunda* (N. Carter,1937) Fott,19??
 = *Heteromastix rotundata* (N. Carter,1937) Manton,19??
 N Kattegatt, Bergen, Isle of Wight, Plymouth, Australia
- N. minuta* (N. Carter,1937) Butcher,19??
 = *Heteromastix minuta* N. Carter,1937
 Baltic, N Kattegatt, Isle of Wight
- Chlorodendrales unknown family
- Asteromonas* Artari,1926
- A. gracilis* Artari,1926
 S Kattegatt, Germany
- SILICOFLAGELLIIDA Borgert,1891
 = DICTYOCHALES Haekel,1894
 * All Pankow,1990 included.
- Dictyochaceae Lemmermann,1901 = Dictyochidae
- Dictyocha* Ehrenberg,1837
 = *Distephanus* Stöhr,1880
 = *Octacis* Schiller,19
 = *Camponotus* Haekel,18
- D. speculum* Ehrenberg,1837
 * Perhaps toxic to fish.
 All Baltic, Öresund, Kattegatt, Skagerrak, Oslofjord, E North Sea, Plymouth
- D. fibula* Ehrenberg,1837
 Liverpool Bay
- EUGLENOZOA Cavalier-Smith,1981
- EUGLENIDA Bütschli,1884
 = EUGLENOPHYCEAE Schoenichen,1925 = EUGLENALES Engler,1898
 * Pankow,1990 totally included
- Calkinsia* Lackey,1960
 * Which suborder?
- C. aereus* Lackey,1960 (TM)
 * Sediment-living marine species.
 ?
- Dolium* Larsen & Patterson,1990
 * Which suborder? Only recorded from marine tropical sediments.
- Gyropagiae* Skuja,1939
 * Which suborder? Only freshwater. Few species.
- EUTREPTIINA Leedale,1967
 = EUTREPTIACEAE Hollande,1942
- Eutreptiella* da Cunha,1914 (TM *E. marina* da Cunha,1914)
 = *Gymnastica* Schiller,19
- E. gymnastica* Thronsen,1969
 Gulf of Finland, Limfjorden, S Kattegatt, Oslofjord, Bergen
- E. braurudii* Thronsen,1969
 S Kattegatt, Limfjorden, Oslofjord
- E. hirudoidea* Butcher,1961
 Tromsö, England
- E. euphyrynea* Walne, Moestrup, Norris & Ettl,1986
 Little Belt
- E. sp. : Edler, Hälfors & Niemi,1984*
 Gulf of Finland
- Eutreptia* Perty,1852
- E. viridis* Perty,1852 (TM)
 German Baltic, Plymouth
- E. lanowii* Steuer,1904
 Oder estuary, Arkona Sea, Belt Sea, Limfjorden, Plymouth
- E. sp. : Edler, Hälfors & Niemi,1984*
 Gulf of Finland
- Distigma* Ehrenberg,1838
 = *Distigmopsis* Hollande,1942 ?= *Ampullomonas* Skvortzov,1957
 ?= *Baranovia* Skvortzov,1957 ?= *Schewiakowia* Skvortzov,1957
 * Colourless. Not mentioned by Chrétiennot-Dinet,1990. Only recorded from freshwater
- EUGLENINA Bütschli,1884
- Euglena* Ehrenberg,1830
 * *Euglenaceae* von Stein,1878-83, emend. Klebs,1883, according to Pankow,1990
- E. viridis* (O.F. Müller,1786) Ehrenberg,1830 (TM)
 = *Cercaria viridis* O.F. Müller,1786
 * Freshwater species, which does not tolerate the full salinity of the area.
 * Main occurrence in eutrophied waters.
 Helsinki harbour, Tärrämäen, Bretagne
- E. triptera* (Dujardin,1841) Klebs,1883
 = *Phacus triptera* Dujardin,1841
 * Freshwater species, which does not tolerate the full salinity of the area.
 German Baltic
- E. oxyuris* Schmarda,1850
 * Freshwater species, which does not tolerate the full salinity of the area.
 Gulf of Finland
- E. spirigera* Ehrenberg,1830
 * Freshwater species, which does not tolerate the full salinity of the area.
 German Baltic
- E. proxima* Dangeard,1901
 = *E. ballica* Schüller,1910
 Kieler Bucht
- E. acus* Ehrenberg,1830
 * Freshwater species, which does not tolerate the full salinity of the area.
 German Baltic
- E. reticulata* Sjöstedt,1925
 * Not recognizable, sensu Pringsheim,1953
 W Baltic
- E. deses* Ehrenberg,1830
 E North Sea
- E. fenestrata* Elenkin,19??
 Plymouth
- E. vermiformis* N. Carter,19??
 Plymouth
- E. obtusa* Schmitz,19??
 * in estuarial mud
 Britain
- Protoeuglena* Subrahmanyam,1954
 @ *Nocilluca scintillans*
 * the Mediterranean form may be another species
 Indian Ocean, ?Mediterranean
- P. noctilucae* Subrahmanyam,1954
 @ *Nocilluca scintillans*
 * the Mediterranean form may be another species
 Indian Ocean, ?Mediterranean
- Astasia* Dujardin,1841 (TM *A. limpida* Dujardin,1841 - fresh water)
 ?= *Acusmonas* Skvortzov,1969 ?= *Astasiella* Michajlow,1964, non Skvortzov,1957
 ?= *Atractonema* von Stein,1878 ?= *Bicudomonas* Skvortzov,1969
 ?= *Hoehnemonas* Skvortzov,1969 ?= *Kolbeana* Skvortzov,1966
 ?= *Liangiana* Skvortzov,1969 ?= *Liuiana* Skvortzov,1969
 ?= *Paramylonomas* Skvortzov,1957 ?= *Parastasiella* Michajlow,1965
- A. sp. : Edler, Hälfors & Niemi,1984*
 Gulf of Finland

- A. doridis** Zerling,1???
- * @ oviparosite of *Archidoris tuberculata*
- * incertae sedis?
- Cyclidiopsis** Korschikow,1917
- * Sometimes recognized as a subgenus of *Astasia* Dujardin
- Euglenopsis** Klebs,1892
- * Sometimes recognized as a subgenus of *Astasia* Dujardin
- Khwakinea** Jahn & McKibben,1937
- * Sometimes recognized as a subgenus of *Astasia* Dujardin
- Pseudastasia** Bourrelly,1985
- * Normally recognized as a subgenus of *Astasia* Dujardin
- Colacium** Ehrenberg,1838
- * Colaciaceae G.M. Smith,1933, according to Pankow,1990
- * Not mentioned by Chrétiennot-Dinet,1990
- C. vesiculosum** f. *vesiculosum* Ehrenberg,1838
- @ epizoic on *Acartia* spp., *Temora longicornis*, *Centropages hamatus*, *Pseudocalanus* sp.
- * Freshwater species, which does not tolerate the full salinity of the area.
- Helsinki, W Baltic, Öresund, Belt Sea
- C. vesiculosum** f. *arbucula* (von Stein,1878) Huber-Pestalozzi,1929
- = *C. arbucula* von Stein,1878
- * Freshwater species, which does not tolerate the full salinity of the area.
- Gulf of Finland
- Hyalophacus** Pringsheim,1936
- * Sometimes considered as a colourless subgenus of *Phacus* Dujardin,1841
- Lepocinclus** Perty,1849 (TM *L. globulus* Perty,1849)
- * Euglenaceae von Stein,1878-83, emend. Klebs,1883, according to Pankow,1990
- L. ovum** var. *ovum* Ehrenberg,18???, emend. Lemmermann,1901
- ?Gulf of Finland, Rügen
- L. ovum** var. *major* (Huber-Pestalozzi,1929) Conrad,1934
- = *L. butschlii* var. *major* Huber-Pestalozzi,1929
- German Baltic
- L. sp.** : Edler, Hälfors & Niemi,1984
- * Freshwater species, which does not tolerate the full salinity of the area.
- Gulf of Finland
- Phacus** Dujardin,1841
- * Euglenaceae von Stein,1878-83, emend. Klebs,1883, according to Pankow,1990
- P. pyrum** (Ehrenberg,1838) von Stein,1878-83
- = *Euglena pyrum* Ehrenberg,1838
- * Freshwater species, which does not tolerate the full salinity of the area.
- ?Gulf of Finland, S Baltic
- P. pleuronectes** (O.F. Müller,1786) Dujardin,1841
- = *Cercaria pleuronectes* O.F. Müller,1786
- * Freshwater species, which does not tolerate the full salinity of the area.
- German & S Baltic
- P. longicauda** (Ehrenberg,1838) Dujardin,1841 (TM)
- = *Euglena longicauda* Ehrenberg,1838
- * Freshwater species, which does not tolerate the full salinity of the area.
- Gulf of Finland
- Klebsiella** Pascher,1931 (TM *K. alligata* Pascher,1931)
- Atraktomonas** Christen,1962
- * Freshwater; monotypic
- Calymonoma** Christen,1959
- * Only freshwater. several species
- Dylakosoma** Skuja,1964
- * Only Freshwater. Two species.
- Parmidium** Christen,1962
- * Only freshwater. Two species.
- Petalomonas** von Stein,1859
- = *Copromonas* Dobell,1908 = *Huber-Pestalozziomona* Skvortzov,1967
- = *Pentamonas* Lackey,1962 = *Scytonomas* Stein,1878
- = *Thylakomonas* Schweiakoff,1892 = *Triangulomonas* Lackey,1940
- P. abeissa** (Dujardin) von Stein,1859 (TM)
- E North Sea, fresh water
- P. minuta** Hollande,1942
- Tvärminne, S Baltic, Kattegatt. E North Sea, N midAtlantic, Fiji, Australia, Brazil
- P. spinifera** (Lackey,1962?)
- = *Pentamonas spinifera* Lackey,1962?
- * sand- and mud-living.
- Plymouth, E & W USA
- P. pusilla** Skuja,1948
- * sand- and mud- & sediment-living.
- Tvärminne, E North Sea, Plymouth
- P. excavata** Skuja,19???
- * Sediment living.
- Latvia, Plymouth
- P. medicocanellata** Skuja,19???
- * sand- and mud- & sediment-living. Widely distributed in fresh water.
- Plymouth
- P. pusilla** (von Stein,1878?)
- = *Scytonomas pusilla* von Stein,1878?
- Plymouth
- Strombomonas** Deflandre,1930
- * Trachelomonadaceae ?1???, according to Pankow,1990
- * Not mentioned by Chrétiennot-Dinet,1990
- = *Trachelomonas deflandrei* Roll,1928
- * Freshwater species, which does not tolerate the full salinity of the area.
- Gulf of Finland
- S. deflandrei** (Roll,1928) Deflandre,1930
- = *Trachelomonas deflandrei* Roll,1928
- * Freshwater species, which does not tolerate the full salinity of the area.
- Gulf of Finland
- T. volvocina** Ehrenberg,1833
- * Freshwater species, which does not tolerate the full salinity of the area.
- Gulf of Finland
- T. hispida** (Perty,1852) von Stein,1878, emend. Deflandre,1926
- = *Chonemonas hispida* Perty,1852
- * Freshwater species, which does not tolerate the full salinity of the area.
- Gulf of Finland
- T. abrupta** Swivenko, sensu Dragesco,19???
- * in brackish water sand
- Bretagne
- RHABDOMONADINA** Leedale,1967
- Rhabdomonas** Fresenius,1858
- * Only freshwater. Several species.
- Menoidium** Perty,1852
- = *Menoidionomas* Skvortzov,1957 ?= *Calawaymonas* Skvortzov,1969
- ?= *Nitzamonas* Skvortzov,1969 ?= *Skijamonas* Skvortzov,1969
- ?= *Wellstella* Skvortzov,1969 ?= *Zimmermanniella* Skvortzov,1969
- * Only freshwater.
- Rhabdospira** Pringsheim,1942 (nomen nudum)
- * Not formally erected for certain species within *Menoidinium* Perty
- SPHENOMONADINA** Leedale,1967
- * No longer separable from Heteronomatina, sensu Larsen & Patterson,1992
- Sphenomonas** von Stein,1878
- * Not mentioned by Chrétiennot-Dinet,1990
- S. elongata** Lackey,19???
- * sand- and mud-living.
- Plymouth, E & W USA
- S. quadrangularis** von Stein,18???
- * In sediments. Also in fresh water.
- Plymouth
- Notosolenus** Stokes,1884
- = *Solenotus* Stokes,1884
- N. cancellatus** Skuja,19
- E North Sea
- N. esulcis** Larsen,19
- E North Sea
- N. lagenos** Skuja,19
- ? E North Sea
- N. rhombicus** Larsen,19
- E North Sea
- N. apocampatus** Stokes,1???
- * sand- and mud-living. Fresh water in USA.
- Plymouth
- Anisonema** Dujardin,1841
- ?= *Aliniella* Skvortzov,1957 ?= *Fidalgomonas* Skvortzov,1969
- ?= *Mattosmonas* Skvortzov,1969
- A. acinus** Dujardin,1841 (TM)
- E North Sea
- A. prosgeobium** Skuja,1939
- Tvärminne, Latvia, S Baltic, Kattegatt, Wadden Sea, Arctic Canada, Fiji, Hawaii, Australia
- A. trepidum** Larsen,19
- E North Sea
- A. emarginatum** Stokes,1???
- * in sand & mud. Freshwater in USA
- Plymouth
- A. ovale** Klebs,1???
- * sand- and mud-living. Also in fresh water.
- Plymouth
- A. pusillum** Stokes,1???
- * sand- and mud-living. Fresh water in USA.
- Plymouth
- Metanema** Senn,1900
- M. sedatum** Larsen,19
- E North Sea
- M. strenuum** (Skuja) Larsen,19
- E North Sea
- M. variabile** Klebs,1???
- Plymouth
- Tropidoscyphus** von Stein,1878
- T. octocostatus** von Stein,18???
- * sand- and mud-living. Also in fresh water.
- Plymouth
- T. sp.**
- E North Sea
- HETERONEMATINA** Leedale,1967
- * No longer separable from Sphenomonadina, sensu Larsen & Patterson,1992
- Heteronema** Dujardin,1841
- = *Zygoselmis* Dujardin,1841
- * Not mentioned by Chrétiennot-Dinet,1990
- H. ovale** Kahl,19
- E North Sea
- H. acus** (Ehrenberg,18???) von Stein,18???
- * in sediment. Also in fresh water.
- Plymouth

- Entosiphon*** von Stein,1878 (ICZN), non Beddome,1864 (ICBN)
 = ***Entosiphonomas*** Larsen, & Patterson,1992 (ICBN)
 ?= *Marsupiogaster* Schewiakoff,1893
 * Not mentioned by Chretiennot-Dinet,1990
- E. limenites*** Norris,1964
 Tromsö, Pacific Ocean
- E. appanatum*** Preisig,19
 ?E North Sea
- E. cuneatiformis*** Lackey,1962
 In sand
 USA
- Dinema*** Perty,1852 (ICZN)
 = ***Dinematomonas*** Silva,1960 (ICBN)
- D. litorale*** (Skuja,19??) Silva,19??
 * sand-living.
 E North Sea, Plymouth
- D. grisoleum*** (Perty,18??)
 = *Dinema grisoleum* Perty,18??
 * sand-living. Chiefly fresh water living.
 Plymouth
- Ploetotia*** Dujardin,1841
 = *Pseudamoeba* Behre,1961 = *Serpenomonas* Triemer,1986
 * Mostly marine, with unclear boundaries with *Anisonema* & *Entosiphon*
- P. corrugata*** Larsen & Patterson,1990
 N midAtlantic
- Peranema*** Dujardin,1841 (ICZN) (preoccupied in ICBN)
 = ***Pseudoperanema*** Christen,1962 (ICBN)
 ?= *Astasiella* Skvortzov,1957, non Michajlow,1964 ?= *Baikowia* Skvortzov,1957
 ?= *Jenningsia* Schaeffer,1918 ?= *Peranemopsis* Lackey,1940
- P. fusiforme*** Larsen,19
 E North Sea
- P. trichophorum*** (Ehrenberg,18) Larsen,19
 * In sediment. Widely distributed in freshwater.
 E North Sea
- P. cuneatum*** (Playfair,1???)
 = *Peranema cuneatum* Playfair,1???
 Plymouth, freshwater in Australia
- Urceolus*** Mereschkowsky,1877
 = *Phialonema* von Stein,1878 = *Urceolopsis* Stokes,1887
- U. costatus*** von Stein,18??
 E North Sea
- U. cyclostomus*** (von Stein,18??) Mereschkowsky,1877?
 * sand- and mud-living. Also in fresh water.
 E North Sea, Plymouth
- U. pascheri*** Skvortzov,19??
 * sand- and mud-living. Also in fresh water.
 E North Sea, Plymouth
- U. sabulosus*** Stokes,19??
 * sand- and mud-living. Widely distributed in fresh water.
 E North Sea, Plymouth
- KINETOPLASTIDA** Honigberg,1963 emend. Vickerman,1976
- BODONINA** Holland,1952 Emend. Vickerman,1976
- Diplomastix** Kent,1880
 * Marine? Which family
- Bodonidae**
 = Bodonacean Lemmermann,1914
- Trypanophis*** Keyselitz,1904
 @ Siphonophora & Chaetognatha
 * Correctly classified?
- T. grobbeni*** (Poche,1903?)
 = *Trypanosoma grobbeni* Poche,1903?
 @ *Chelopes appendiculata* Eschscholtz, *Nanomia bijuga* & other siphonophores
 Mediterranean
- T. major*** Duboscq & Rose,1926
 @ *Abylopsis pentagona*
 ?France
- Bodoninae**
- Bodo*** Ehrenberg,1832
 = *Prowazekia* Hartmann & Chagas,1910
- B. ovatus*** (Dujardin,1841) von Stein,1878
 = *Heteromita ovata* Dujardin,1841
 Finland
- B. curvifilis*** Griesmann,1913
 Tärminne, Denmark, Bergen, Tromsö, Greenland, midAtlantic
- B. designis*** Skuja,1948
 Tärminne, Denmark, midAtlantic, widespread in arctic, temperate & tropic salt & fresh waters
- B. saliens*** Larsen & Patterson,1990
 Tärminne, Denmark, England, Greenland, midAtlantic, Arctic Canada, Tenerife, Belize, Brazil, Hawaii, Australia
- B. saltans*** Ehrenberg,1832 (TM)
 ?= *Pleuromonas jaculans* Perty,1852
 Tärminne, Denmark, Plymouth, midAtlantic, Belize, widespread in fresh water & soil
- B. edax*** Klebs,1???
 Plymouth
- B. globosus*** von Stein,18??
 Plymouth
- B. insidiosus*** Hollande & Enjumet,1953 (nomen nudum?)
 @ in protoplasm of radiolarian genus *Collozoum* or ?freeliving
- Ichthyobodo*** Pinto,1928 (TM) *Costia necatrix* Henneguy,1888 - fresh water)
- I.*** sp. : Bullock & Robertson,1928
 @ *Pleuronectes platessa*
 * the authors identified this population with the type species, *I. necatrix* (Henneguy,1888), which however is a fresh water species, with no affinity for NaCl-solutions
 Scotland
- Pleuromonas*** Perty,1852
- P. jaculans*** Perty,1852 (TM)
 ?
- Rhynchobodo*** Lackey,1940, emend. Vørs,1992
- R. taeniatata*** (Skuja,1956) Vørs,1992
 = *Cryptaulax taeniatata* Skuja,1956 (TM)
 Tärminne, Swedish fresh water
- R. agilis*** Lackey,1940
 = *R. armata* Brugerolle,1985
 = *Phylloimitus apiculatus* : Mylnikov,1986, non Skuja,1948
 ?
- Rhynchomonas*** Klebs,1892
- R. nasuta*** (Stokes,1888) Klebs,1892
 = *Heteromita nasuta* Stokes,1888
 Tärminne, Oslofjord, Bergen, Tromsö, Plymouth, Greenland, N midAtlantic, Arctic Canada, N America, Brazil, Australia, Fiji, Hawaii
- R. mutabilis*** Griessmann,1???
- Cryptobiinae**
- Dimastigella*** Sandon,1928
 * Only one ?soil-living species
- Cryptobia*** Leidy,1846
 = *Trypanoplasma* Laveran & Mesnil,1901 (p.p.)
 @ in freshwater invertebrates or freshwater & marine fish
- C. dahli*** (Müller,1888) Alexeieff,1912
 @ stomach of *Cyclopterus lumpus*
 Baltic, Belt Sea, Kattegatt, Skagerrak, North Sea
- C. congeri*** (Elmhirst & Martin,1910)
 @ stomach of *Conger niger* (= *Conger conger*)
 E Scotland
- C. intestinalis*** Léger,1905
 @ *Box boops*
 Mediterranean
- C. salmonis*** Katz,1951
 @ *Salmo gairdneri* & other pacific salmonids (transmitted by leeches)
 ?
- C. antedonae*** Lecal,1980
 @ coelomic fluid of *Antedon bifida*
 ?
- C. bialata*** Ruinen,1938
 * Free-living
 ?
- Procryptobia*** Vickerman,1978
- Cephalothamniidae**
- Cephalothamnum*** Stein,1878
- TRYPANOSOMATINA** Kent,1880
- Trypanosomatidae** Doflein,1911
- Leptomonas*** Kent,1880
 @ no host change, chiefly intestinal parasites
- L. patellae*** Porter,1914
 @ intestinal tract & digestive gland of *Patella vulgata*
 ?
- L. karyophilus*** Gillies & Hanson,1963
 @ *Paramecium trichium*
 ?
- L.*** sp. : Chatton,1924
 @ a nematode species
 Roscoff
- Trypanosoma*** Gruby,1843
 @ host changing blood parasites
- T. murmanensis*** Nikitin,1927
 @ *Gadus morhua* Cottidae & several other fish species ; vector : *Johanssonia* sp. & *Myzobdella* sp.
 Murmansk, E North America
- T. raiiae*** Laveran & Mesnil,1902
 @ *Raja* spp. ; vector : *Pontobdella muricata*
 Shetlands, Engl. Channel
- T. giganteum*** Neumann,1909
 @ *Raja oxyrinchus* ; vector : *Pontobdella muricata*
 Mediterranean
- T. scyllii*** Laveran & Mesnil,1902
 @ *Scyliorhinus* spp. ; vector : *Pontobdella muricata*
 Engl. Channel
- T. callionymi*** Brumpt & Lebailly,1904
 @ *Callionymus lyra* ; vector : ?
 Engl. Channel
- T. cataphracti*** Henry,1913
 @ *Agonus cataphractus* ; vector : ?
 Plymouth
- T. granulosum*** Laveran & Mesnil,1902
 @ *Anguilla anguilla* ; vector: *Hemiclepsis marginata* & *Piscicola geometra*

- Engl. Channel
- T. laternae** Lebailly,1904
@ *Arnoglossus laterna* ; vector: ?
Shetlands, Engl. Channel
- T. delagei** Brumpt & Lebailly,1904
@ *Blennius pholis* ; vector: ?
Engl. Channel
- T. bothi** Lebailly,1905
@ *Bothus rhombus* ; vector: ?
Engl. Channel
- T. cotti** Brumpt & Lebailly,1904
@ *Taurulus bubalis* ; vector: *Trachelobdella punctata*
Engl. Channel
- T. limandae** Brumpt & Lebailly,1904
@ *Hippoglossoides platessoides* ; vector: ?
Engl. Channel
- T. aeglefini** Henry,1913
@ *Melanogrammus aeglefinus* ; vector: ?
Shetlands
- T. gobii** Brumpt & Lebailly,1904
@ *Gobius niger* ; vector: ?
Engl. Channel
- T. flesi** Lebailly,1904
@ *Platichthys flesus* ; vector: ?
Engl. Channel, Millport
- T. platessa** Lebailly,1904
@ *Pleuronectes platessa* ; vector: ?
Engl. Channel, Millport
- T. soleae** Laveran & Mesnil,1901
@ *Solea solea* ; vector: *Trachelobdella punctata* (= *Callobdella p.*) & other leeches
Engl. Channel
- T. yakimovi** Wladimiroff,1910
@ *Syngnathus acus* ; vector: ?
Gulf of Naples
- T. trigiae** Neumann,1909
@ *Triglidae* e.g. *Eurigla gurnardus* ; vector: ?
Shetlands, Mediterranean
- T. zeugopteri** Henry,1913
@ *Zeugopterus punctatus* ; vector: ?
Isle of Man
- T. testudinis** ??,???
@ *Chelonia mydas*
?
- ZOOMASTIGOPHOREA** Calkins
- Bodomorpha** Hollande, in Grassé,1952 (TM *Pseudobodo minima* Hollande,1942)
= *Pseudobodo* Hollande,1942, non Griessmann,1913
- Pseudospora** Cienkowski,1865
Correct classis? Which order & family?
- ?P. sp. : Parke & den Hartog-Adams,1965
@ *Halosphaera* spp.
SW Britain, Biscay, Mediterranean
- CHOANOFAGELLIDA** Kent,1880
= CRASPEDOPHYCEAE (Chadefaud,1960) Christensen,1962
= CRASPEDOMONADALES von Stein,1878
= STYLOCHROMONADALES
* All Pankow,1990 included.
- Codonosigidae* Kent,1880
= *Codonosigaceae* Kent,1880
- Astrosiga** Kent,1878
- Cladospongia** Iyengar & Ramanathan,1940
- Codosiga** James-Clark,1868
- C. gracilis** (Kent) de Saedeleer
Parke & Leadbeater,1977 area
- C. pyriformis** Kent,1880
Parke & Leadbeater,1977 area
- C. botrytis** (Ehrenberg,18??) Kent,1880
Tårmimne, Germany, Denmark, Italy, temperate fresh waters
- Codonocladium** von Stein,1878
* Considered to be a synonym of *Codosiga* James-Clark, by most authors
- C. cymosum** (Kent) Boucaud-Camou,1967?
Parke & Leadbeater,1977 area
- Desmarella** Kent,1878
= *Codonodesmus* von Stein,1878
= *Kentia* Schiller,1953
- D. moniliformis** (Kent,1878)
Oslofjord, Plymouth
- Kentrosiga** Schiller,1953
- K. thienemanni** Schiller,19
Parke & Leadbeater,1977 area
- Monosiga** Kent,1882
- M. ovata** Kent,1882
* Considered a synonyme of *Codosiga gracilis* (Kent,1880-82) de Saedeleer,1927, by Thronsen,1974
Bergen, Plymouth, Luc-sur-Mer
- M. brevicollis** Ruinen,1938
Oslofjord
- M. lagunula** Thronsen,1969
Oslofjord, W Norway
- M. marina** var. *marina* Grönvæd,1952
Skagerrak, North Sea, Norw. Sea, N Norway
- M. marina** Grönvæd var. *minima* Paasche,1961
W Norway, Norw. Sea
- M. mikropelagica** Thronsen,1974
Oslofjord, Bergen
- M. sinuosa** Möbius,18
Parke & Leadbeater,1977 area
- Proterospongia** Kent,1882 (TM *Protospongia haecellic* Kent,1881 - fresh water)
- P. dybsoensis** (Grönvæd,1956) Loeblich III,1967
= *Protospongia dybsoensis* Grönvæd,1956
Dybø Fjord (Denmark)
- P. nana** (Braarud,1935) Tuzet,1963
= *Protospongia nana* Braarud,1935
E Greenland
- P. choanojuncta** Leadbeater,1983
Lincolnshire
- Sphaerodendron** Zhukov,1978
- Sphaeroeca** Lauterborn,1894 (TM *S. volvox* Lauterborn,1894 - fresh water)
* Possibly a synonym of *Proterospongia* Kent
- S. salina** Bourcet,1947
* pelagic brackish water species
?France
- Codonosigopsis** Senn, in Engler & Prantl,1900
= *Dicraspedella* Ellis,1930
- C. stokesii** (Ellis,1930)
= *Dicraspedella stokesii* Ellis,1930
Oslofjord
- Diplosiga** Frenzel,1891
- Salpingoecidae** Kent,1880
= *Salpingoecaceae*
- Choanoeca** Ellis,1930
- C. perplexa** Ellis,1930
Parke & Leadbeater,1977 area
- Diploeca** Ellis,1930
- Pachysoeca** Ellis,1930
* Considered to be a synonym of *Salpingoeca* James-Clark, by some authors
- P. convallaria** (von Stein,18) Ellis,19
Parke & Leadbeater,1977 area
- P. depressa** Ellis,19
Parke & Leadbeater,1977 area
- Salpingoeca** James-Clark,1868
?= *Lagenoeca* Kent,1881
- S. aggregata** Valkanov,1970
Tårmimne, Bulgaria
- S. amphoridium** James-Clark,1867
= *S. ampullacea* Stein,1878 = *S. ampulloides* Bicudo & Bicudo,1983
Tårmimne, Denmark, Fresh water in Europe, USA & Australie
- S. collaris** Stokes,1894
Tårmimne, New Jersey fresh water
- S. inquillata** Kent,1880-82
Oslofjord, S England
- S. cardiforme** Ellis,19
Parke & Leadbeater,1977 area
- S. cornuta** Kent,18
Parke & Leadbeater,1977 area
- S. huxleyi** Ellis,1930
= *S. caudiculata* Bicudo & Bicudo,1983
Parke & Leadbeater,1977 area
- S. infusionum** Kent,1880
= *S. longipes* Kent,18
Parke & Leadbeater,1977 area, N midAtlantic
- S. marina** James-Clark,1867
Tårmimne, Denmark, England, France, USA
- S. megacheila** Ellis,19
Parke & Leadbeater,1977 area
- S. napiformis** Kent,18
= *S. desadeleri*
Parke & Leadbeater,1977 area
- S. procera** Möbius,18
Parke & Leadbeater,1977 area
- S. stenotheca** Boucaud-Camou,1966
= *S. lansi* Ertl,1970
Parke & Leadbeater,1977 area
- S. teres** Kent,18
Parke & Leadbeater,1977 area
- S. tuba** Kent,1880
= *S. petiolata* Kent,18
Parke & Leadbeater,1977 area
- S. undulata** Ellis,19
Parke & Leadbeater,1977 area
- S. urceolata** Kent,18
Parke & Leadbeater,1977 area

- Diplosigopsis* Francé, 1897
Aulomonas Lackey, 1942
Steloxomonas Lackey, 1942
Acanthoecidae Norris, 1965
= *Acanthoecaceae* Norris, 1965
Acanthocoris Hasa & Takahashi, 1984
= *Pseudomicrosporella* Scagel & Stein, 1961
A. apoda (Leadbeater, 1972) Hasa & Takahashi, 1984
= *Acanthoecopsis apoda* Leadbeater, 1972 (TM)
Tvärrminne, Denmark, Oslofjord, S Norway, Bergen, W Greenland, Mediterranean
A. campanula (Espeland, 1986) Thomsen, in & Thomsen, Buck & Chavez, 1991
= *Acanthoecopsis campanula* Espeland, in Espeland & Thronsen, 1986
Kattegatt, S Norway
A. haarakiana Thomsen, in Thomsen, Buck & Chavez, 1991
Kattegatt, California
A. unguiculata (Thomsen, 1973) Hasa & Takahashi, 1984
= *Acanthoecopsis unguiculata* Thomsen, 1973
= *Pseudomicrosporella ornata* Scagel & Stein, 1961
Tvärrminne, Isefjord, Oslofjord, S Norway, Bergen, W Greenland, midAtlantic
A. asymmetrica (Thomsen, 1979) Hasa & Takahashi, 1984
= *Acanthoecopsis asymmetrica* Thomsen, 1979
Tvärrminne, Kattegatt
A. sp. n. Thomsen, in Thomsen, Buck & Chavez, 1991
Baltic
Acanthoecea Ellis, 1930
A. brevipoda Ellis, 1930
Frederikshavn, Britain
A. spectabilis Ellis, 1930 (TM)
Tvärrminne, Isefjord, Bergen, Britain, Greenland, S France
Amoenoscopa Hara & Takahashi, 1987
A. caudata Hara & Takahashi, 1987 (TM)
Denmark, Japan
Apheloecion Thomsen, in Thomsen & Boonruang, 1983 (TM) *A. quadrispinum* Thomsen, in Thomsen & Boonruang, 1983 - tropical-subtropical seas
Bicosta Leadbeater, 1978
B. spinifera (Thronsen, 1970) Leadbeater, 1978
= *Salpingoeca spinifera* Thronsen, 1970 (TM)
Isefjord, S, W & N Norway, off E & W Bear Island, midAtlantic
B. minor (Reynolds, 1976) Leadbeater, 1978
= *Salpingoeca minor* Reynolds, 1976, non Dangeard, 1910
= *Salpingoeca virgata* Parke & Leadbeater, 1977
Tvärrminne, Denmark, S & W Norway, Arctic Seas
B. antennigera Moestrup, 1979
Denmark
Calliantha Leadbeater, 1978
C. natans (Gröntved, 1956) Leadbeater, 1978
= *Salpingoeca natans* Gröntved, 1956 (TM)
Tvärrminne, Isefjord, Fredrikshavn & Venø Bugt (Denmark), ?W Norway, Greenland
C. simplex Manton & Oates, 1979
= *Salpingoeca* sp. : Thomsen, 1973
Tvärrminne, Isefjord, Greenland, Mediterranean
C. longicaudata (Leadbeater, 1975) Leadbeater, 1978
Denmark, N Atlantic
Calotheca Thomsen & Moestrup, 1983 (TM) *Calotheca alata* Thomsen & Moestrup, 1983 - Indo-Pacific
* Monotypic warmwater genus
Campyloacantha Hara & Takahashi, 1987 (TM) *Parvicorbicula spinifera* Leadbeater, 1973
Campanoeca Thronsen, 1974
C. dilatata Thronsen, 1974 (TM)
= *Polyfibula sphyrrelata* (Thomsen, 1973) Manton & Bremer, 1981
Oslofjord
Conion Thomsen, 1982 (TM) *Conion groenlandicum* Thomsen, 1982 - W Greenland
Cosmoeca Thomsen, in Thomsen & Boonruang, 1984
C. ventricosa Thomsen, in Thomsen & Boonruang, 1984
= *Pleurasiga orculaformis* Schiller, 1925 (TM) of *Pleurasiga* Schiller, 1925
Tvärrminne, Isefjord, S Norway, Greenland, Mediterranean
C. norvegica Thomsen, in Thomsen & Boonruang, 1984 (TM)
Baltic, Denmark, N Atlantic
Crinolina Thomsen, 1976
C. iseifiordensis Thomsen, 1976 (TM)
Isefjorden, S Norway, W Greenland
Crucispina Espeland, in Espeland & Thronsen, 1986
C. cruciformis (Leadbeater, 1974) Espeland & Thronsen, 1986
= *Salpingoeca cruciformis* Leadbeater, 1974 (TM)
S Norway, Denmark, Algeria
Diaphanoeca Ellis, 1930
= *Microsporella* Scagel & Stein, 1961 (TM) *M. fiordensis* Scagel & Stein, 1961 - British Columbia
D. grandis var. *grandis* Ellis, 1930 (TM)
Tvärrminne, Isefjord, Oslofjord, S, W & N Norway, Britain, Mediterranean, Black Sea, midAtlantic
D. grandis Ellis, 1930 var. *minor* Thronsen, 1974
Oslofjord
D. sphaerica Thomsen, 1982
= *Campanoeca pedicellata* (Leadbeater, 1972) Thronsen, 1974
non = *Diphanoeca pedicellata* Leadbeater, 1972
Tvärrminne, Kieler Bucht, Denmark, Oslofjord, Bergen
D. pedicellata Leadbeater, 1972
Finland, all Baltic, Denmark
D. undulata Thomsen, 1982
Denmark, W Greenland, California
Diplotheca Valkanov, 1970
D. costata Valkanov, 1970 (TM)
Tvärrminne, Isefjord, Frederikshavn, Mediterranean, Black Sea
Kakoeca Buck & Marchant, in Buck & al., 1991 (TM) *Kakoeca antarctica* Buck & Marchant, in Buck & al., 1991 - Antarctic
Monocosta Thomsen, 1979
M. fennica Thomsen, 1979 (TM)
Tvärrminne, Denmark
Nannoeca Thomsen, 1988
N. minuta (Leadbeater, 1972) Thomsen, 1988
= *Pleurasiga minuta* (Leadbeater, 1972) Leadbeater, in Manton, Sutherland & Leadbeater, 1976
= *Pleurasiga minima* Thronsen, 1970 var. *minuta* Leadbeater, 1972 (TM)
Isefjord, Frederikshavn, S Norway, Mediterranean
Parvicorbicula Deflandre, 1960
= *Corbicula* Meunier, 1910
P. socialis (Meunier, 1910) Deflandre, 1960
= *Corbicula socialis* Meunier, 1910 (TM)
S Baltic, Belt Sea, Isefjord, Kattegatt, Bergen, off W Bear Island, E Greenland, Denmark Strait, Frans Joseph Land, Kara Sea, between Ireland & Portugal, Mediterranean
P. quadricostata Thronsen, 1970
Denmark, Oslofjord, S Norway, off W Bear Island
P. circularis Thomsen, 1976, emend. Thomsen, Buck, Coale, Garrison & Gowing, 1990
Tvärrminne, Isefjord
P. superpositus Booth, 1990
Denmark, N Pacific
P. pedunculata Leadbeater, 1980
S Bretagne
Platyleura Thomsen, in Thomsen & Boonruang, 1983
P. infundibuliformis (Leadbeater, 1974) Thomsen, in Thomsen & Boonruang, 1983
= *Parvicorbicula infundibuliformis* Leadbeater, 1974 (TM)
Denmark, S Norway
Pleurasiga Schiller, 1925 (TM) *P. orculaformis* Schiller, 1925 - Adriatic
P. minima Thronsen, 1970
Denmark, S & W Norway, off W Bear Island, Barents Sea, Sargasso Sea
P. reynoldstii Thronsen, 1970
Isefjord, W Norway, off W Bear Island, Mediterranean
P. echinocostata Espeland, in Espeland & Thronsen, 1986
Denmark, S Norway, Algeria
P. tricaudata Booth, 1990
Subarctic Pacific
Polyfibula Manton & Bremer, 1981
P. sphyrrelata (Thomsen, 1973) Manton & Bremer, 1981
= *Pleurasiga sphyrrelata* Thomsen, 1973 (TM)
Tvärrminne, Isefjord
Polyoeca Kent, 1881
= *Acanthoecopsis* Norris, 1965 (p.p.)
P. dichotoma Kent, 1881 (TM)
= *P. dumosa* Dunkerly, 1910
= *Acanthoecopsis spiculifera* Norris, 1965 (TM) of *Acanthoecopsis* Norris, 1965
= *Salpingoeca tintinnabulum* Kent, 1881?
Denmark, S Bretagne, U.S. Pacific
Saepicula Leadbeater, 1980
S. pulchra Leadbeater, 1980 (TM)
Denmark, Bretagne
Saroeca Thomsen, 1979
S. attenuata Thomsen, 1979 (TM)
Tvärrminne, Öresund
Savilla Loeblich III, 1967
= *Ellisiella* Norris, 1965, non Gray, 1858 (Coelenterata)
S. parva (Ellis, 1930) Loeblich III, 1967
= *Diaphanoeca parva* Ellis, 1930 (TM)
= *Ellisiella parva* (Ellis, 1930) Norris, 1965
Tvärrminne, Isefjorden, Frederikshavn, Britain
S. micropora (Norris, 1965) Leadbeater, 1975
= *Pleurasiga micropora* Norris, 1965
Tvärrminne, Denmark, Parke & Leadbeater, 1977 area
Spiraloecion Marchant & Perrin, 1986 (TM) *S. didymostatum* Marchant & Perrin, 1986 - Antarctic
Stephanacantha Thomsen, in Thomsen & Boonruang, 1983
S. campaniformis (Leadbeater, 1973) Thomsen, in Thomsen & Boonruang, 1983
= *Parvicorbicula campaniformis* Leadbeater, 1973 (TM)
S Norway, Mediterranean
Stephanoecca Ellis, 1930
= *Sporetloeca* Norris, 1965
S. ampulla (Kent, 1880) Ellis, 1930
= *Salpingoeca ampulla* Kent, 1880 (TM)
Oslofjord, Engl. Channel
S. aphelis Thomsen, Buck & Chavez, 1991
Tvärrminne, Isefjord

- S. complexa* (Norris,1965) Thronsdene,1974
= *Pleurasiqa complexa* Norris,1965
Oslofjord
- S. cupula* (Leadbeater,1972) Thomsen,1988
= *Pleurasiqa cupula* Leadbeater,1972
Isfjord, Frederikshavn
- S. diplocostata* var. *diplocostata* Ellis,1930
= *S. pedicellata* Leadbeater,1972
Isfjord, Frederikshavn, Bergen, Britain, Mediterranean
- S. diplocostata* Ellis,1930 var. *paucicostata* Thronsdene,1969
Tvärminne, Isfjord, Oslofjord, Mandal
- S. elegans* (Norris,1965) Thronsdene,1974
= *Pleurasiqa elegans* Norris,1965
Isfjord, Bergen, Mediterranean, US Pacific
- S. kentii* Ellis,1930
Oslofjord
- S. urnula* Thomsen,1973
Tvärminne, Isfjord
- S. norrisii* Thomsen,1973
= *Sportelloeca kentii* : (Ellis,1930) Norris,1965, non *Stephanoea kentii* Ellis,1930
Isfjord
- S. campanula* (Kent,1880) Boucaud-Camou,1967
= *Salpingoeca campanula* Kent,1880
Engl. Channel
- S. constricta* Ellis,1930
Parke & Leadbeater,1977 area
- Syndetophyllum* Thomsen & Moestrup,1983 (TM *Parvicorbicula pulchella* Leadbeater,1974 - Indo-Pacific)
- CERCOMONADIDA Vickerman,1983, in Lee,1985
* With uncertain taxonomical position according to Patterson & Zöllfel,1992
- Cercomonadidae Kent = Cercobodonidae Hollande,1942
- Cercomonas* Dujardin,1841
?= *Dinastigamoeba* Blochmann,1894 ?= *Cercobodo* Krassilstchick,1886
?= *Prismatomonas* Massart,1920 ?= *Mukdeniamonas* Skvortzow,1960
?= *Changia* Skvortzow,1960 ?= *Reptomonas* Kent,1880
?= *Cercomasix* Lemmermann,1913 ?= *Helkesimastix* Woodcock & Lapage,1914
- C. crassicauda* (Alexeieff,1???)
= *Cercobodo crassicauda* (Alexeieff,1???) Lemmermann,1???
- Plymouth
- C. varians* Skuja,1???
- ?
Heteromita Dujardin,1841
- H. reniformis* (Zhukov, ?in Zhukov & Mylnikov,1983)
= *Bodomorpha reniformis* Zhukov, ?in Zhukov & Mylnikov,1983?
?
- Massisteria* Larsen & Patterson,1990
- M. marina* Larsen & Patterson,1990
Tvärminne, Limfjorden, England, Ireland, deep Atlantic, Brazil, Panama, Fiji, Hawaii, Australia
- PROTEROMONADIDA Grassé, in Grassé,1952 emend. Vickerman,19??
@ Parasitic in amphibians, anurans & urodeans
* Not marine
- RETORTAMONADIDA Grassé, in Grassé,1952
- Retortamonadidae Wenrich,1932
@ chiefly in vertebrates & insects, but *Chilomastix* is reported from e.g. *Box salpa* & *Toxopneustes esculentus*
- Chilomastix* Alexeieff,1912
- C. motellae* Alexeieff,1912
@ in alimentary canal of *Gaidropsaurus mediterraneus* & *Ciliata mustela*
Roscoff
- C. caulleryi* ?Alexeieff,1912? (TM)
@ ?
?
- C. undulata* Skuja,1956
* Free-living
?
- C. sp.* : Martin & Robertson,1911
@ in alimentary canal of *Pollachius virens*
E Scotland
- C. bocis* (Brumpt,1912)
@ in alimentary canal of *Box salpa*
S France
- Retortomonas* Grassi,1879
- Cochlosomidae Tyzzer,1930
@ intestinal bird parasites
* Not strictly marine
- TRICHOMONADIDA Kirby,1947
- Monoceromonadidae
@ In alimentary tracts of invertebrates & vertebrates
- Monoceromonadinae* Kirby,1947
- Monoceromonas* Grassi,1879
= *Eutrichomastix* Kofoed & Swezy,1915
- M. motellae* (Alexeieff,1910)
@ in rectum of *Gaidropsaurus mediterraneus*
Roscoff
- M. salpae* (Alexeieff,1910)
@ in rectum of *Box salpa*
Mediterranean
- M. molae* Noble & Noble,1966
- @ endocommensal in hindgut of *Mola mola*
California
- Protrichomonas** Alexeieff,1911
- P. legeri* (Alexeieff,1910) Alexeieff,1911
@ endocommensal in stomach of *Box boops*
Mediterranean
- Trimitus* Alexeieff,1910
= *Tricercomitus* Kirby,1931
- T. motellae* Alexeieff,1910
@ in rectum of *Gaidropsaurus mediterraneus*
Roscoff
- Chilomitinae**
* Not marine?
- Hypotrichomonadinae
@ in land tortoises & squamate reptiles
* Not marine
- Trichomonadidae** Wenyon,1926, em. Grassé,1952
@ In alimentary tracts of invertebrates & vertebrates
- Trichomonadinae** Chalmers & Pekkola,1917
- Trichomonas** Donné,1837
- T. prowazeki* Alexeieff,19??
@ *Box salpa*
Banyuls-sur-mer
- Tririchomonadinae**
* Not marine?
- Trichomitopsisinae**
* Not marine?
- Pentatrichomonoidinae**
* Not marine?
- Devescovinidae** Kirby,1931
@ In hindgut of termites
* Not marine
- Calonymphidae** Grassé,1911
@ In hindgut of termites
* Not marine
- OXYMONADIDA** Grassé, in Grassé,1952
- Polyamystigidae** Bütschli,1884
* Not marine?
- Pvsonymphidae** Grassé,1892
* Not marine?
- Oxymonadidae** Kirby,1928
@ Insect endosymbionts
* Not marine
- DIPLOMONADIDA**
- DIPLOMONADINA**
- Turbomonas* Prowse,1962
* Marine. Which family?
- Hexamitidae**
- Gyromonas* Seligo,1886
- Hexamita* Dujardin,1838
= *Hexamitus* Bütschli,1878 = *Urophagus* Klebs,1892
- H. inflata* Dujardin,1841
* a brackish water saprobionte
Marennes (France)
- H. nelsoni* Schlicht & Mackin,1968
@ held responsible for "pit disease" in *Ostrea edulis*, *Crassostrea* spp., *Ostrea lurida*
USA, Netherlands
- H. bovis* Lavier,1936
@ *Box boops*
Mediterranean
- H. motellae* Lavier,1936
@ *Gaidropsaurus mediterraneus*
Mediterranean
- H. salmonis* (Moore,1923) Kulda & Lom,1964
@ causes heavy epizootics in freshwater salmonid cultures
?
- Octomitus* von Prowazek,1904
* Correct family & order?
- O. motellae* Alexeieff,1910
@ in rectum of *Ciliata mustela* & *Gaidropsaurus mediterraneus*
Roscoff
- Trigonomonas** Klebs,1892
- Treptomonas** Dujardin,1841
- (RHIZOMASTIGIDA Doflein)
* Artificial order from which included taxa has gone to different other groups.
- HYPERMASTIGIDA** Grassé & Foa
- Joenidae* Janicki,1915
@ termite symbionts
* Not marine
- Lophomonadidae* Kent,1880
@ cockroach symbionts
* Not marine
- Rhizonymphidae* Grassé & Hollande,1951
@ termite symbionts

- * Not marine
Hoplonymphidae Light,1926
 @ termite & cockroach symbionts
 * Not marine
- Staurojoeninidae** Grassi,1917
 @ termite & cockroach symbionts
 * Not marine
- Trichonymphidae** Kent,1880
 @ insect symbionts
 * Not marine
- Eucomonymphidae** Cleveland, Hall, Sanders & Collier,1934
 @ termite symbionts
 * Not marine
- Teratonympidae** Koidzumi,1921
 @ termite symbionts
 * Not marine
- Spirotrichosomidae**
- Spirotrichonymphidae** Grassi,1917
 @ termite & cockroach symbionts
 * Not marine
- Holomastigotoididae** Janicki,1915
 @ termite symbionts
 * Not marine
- EBRIIDA** Poche,19??
 = **EBRIALES** Honigberg & al.,1964
 * Taxon incertus sedis, sensu Patterson & Zölfel,1992
- Ebriidae** De flandre,1936
 = **Ebriaceae** Lemmermann,1901
 * Monogenetic & monospecific family
- Ebria** Borgert,1891
 = *Plectanum* Haeckel,1881 (p.p.)
- E. tripartita** (Schumann,1867) Lemmermann,1900
 = *Dicyrocha tripartita* Schumann,1867
 = *Dicyroda formix* Möbius,1887 (?)
 = *Ebria formix* (Möbius,1887) Borgert,1891
 All Baltic, Öresund, Kattegatt, Limfjorden, cosmopolitan
- Hermesinaceae** = **Ebriopsidae** Deflandre,1950
- Hermesinum** Zacharias,1906 (TM*H. adriaticum* Zacharias,1906 - Adriatic)
- OPALINATA** Corliss & Balamuth,1963
 @ endocommensals in posterior portion of digestive tracts of chiefly amphibians
 * Only two species known from marine fishes
- Opalinidae** Claus,1874
- Opalininae** Claus,1874 (Metcalf,1920)
 @ parasitizing anuran frogs & toads
 * Not marine
- Protoopalininae** Metcalf,1920
- Protoopalina** Metcalf,1918
- P. saturnalis** (Léger & Duboscq,1904)
 = *Opalina saturnalis* Léger & Duboscq,1904
 @ Box hoops Linnaeus
 Mediterranean
- P. duboscqui** Lavier,1936
 @ *Gadus capelanus* Riso
 * This is probably a synonym of *P. saturnalis* (Léger & Duboscq,1904)
 Mediterranean (Banyuls-sur-Mer)
- SARCODINA** Schmarda,1871
- Podostoma** Claparède & Lachmann,1857
 * Marine? Which taxon?
- RHIZOPODEA** von Siebold,1845
- Gruberella** Page,1983
 * Not includable in any existing family of Gymnamoebia or Leptomyxida"
- G. flavescent** (Gruber,1889)
 Cornwall, Mediterranean
- Nolandella** Page,1983
 * Not includable in any existing family of Gymnamoebia or Leptomyxida"
- N. hibernica** (Page,1980)
 = *Hartmannella hibernica* Page,1980
 Ireland
- LOBOSEA** Carpenter,1861
- GYMNAMOEBA** Haeckel,1862
- Euhyperamoeba** Seravin & Goodkov,1987
 = *Hyperamoeba* Goodkov, Seravin & Railkin,1982, non Aléxieff,1924 (Flagellata)
 * Which taxon within Gymnamoebia?
- E. fallax** (Goodkov, Seravin & Railkin,1982)
 = *Hyperamoeba fallax* Goodkov, Seravin & Railkin,1982 (TM)
 * Marine.
 ?
- AMOEVIDA** Ehrenberg,1830, Kent,1880
- TUBULINA** Bovee & Jahn,1966
- Amoebidae** Diesing,1848
- Amoeba** Ehrenberg,1830
 = *Amiba* Bory de Siv. Vincent,1822
 * Marine species ascribed to this genus do not belong there
- "*A*". *lescherae* Taylor & Hayes,1944
- * in brackish tide pools. According to Page,1976 is this almost certainly a synonym of the freshwater species
Amoeba proteus Leidy,1878.
- "A."** *biddulphiae* Zuelzer,1927
 @ *Biddulpha sinensis* Grév.
 North Sea
- "A."** *communis* Rhumbler,1902
 Marine aquarium containing Helgoland material
- "A."** *crystalligera* Gruber,18??
 Baltic aquarium, North Sea water aquarium
- "A."** *flava* Gruber,18??
 Baltic aquarium
- "A."** *radiosa* Ehrenberg,18??
 Baltic
- "A."** *verrucosa* Ehrenberg,18??
 Baltic water aquarium
- "A."** *villosa* Wallich,18??
 Baltic aquarium
- "A."** *prehensilis* Möbius,18??
 Kieler Bucht
- "A."** *clavellinae* Huxley,1920
 @ *Clavelina lepadiformis*
 Mediterranean
- Trichamoeba** Fromentel,1874
- T. schaefferi** Radir,1927
 California
- Hydramoeba** Reynolds & Looper,1928
- Metachaos** Schaeffer,1926 (TM*M. discoides* ?,1???)
- Styamoeba** Sawyer,1975
- Entamoebidae** Chatton,1925
- Entamoeba** Cassigrandi & Barbagallo,1895
- E. gadii** Bullock,1966
 @ endocommensal in rectum of *Pollachius virens*
 Atlantic ocean
- E. molae** Noble & Noble,1966
 @ endocommensal in hindgut of *Mola mola*
 Pacific Ocean
- E. phallusiae** Mackinnon & Ray,1931
 @ *Phallusia mammillata*
 Plymouth
- Hartmannellidae** Volkonsky,1931 emend. Page,1974
- Hartmannella** Alexeieff,1912, emend. Page,1974
- H. abertawensis** Page,1980
 * in sandy beaches
 S Wales
- Janickina** Chatton, in Grassé,1953
- J. pigmentifera** (Grassi,1879)
 @ testis of *Spadella* spp. & *Sagitta* spp.
 Atlantic, Mediterranean
- J. chaetognathi** (Janicki,1912)
 @ testis of *Spadella* spp.
 Atlantic, Mediterranean
- Saccamoeba** Frenzel,1892, emend. Bovee,1972 & Page,1974
- Rhizamoeba** Page,1972
- R. saxonica** Page,1974
 North Sea
- R. globifera** (Gruber,1889)
 = *Amoeba globifera* Gruber,1889
 Mediterranean
- THECINA** Bovee & Jahn,1966
- Thecamoebidae** Schaeffer,1926
- Thecamoeba** Fromentel,1874
 = *Sriamoeba* Bovee & Jahn,1966 (invalid name)
- T. orbis** Schaeffer,1926
 Tärminne, North Sea, W North Atlantic, Gulf of Mexico
- T. hilla** Schaeffer,1926
 North Sea, W North Atlantic, Gulf of Mexico
- T. pulchra** (Biemacka,1963)
 Gdansk brackish water
- T. verrucosa** : Levander,1894, non ?,1???
- Gulf of Finland
 ? potentially worldwide
- Clydonella** Sawyer,1975 (TM*Rugipes vivax* Schaeffer,1926)
- Vannella** Bovee,1965
- V. devonica** Page,1979
 Devon (Channel coast)
- V. aberdonica** Page,1980
 Tärminne, Denmark, Aberdeen beach, Greenland, Belize

- V. caledonica* Page,1979
W Scotland
- V. platypodia* Gläser,1912
Tärminne, Denmark, European & N American fresh water
- V. septentrionalis* Page,1980
River Don (Aberdeen) estuary
- V. simplex* Wohlfarth-Bottermann,1960
Tärminne, Denmark, European & N American fresh water
- V. anglica* Page,1980
Suffolk brackish channel at high tide
- Platyamoeba* Page,1969
= *Lingulamoeba* Sawyer,1975
- P. plurinucleolus* Page,1974
W Britain, W Atlantic
- P. calycinucleolus* Page,1974
North Sea
- P. bursella* Page,1974
E coast of England
- P. flabellata* Page,1974
North Sea W North Atlantic
- FLABELLINA** Page,1976
- Hyalodiscidae** Poche,1913 emend. Page,1976
- Hyalodiscus** Hertwig & Lesser,1874 (TM)
* freshwater species
Baltic
- Gibbodiscus** Schaeffer,1926
- Unda** Schaeffer,1926
- Flabellulidae** Bovee,1970
- Flabellula** Schaeffer,1926
- F. trinovantica* Page,1980
E England
- F. demetica* Page,1980
Tärminne, S Wales, Arctic Canada
- F. citata* Schaeffer,1926 (TM)
?E Britain, W North Atlantic
- F. calcinsi* (Hogue,1914)
?E Britain, W North Atlantic
- Paraflabellula* Page & Willumsen, in Page, 1983
- P. reniformis* (Schmöller,1964)
= *Flabellula reniformis* : Auctt.
* "apparently very euryhaline"
Baltic
- Flamella* Schaeffer,1926 (TM) *F. magnifica* Schaeffer,1926
- CONOPODINA** Bovee & Jahn,1966
- Mayorellidae** Schaeffer,1926
- Mayorella* Schaeffer,1926
= *Hollandella* Page,1981
- M. gemmifera* Schaeffer,1926
Cornwall, Norfolk, W North Atlantic, Gulf of Mexico
- M. pussardi* Hollande, Nicolas & Escaig,1981
Mediterranean
- M.?* *alveolata* (Mereshowsky,1879)
= *Amoeba alveolata* Mereshowsky,1879
White Sea
- M.?* *mustela* (Delphy,1938)
= *Amoeba mustela* Delphy,1938
Biscay
- Dactylamoeba* Korotneff,1880
- Vexillifera* Schaeffer,1926
- V. minutissima* Bovee & Sawyer,1979
S Wales, W North Atlantic, Gulf of Mexico
- V. armata* Page,1979
S Devon
- V. telmatalassa* Bovee,1956
North Sea, W North Atlantic, Gulf of Mexico
- Pseudoparamoeba* Page,1979
- P. pagei* (Sawyer,1975)
S Devon, W North Atlantic, Gulf of Mexico
- Triaenamoeba* Bovee,1970
- Striolatus* Schaeffer,1926
- Dinamoeba*
- Boveella* Sawyer,1975
- Dactylosphaerium* Hertwig & Lesser,1874
- Mastigamoebidae**
- Mastigamoeba* Schulze,1875
- Mastigella* Frenzel,1897
- Paramoebidae** Poche,1913
- Paramoeba** Schaudinn,1896
- P. eilhardi* Schaudinn,1896
Kattegatt, Mediterranean, Indian Ocean
- P. aestuarina* Page,1970
North Atlantic, Gulf of Mexico
- P. pemaquidensis* Page,1970
* "one of the most common marine amoebae"
W Baltic, Britain, North Atlantic, Gulf of Mexico
- P. invadens* Jones,1985
@ causing mass mortalities of *Strongylocentrotus droebachiensis* along Atlantic coast of Nova Scotia
Nova Scotia
- P. perniciosa* Sprague, Beckett & Sawyer,1969
@ causing gray crab disease in *Callinectes sapidus*, *Cancer irroratus* & *Homarus americanus*
North Carolina
- ACANTHAPODINA** Page,1976
- Acanthamoebidae** Sawyer & Griffin,1975
- Acanthamoeba* Volkonsky,1931, emend. Page,1967
* pathogenic genus
- A. gigantea* Schmöller,1964
Baltic
- A. polyphaga* (Puschkarew,1???)
Britain
- Protacanthamoeba* Page,1981
- P. caledonica* Page,1981
Morar estuary
- Echinamoebidae** Page,1975
* No marine species known.
- SCHIZOPYRENIDA** Singh,1952
* This taxon should possibly be arranged among Labyrinthulids instead
* According to Patterson & Zoffel, in Patterson & Larsen,1992 is this taxon together with the slime mould family Acerasidae constituting a natural grouping Heterolobosea Page & Blanton,1985. However Heterolobosea is a taxon incertae sedis in relation to other taxa.
- Percolomonas* Fenchel & Patterson,1986
* Which family?
- P. cosmopolitus* (Ruijten,1938) Fenchel & Patterson,1986
= *Tetramitus cosmopolitus* Ruijten,1938 (TM)
= *Tetramitus ovoides* Ruijten,1938
Aarhus bay
- Vahlkampfiidae** Jollos,1917, Zulueta,1917
- Adelphamoeba* Napolitano, Wall & Ganz,1970
* Marine?
- Didascalus* Singh,1952
* Marine?
- Heteramoeba* Droop,1962
- H. clara* Droop,1962
Finland, Scotland
- Naegleria* Aléxieff,1912
* Marine?
- Paratetramitus* Derbyshire, Page & Goodfellow,1976
- P. jugosus* (Page,1967) Derbyshire, Page & Goodfellow,1976 (TM)
euryhaline freshwater & soil species from Europe & N America
- Tetramastigamoeba* Singh & Hanumaiah,1977 (TM) *Tetramastigamoeba hoarei* Singh & Hanumaiah,1977
* Marine?
- Tetramitus* Perty,1852 (TM) *Tetramitus rostratus* Perty,1852
= *Copromastix* Aragão,1909
- T. salinus* Entz,1883
* may grow in 20% S.
?
- Trimastigamoeba* Whitmore,1911
* Marine?
- Vahlkampfia* Chatton & Lalung-Bonnaire,1912
- V. baltica* Schmöller,1961
European Atlantic
- V. salina* (Hamburger,1905)
European Atlantic, Mediterranean
- V. longicauda* Schmöller,1964
Warnemünde
- V. dumnonica* Page,1983
Cornwall
- V. trilaminata* Schmöller, Jonas & Ludvik,1983
- V. discorbini* Le Calvez,1939
@ *Rosalina mediterraneensis* d'Orbigny
Banyuls-sur-mer
?
- V. paedophthora* ?Cauillery,1906
@ *Peltogaster curvatus* Kossmann
?
- V. mucicola* (Chatton,1910)
= *Amoeba mucicola* Chatton,1910
@ *Sympodus* spp. & *Trichodina labrorum*
?
- V. damariscottae* Page,19??

- ?Britain
- Willertia** de Jonckheere, Dive, Pussard & Vickermann,1984
=? *Protonegleria* Michel & Raether,1985
* Marine?
- PELIOBIONTIDA** Page,1976
- Pelomyxidae** Schultze,1877
- Pelomyxa** Greeff,1874 (TM *P. palustris* Greeff,1874)
* Marine species ascribed to this genus do not belong there
- "**P". ostendensis** (Kufferath,1952)
= *Amoeba ostendensis* Kufferath,1952
North Sea coastal waters
- "**P". marina** Delphy,1938
Arcachon
- TESTACEALOBOSIA** de Saedeleer,1934
- ARCELLINIDA** Kent,1880
- ARCELLINA** Haeckel,1894
- Cochliopodiidae** Taránek,1882
- Cochliopodium** Hertwig & Lesser,1874
- C. bilimbosum** (Auerbach,1856) Leidy,1879
= *Amoeba bilimbosa* Auerbach,1856
= *C. pellucidum* Hertwig & Lesser,1874 (TM - fresh water)
* Systematic classification sensu Page,1987; Bark,1973 & D Patterson suggest classification among Gymnamoebia.
Tärminne, Europeen & N American fresh water
- C. "sp. 2"** : Bark,1973
Kattegatt
- C.** sp. : Rhumbler,1923/25
Marine aquarium with Helgoland material
- Chlamydamoeba** Collin,1912 (TM *C. tentaculifera* Collin,1912 - marine)
- Gocevia** Valkanov,1932 (TM *G. pontica* Valkanov,1932 - marine)
- Microcoryciidae** de Saedeleer,1934
* fresh water genera only?
- Arcellidae** Ehrenberg,1832
- Arcella** Ehrenberg,1832 (TM)
* in estuarine sediments
Britain
- Hyalospheniidae** Schultze,1877 = *Cryptodifflugiidae* Rhumbler,1923
- Cryptodifflugia** Penard,1890 (TM *C. oviformis* Penard,1890)
= *Geococcus* Francé,1913, non Green,1902 (TM *G. vulgaris* Francé,1913)
- C. lanceolata** Golemansky,1970
* in beach sediments
North Sea, Black Sea
- Difflugiella** Cash,1904 (TM *D. apiculata* Cash,1904)
- D. collum** Chardez,1971
Belgian shores, W France
- D. lamingerae** Chardez,1977
Belgian shores
- D. psammophila** Golemansky,1970
North Sea, Black Sea
- Pseudowaillesella** Sudzuki,1979
- DIFFFLUGINA**
- Difflugiidae** Wallich,1864, Awerintzew,1906
- Difflugia** Leclerc, in Lamarck,1816 (TM *D. protoeiformis* Lamarck,1816)
- D. submarina** Chardez & Thomas,1980
W France
- D. subterranea** Golemansky,1970, var. *inflata* Decloitre,1975
French Mediterranean
- Centropyxiidae** Jung,1942, Deflandre,1953
- Centropyxis** von Stein,1859 (TM *Arcella aculeata* Ehrenberg,1832)
= *Echinopyxis* Claparède & Lachmann,1859
= *Millettella* Rhumbler,1904 (TM *Reophax pleurostomelloides* Millett,1899)
= *Armillentula* Rhumbler,1913 (TM *Reophax pleurostomelloides* Millett,1899)
* a fresh water genus, but may be found washed into shallow marine sediments
- Centropyxiella** Valkanov,1970
- C. arenaria** Valkanov,1970
North Sea shores
- C. gibbula** Valkanov,1970
North Sea shores
- C. gibbulina** Chardez & Thomas,1980
W France
- C. golemanskyi** Chardez,1977
Belgian shores
- C. oopyxiformis** Chardez,1977
Belgian shores
- C. platystoma** Golemansky,1981
- Baltic (Poland), Black Sea
- Micropsammella** Golemansky,1970
- M. minima** Chardez & Thomas,1980
W France
- M. retorta** Golemansky,1970
North Sea Shores
- Paraquadridiidae** Deflandre,1953
- Nebelidae** Taránek,1882 = *Hyalospheniidae* Schulze,1877
- Pomoriella** Golemansky,1970
- P. valkanovi** Golemansky,1970
Baltic, Black Sea
- Hyalosphaenia** von Stein,1857 (TM *H. cuneata* von Stein, in Schultze,1875 = *Difflugia ligata* Tatem,1870)
= *Catharia* Leidy,1874, non Lederer,1863 (TM *Difflugia ligata* Tatem,1870)
- H. savoiei** Chardez,1978
Belgian shores
- Lesquerella** Chardez & Thomas,1980
- L. mesopsammophila** Chardez & Thomas,1980
W France
- Plagiopyxidae** Bonnet & Thomas,1960
* No marin taxa?
- PHRYGANELLINA**
- Phryganellidae** Jung,1942
- Phryganella** Penard,1902 (TM *P. nidulus* Penard,1902)
- P. marinus** Chardez,1971
North Sea
- TRICHOSIDA** Möbius,1889
- Trichospaeridae** Möbius,1878
- Trichosphaerium** Schneider,1878
= *Pontifex* Schaeffer,1926
- T. sieboldi** Schneider,18??
Baltic, North Sea, Mediterranean
- T. micrum** Angell,1975
?
- T. platyxyrum** Angell,1976
?
- ACARPOMYXEA** Page,1976
- STEREOMYXIDA** Grell,1966
- Stereomyxidae** Grell,1966
- Stereomyxa** Grell,1966
- Corallomyxa** Grell,1966
- C. chattoni** Grell & Benwitz,1978
Cornwall, Mediterranean
- LEPTOMYXIDA** Pussard,1976
* Only freshwater & soil organisms?
- Leptomyxidae** Goodey,1915, Pussard & Pons,1976
- MYCETOZOA** de Bary,1859
* No marine species?
- PROTOSTELIA** L. Olive,1970
= **PROTOSTELIIDAE**
- PROTOSTELIIDA** Olive & Stoianovich,1966
= **PROTOSTELIALES**
- Cavosteliidae** L. Olive,1964
= **Cavosteliaceae**
- Protostellidae** Olive & Stoianovich,1966
= **Protosteliaceae**
- Ceratiomyxidae** Schröter,1889
= **Ceratiomyxaceae**
- Ceratiomyxella**
- C. tahitiensis** Olive & Stoianovich,19??
Tahiti
- DICTYOSTELIIDA** L. Olive,1970
= **DICTYOSTELIALES**
- Acytosteliidae** Raper, in Raper & Quinlan,1958
= **Acytosteliaceae**
- Dictyosteliidae** Rostafinski,1875
= **Dictyosteliaceae**
- GUTTULINIA** L. Olive,1970
= **GUTTULINIDAE**
- GUTTULINIDA** L. Olive,1970
= **GUTTULINALES**
- Guttulinopsidae** L. Olive,1970
= **Guttulinopsidaceae**
- Guttulinidae** Zopf,1884
= **Guttulinaceae**
- MYXOGASTRIA** L. Olive,1970

= **MYXOGASTRIDAE**

ECHINOSTELIIDA
= ECHINOSTELIALES

LICEIDA
= LICEALES

TRICHIDA
= TRICHIALES

STEMONITIDA
= STEMONITALES

PHYSARIDA
= PHYSARALES

PLASMODIOPHORIDA Cooke,1928

= PLASMODIOPHORALES

* Group of uncertain relationship.

Pyrrhosus Juel,1901

* very doubtfully a member of this order. May be a *Labyrinthula* or *Sorolpidium*

P. marinus Juel,1901

@ *Cystoclonium purpurascens* & ?*Ceranium rubrum*

Sweden

Plasmodiophoridae Zopf,1884

Plasmodiophora Woronin,1877

= *Ostenfeldiella* Ferdinandsen & Winge,1914

P. diplantherae (Ferdinandsen & Winge) Ivimey-Cook,1932

= *Ostenfeldiella diplantherae* Ferdinandsen & Winge,1914

@ *Halodule* spp.

West Indies

P. bicaudata J. Feldmann,1940

@ *Zostera nana* Roth.

France, W Africa

P. halophilae Ferdinandsen & Winge,1913

@ *Halophila* spp.

Java

P. maritima G. Feldmann,1958

@ *Triglochin maritima*

France

Phagomyxa Karling,1944

P. algarum Karling,1944

@ endosymbiont in several filamentous brown algae, e.g. *Bellerochea malleus* (Brightwell) Heurck USA

Tetramyxta Goebel,1884

= *Molliardia* Maire & Tison,1911

= *Thecaphora* Setchell,1924

T. parasitica Goebel,1884 (TM)

= *Thecaphora ruppiae* Setchell,1924

@ *Ruppia*, *Zannichellia* & *Potamogeton panormitanus*?

T. triglochinis Molliard,1909

= *Molliardia triglochinis* (Molliard) Maire & Tison,1911

@ *Triglochin palustris* & *Triglochin maritima*

* incompletely known. Provisional genus.

?

FILOSEA Leidy,1879

ACONCHULINIDA de Saedeleer,1934

Vampyrellidae Doflein,1902

* No marine species?

Reticulosidae Cash & Hopkinson,1905 = Penardiidae Loeblich & Tappan,1961

* No marine species?

GROMIIDA Claparéde & Lachmann,1859

GROMIINA Delage & Hérouard,1896

* With organic tests & filose pseudopodia.

Gromiidae Reuss,1862

Gromia Dujardin,1835

= *Arcellina* DuPlessis,1876, non Carter,1856 (TM) *A. marina* DuPlessis,1876

= *Hyalopus* Schaudinn,1894 (TM) *Gromia dujardini* M. Schultze,1854

G. oviformis Dujardin,1835 (TM)

= *Gromia dujardini* M. Schultze,1854

Bergen, Engl. Channel, Mediterranean, Cosmopolitan

Amoebogromia Giard,1900

A. cinnabarina Giard,1900 (TM)

@ associated with cirripeds

France

Bargoniella Loeblich & Tappan,1961

= *Salpicola* Bargoni,1894, non Richiardi,1880

B. amylacea (Bargoni,1894)

= *Salpicola amylacea* Bargoni,1894 (TM)

@ on *Salpa micronota* & *Salpa democratica*

Mediterranean

Lagunculina Rhumbler,1904

= *Ovulina* Gruber,1884, non Ehrenberg,1845

L. urnula (Gruber,1884)

= *Ovulina urnula* Gruber,1884 (TM)

Mediterranean

Urnulina Gruber,1884

U. diffugiaeformis Gruber,1884 (TM)

North Sea, Mediterranean

Ogdeniella (Golemansky,1970)

* Correct family?

O. carinata (Chardez,1977)

North Sea

O. elegans (Golemansky,1970)

North Sea, Black Sea

O. lucida (Golemansky,1970)

North Sea, Black Sea

O. pusilla (Chardez,1977)

Belgian shores

O. taschevi (Golemansky,1970)

North Sea, Black Sea

Chlamydophryidae de Saedeleer,1934

Lagenidiopsis Golemansky,1974

L. valkanovi Golemansky,1974

North Sea, Black Sea

Pseudolagenidiopsis Sudzuki,1979

* Correct family?

Pseudodifflugidae de Saedeleer,1934

Rhumbleriella Golemansky,1970

R. filosa Golemansky,1970

North Sea, Black Sea

Volutellidae Bovee, in Lee, Hutter & Bovee,1985

Volutella Sudzuki,1979

?= *Voluta* Chardez,1972, non Linnaeus,1758

Pseudovolutella Sudzuki,1979

* Correct family?

Psammonobiidae Golemansky,1974

* Encompassing sand dwelling species.

Psammonobiotus Golemansky,1967

P. balticus Golemansky,1973

Baltic, Black Sea

P. communis Golemansky,1967

North Sea, Black Sea

P. golemanskyi Chardez,1971

North Sea

P. linearis Golemansky,1970

North Sea, Black Sea

P. minutus Golemansky,1970

North Sea, Black Sea

P. plana Chardez,1971

North Sea

P. septentrionalis Chardez,1977

North Sea, Black Sea

Pseudocorythion Valkanov,1970

P. acutum (Wailes,1927)

North Sea, Black Sea

P. undulacollis Chardez & Thomas,1980

W France

P. walesi Golemansky,1971

North Sea, Black Sea

Messemvriella Golemansky,1973

Chardezia Golemansky,1970

Corythionella Golemansky,1970

C. acolla Golemansky,1970

North Sea, Black Sea

C. anteroplana Chardez,1977

Belgian shores

C. minima Golemansky,1970

North Sea, Black Sea

C. sudzukii Chardez,1977

North Sea

Corythionelloides Sudzuki,1979

Micramphora Valkanov,1970

M. amphoriformis Chardez & Thomas,1980

W France

M. atlantica Chardez & Thomas,1980

W France

M. pontica Valkanov,1970

North Sea, Black Sea

Ovulina Anderson, Rogerson & Hannah,1996

Firth of Clyde

O. parva Anderson, Rogerson & Hannah,1996 (TM - Firth of Clyde)

Firth of Clyde

EUGLYPHINA Bovee, in Lee, Hutter & Bovee,1985

- Euglyphidae Wallich,1864
* No marine species?
- Cyphoderidae de Saedeleer,1934
- Cyphoderia Schlumberger,1845
- C. ampulla** (Ehrenberg,1840)
= *Difflugia ampulla* Ehrenberg,1840
= *C. marginatacea* Schlumberger,1845 (TM)
* also in fresh water
Kieler Bucht
- C. truncata** Schulze,1875
* chiefly a fresh water species
Kiel, W  rtemb  ndt
- C. compressa** Golemansky,1979
North Sea, Black Sea
- C. littoralis** Golemansky,1973
North Sea, Black Sea
- Campascus** Leidy,1877 (TM *C. cornutus* Leidy,1877 - fresh water)
- C. interstitialis** Golemansky,1981
North Sea, Black Sea
- Pseudocyphoderia** Sudzuki,1979
- Paulinellidae de Saedeleer,1934
- Paulinella** Lauterborn,1895
- P. ovalis** (Wulff,1919) Johnson, Hargraves & Sieburth,1988
= *Calycomonas ovalis* Wulff,1919
Bothnian Sea, Tvr  minne, Gulf of Finland, N Baltic, Kieler Bucht, S Kattegatt, Oslofjord, S & N Norway, Barents Sea, Belgium, Massachusetts, New Zealand, Japan
- P. gracilis** (Lohmann,1908) V  rs,1992
= *Calycomonas gracilis* Lohmann,1908 (TM of *Calycomonas* Lohmann,1908)
S Norway, Park & Dixon,1976 area
- CRISTIDISCOIDIADA Page,1987
- Pompholyxophryidae Page,1987
- Rabdiophrys** Rainer,1968
= *Pinaciophora* Thomsen,1978, non Greeff,1875
- R. candelabrum** (Thomsen,1978) Roijackers & Siemersma,1988
Denmark (Dyb  s fjord), Tvr  minne
- R. multicosta** (Thomsen,1978) Roijackers & Siemersma,1988
= *Pinaciophora multicosta* Thomsen,1978
Denmark (Dyb  s fjord), Tvr  minne
- R. reticulata** (Thomsen,1979) Roijackers & Siemersma,1988
= *Pinaciophora reticulata* Thomsen,1979
Tvr  minne, S Baltic, USA, Arctic Canada
- R. thomseni** Roijackers & Siemersma,1988
Tvr  minne, S Baltic
- R. denticulata** (Thomsen,1978) Roijackers & Siemersma,1988
= *Pinaciophora denticulata* Thomsen,1978
Tvr  minne, Denmark (Dyb  s fjord), ?Peru
- R. paucipora** (Thomsen,1978) Roijackers & Siemersma,1988
= *Pinaciophora paucipora* Thomsen,1978
Denmark (Dyb  s fjord)
- R. spiculata** (Manton & Sutherland,1978) Roijackers & Siemersma,1988
= *Pinaciophora spiculata* Manton & Sutherland,1978
Tvr  minne, Denmark, North American Arctic Ocean
- R. "circulata"** Thomsen (nom. nudum) : V  rs,1992
Kattegatt
- Pompholyxophryx** Archer,1869 (TM ?*P. punicus* Archer,1869?)
- XENOPHYOPHOREA** Schulze,1904
- Aschemonellidae Eimer & Fickert,1899
* Which order?
- Aschemonellinae Eimer & Fickert,1899
- Aschemonella** Brady,1879 (TM *A. scabra* Brady,1879)
* The type species was moved from Foraminifera to Xenophyophorea by Gooday,1987
- A. catenata** Norman,18??
Sognefjord at 1200m according to Kjaer,1900
- PSAMMINIDA
- Psammettidae
- Psammetta** Schulze,1905 (TM *P. erythrocytomorpha* Schulze,1905 - Ind. Ocean)
- Psamminidae Haeckel,1889
- Psammina** Haeckel,1889 (TM *P. globigerina* Haeckel,1889)
= *Psammoplakina* Haeckel,1889 (TM *P. discoidea* Haeckel,1889 = *Psammina plakina* Haeckel,1889)
- Reticulamina** ?19??
- R. labyrinthica** (Tendal,1972)
off W Africa at 3000-4000 m.
- Holopssamma** Carter,1885 (TM *H. laevis* Carter,1885 - S Australia)
* Correct family?
- Psammopemna** Marshall,1881 (TM *P. densum* Marshall,1881 - trop. Atl. & Pacif.)
* Correct family?
- Syringamminidae
- Syringammina** Brady,1883
- S. fragillissima** Brady,1883 (TM)
Faeroe Channel
- Aschemonella** Brady,1879 (TM *A. scabra* Brady,1879 - N Pacific)
* Family tentative. The genus has long been thought to be a foraminiferan.
- A. ramuliformis** Brady,1884
?
- Cerelasimidae
- Cerelasma** Haeckel,1889 (TM *C. gyrophaera* Haeckel,1889 - Pacific)
- C. massa** Tendal,1972
off Biscay at >4000 m.
- STANNOMIDA
- Stannomidae Haeckel,1889
- Stannoma** Haeckel,1889 (TM *S. dendroides* Haeckel,1889 - Pacific)
= *Stannoplegma* Haeckel,1889 (TM *Stannoma coralloides* Haeckel,1889 - Pacific)
- Stannarium** Haeckel,1889 (TM *S. concretum* Haeckel,1889)
- Stannophyllum** Haeckel,1889 (TM *S. zonarium* Haeckel,1889 - Pacific)
= *Psammophyllum* Haeckel,1889 (TM *P. flustraceum* Haeckel,1889)
= *Neusina* Go  s,1892 (TM *N. agassizi* Go  s,1892)
- S. globigerinum** Haeckel,1889
central Atlantic at >5000 m.
- SCHIZOCLADEA** Cedhagen & Mattson,1992
- SCHIZOCLADIDA** Cedhagen & Mattson,1992
- Schizocladidae Cedhagen & Mattson,1992
- Schizocladus** Cedhagen & Mattson,1992
- S. sublittoralis** Cedhagen & Mattson,1992
Gullmarfjord
- GRANULORETICULOSEA** de Saedeleer,1934
- ATHALAMIDA** Haeckel,1862
- Biomixida** Loeblich & Tappan,1961
- Arachnula** Cienkowsky,1876
- A. impatiens** Cienkowsky,1876 (TM)
= *Biomixxa vagans* : M  bius,1888, non Leidy,1875
* also in fresh water
S Baltic, Black Sea
- Chlamydomyxidae
- Rhizoplasma** Verworn,1896
- R. kaiseri** Verworn,1896 (TM)
?
- Pontomyxa** Topsent,1892
- P. flava** Topsent,1892 (TM)
@ commensal in tunicates
?
- MONOTHALAMIDA Haeckel,1862
- Amphitrematidae** Poche,1913
* no marine species?
- Lieberkuhnidae de Saedeleer,1934
- Lieberkuhnia** Clapar  de & Lachmann,1859
- L. wagneri** Clapar  de & Lachmann,1859 (TM)
* fresh water & marine species
?
- Microgromiidae de Saedeleer,1934
* no marine species?
- FORAMINIFERA** d'Orbigny,1826, Eichwald,1830
- ALLOGROMIINA** Loeblich & Tappan,1961
- Lagynacea
- Lagynidae Schultze,1854
- Boderia** T.S. Wright,1867
- B. turneri** T.S. Wright,1867 (TM)
@ *Corallina*
North Sea, Plymouth
- Kibisidyles** Jepps,1934
- K. marinus** Jepps,1934 (TM)
* in surface film or attached to floating objects
Europe, e.g. Scotland (Millport)
- Lagynis** Schultze,1854
= *Platoum* Schultze,1875
- L. baltica** Schultze,1854 (TM)
= *Platoum parvum* Schultze,1875
S Baltic
- Myxotheca** Schaudinn,1893
= *Armyxothecum* Rhumbler,1913
- M. arenilega** Schaudinn,1893 (TM)
Bergen
- Ophiotuba** Rhumbler,1894
- O. gelatinosa** Rhumbler,1894 (TM)
* in *Saccammina*-shells
North Sea, Bergen, Kristiansand
- Pilalla** Rhumbler,1935

- P. exigua* Rhumbler,1935 (TM)
Kieler Bucht, Helgoland
- Rhumblerinella* Schmidt,1929
- R. bacillifera* Schmidt,1929
North Sea
- Schultzella* Rhumbler,1904
= *Schulzia* Gruber,1888, non Grimm,1876, nec Graff,1882
= *Arsctzellum* Rhumbler,1913
- S. diffluens* (Gruber,1884)
= *Lieberkuehnia diffluens* Gruber,1884 (TM)
Europe
- Marenda* Nyholm,1951
- M. nematoidea* Nyholm,1951 (TM)
W France
- Allogromiidae Rhumbler,1904
- Allogromiinae Rhumbler,1904
- Allogromia* Rhumbler,1904
= *Craterina* Gruber,1884, non Curtis,1826, nec Bory de St. Vincent,1827
= *Arrugromonium* Rhumbler,1913
- A. mollis* (Gruber,1884)
= *Craterina mollis* Gruber,1884 (TM)
Bergen
- A. ovoidea* Rhumbler,19??
Britain
- A. crystallifera* Dahlgren,1962
Gullmarsfjord at 80-120 m.
- A. sp. A* : Hedley,Ogden & Wakefield,1972
* on *Corallina*
Plymouth
- A. sp. B* : Hedley,Ogden & Wakefield,1972
* on *Corallina*
Plymouth
- Cylindrogullmia* Nyholm,1974
- C. alba* Nyholm,1974 (TM)
Gullmar Fjord
- Gloiogullmia* Nyholm,1974
- G. eurystoma* Nyholm,1974 (TM)
Gullmar Fjord
- Lieberkuehnia* Claparède & Lachmann,1859
- L. wagneri* Claparède & Lachmann,1859 (TM)
* Marine?
Europe
- Marsupulina* Rhumbler,1904
- M. schultzei* Rhumbler,1904 (TM)
North Sea, Mediterranean
- Pleurophysa* Claparède & Lachmann,1859
- P. sphaerica* Claparède & Lachmann,1859 (TM)
* Marine?
Germany
- P. lageniformis* Schulze,18??
S Baltic
- P. blanckenbergi* ??,1???
- Belgium
- Rhynchogromia* Rhumbler,1894
- R. variabilis* Rhumbler,1894 (TM)
* in Saccammina-shells
North Sea, Bergen, Kristiansand
- Rhynchosaccus* Rhumbler,1894
- R. immigrans* Rhumbler,1894 (TM)
* in Saccammina-shells
North Sea, Bergen, Kristiansand
- Sphairogullmia* Nyholm,1974
- S. aurea* Nyholm,1974 (TM)
Gullmar Fjord
- Shepheardellinae Loeblich & Tappan,1984
- Nemogullmia* Nyholm,1953
- N. longevariabilis* Nyholm,1953 (TM)
Gullmar Fjord
- Phainogullmia* Nyholm,1955
- P. aurata* Nyholm,1955 (TM)
Gullmar Fjord
- Shepheardella* Siddall,1880
- S. taeniformis* Siddall,1880 (TM)
Bergen, England
- Tinogullmia* Nyholm,1954
- T. hyalina* Nyholm,1954 (TM)
Gullmar Fjord
- Argillotubinae Avnimelech,1952
- Dactylosaccus* Rhumbler,1894
- D. vermiciformis* Rhumbler,1894 (TM)
* in Saccammina-shells
North Sea, Bergen, Kristiansand
- Dendrotuba* Rhumbler,1894
- D. nodulosa* Rhumbler,1894 (TM)
* in Saccammina-shells
Bergen, Kristiansand, N Atlantic
- Micatuba* Avnimelech,1952
?= *Rhabdogromia* Valkanova,1964
- M. flexilis* (Höglund,1947)
= *Bathyiphon flexilis* Höglund,1947 (TM)
Sweden 7-109m, Oslofjord, Bulgaria 0-12 m
- Micrometula* Nyholm,1952
- M. hyalostriata* Nyholm,1952 (TM)
Gullmar Fjord at 30-118 m.
- Nodellum* Rhumbler,1913
= *Chitinosiphon* Thalmann & Bermúdez,1954
- N. membranacea* (Brady,1879)
= *Reophax membranacea* Brady,1879 (TM)
= *Chitinosiphon rufescens* Thalmann & Bermúdez,1954
Atlantic, deep water
- Hospitellidae* Loeblich & Tappan,1984
- Hospiella* Rhumbler,1911
- H. fulva* Rhumbler,1911 (TM)
Atlantic
- Placopsilinella* Earland,1934
- P. aurantiaca* Earland,1934 (TM)
Atlantic
- Thalamophaga* Rhumbler,1911
= *Orbitophaga* Rhumbler,1911 = *Marsupophaga* Rhumbler,1911
= *Tubophaga* Rhumbler,1911 = *Nummophaga* Rhumbler,1911
- T. ramosa* Rhumbler,1911 (TM)
Atlantic
- Phtanotrichidae* Arnold,1974
* No species in the area?
- TEXTULARIINA Delage & Hérouard,1896
- Astrorhizaceae Sandahl,1858
- Astrorhizidae Sandahl,1858
- Astrorhizinae Sandahl,1858
- Astrorhiza* Sandahl,1858
= *Arenistella* Fischer, in de Folin & Perier,1875 (TM) *A. agglutinans* Fischer,1875
= *Astrodiscus* Schultze,1875, non Ludwig,1866 (TM) *A. arenaceus* Schultze,1875
= *Haekelia* Bessels,1875
- A. limicola* Sandahl,1858 (TM)
= *Amoeba gigantea* Sandahl,1858
= *Megamoebomyxa argillobia* Nyholm,1950
= *Astrorhiza arenifera* Stschedrina,1946
= *Astrorhiza sabulifera* Stschedrina,1946
= *Astrorhiza arctica* Stschedrina,1958
= *Haekelia gigantea* Bessels,1875 (TM of *Haekelia* Bessels,1875)
- Kattegatt-Bohusl., Bergen, Barents Sea
- A. arenaria* Norman,1876
Skagerrak at 230-510 m., Oslofjord at 78-200 m., Boknafjord, Bergen-Hammerfest, North Sea, Shetlands, off Faeroes, Spitsbergen, Barents Sea, Kara Sea, Iceland, E Greenland, off SW Ireland, off NW Africa
- A. angulosa* Brady,1881
* Correct genus?
off Britain at 630 fathoms
- Astrorhizoides* Shchedrina,1969
- A. cornuta* (Brady,1879)
= *Astrorhiza cornuta* Brady,1879 (TM)
Faeroe Channel, Rockall Bank, Arctic Seas
- Pelosina* Brady,1879
= *Pelosinella* Parr,1950 (TM) *P. bicaudata* Parr,1950
= *Globosiphon* Avnimelech,1952 (TM) *Pelosina variabilis* Brady,1879 var. *sphaeriloculum* Höglund,1947
?= *Millettina* Avnimelech,1952 (TM) *Pelosina distoma* Millett,1904
- P. arborescens* Pearcey,1914
Kattegatt, Gullmarfjord, Koster Channel, Oslofjord, W Norway, W Scotland, Faeroe Channel
- P. fusiformis* Earland,1933
Gullmarfjord at 35-55 m., ?Oslofjord
- P. variabilis* Brady,1879 (TM)
Skagerrak at 230-500 m., Oslofjord, W Norway, Franz Josef's Land, W Scotland, Biscay, Arctic Ocean, cosmopolitan
- P. variabilis* Brady,1879 var. *sphaeriloculum* Höglund,1947
border of Skagerrak & North Sea at 500-510 m.
- P. rotundata* Brady,1879
off S Rockall Bank at 640 fathoms
- Radicula* Christiansen,1958
- R. limosa* Christiansen,1958 (TM)
Drøbak
- Saccodendron* Rhumbler,1935
- S. heronalleni* Rhumbler,1935 (TM)
Kieler Bucht, Skagerrak at 204-700 m., Oslofjord
- S. heronalleni* Rhumbler,1935, forma *latericum* Rhumbler,19?? : Gabel,1971
North Sea

- Vanhoeffenellinae Saidova,1981
- Ampithenestrella** Rhumbler,1935
- A. wiesneri** Rhumbler,1935 (TM)
* Based on a single specimen. Not refound.
off Germany
- Causia** Rhumbler,1938
- C. injusticata** Rhumbler,1938 (TM)
German North Sea coast
- Vanhoeffenella** Rhumbler,1905
- V. gaussi** Rhumbler,1905 (TM)
Kieler Bucht at 40 m., Drøbak, Iceland at 1889-1996 m., Cornwall
- Bathysiphonidae Avnimelech,1952
- Astrorhizinilla** Saidova,1975 (TM) *Psammosiphonella aetherea* Saidova,1970
* genus cosmopolitan at 1260-2000 m.
- Bathysiphon** M. Sars, in G.O. Sars,1872
= *Psammosiphonella* Avnimelech,1952 (TM) *Bathysiphon arenaceus* Cushman,1927
= *Bogdanowiczia* Pishvanova & Vyavol, in Vyavol & Pishvanova,1967 (TM) *B. pocutiva* Pishvanova & Vyavol, in Vyavol & Pishvanova,1967
= *Flagrina* Vyavol,1968 (TM) *F. staminea* Vyavol,1968
- B. argentea** Heron-Alen & Earland,1913
Gullmarfjord at 33-118 m., Skagerrak, Oslofjord, W Norway, N North Sea, Scotland, W Ireland
- B. filiformis** M. Sars, in G.O. Sars,1872 (TM)
Drøbak, Hardangerfjord, Bergen, Sognefjord, off SE Ireland, Biscay, Mediterranean
- B. minutus** Pearcey,1900
Kattegatt at 32 m., Gullmarfjord at 7-109 m., Skagerrak, Oslofjord, W Scotland
- B. crassatina** (Brady,1881)
= *Astrotricha crassatina* Brady,1881
= *Rhabdammina crassatina* : Eimer & Fickert,1899
* Correct genus?
W Norway, Faeroe Channel, Spitsbergen, Arctic Ocean
- Rhabdamminella** de Folin,1887
= *Pseudomarsipella* Saidova,1975 (TM) *Marsipella cylindrica* Brady,1882
- R. cylindrica** (Brady, in Tizard & Murray,1882)
= *Marsipella cylindrica* Brady, in Tizard & Murray,1882
= *Rhabdamminella prismaeignosa* de Folin,1887 (TM)
Drøbak, NW North Sea, Faeroe Channel
- Rhabdamminidae** Brady,1884
- Rhabdammininae** Brady,1884
- Marsipella** Norman,1878
* Genus cosmopolitan at 108-3216 m
- M. elongata** Norman,1878 (TM)
Faeroe Channel, off W Ireland, Wales, Rockall Bank
- M. arenaria** Christiansen,1958
Drøbak
- M. spiralis** Heron-Alen & Earland,1912
Gullmarfjord at 60-70 m., Koster Channel at 200 m., Skagerrak at 196-626 m., NW North Sea at 330-390 m.
- Oculosiphon** Avnimelech,1952 (TM) *Rhabdammina linearis* Brady,1879
- O. linearis** (Brady,1879)
Hardangerfjord, Bergen, W Ireland
- Rhabdammina** M. Sars, in Carpenter, 1869
- R. abyssorum** M. Sars, in Carpenter,1869 (TM)
Skagerrak at 500-700 m., Norway, Barents Sea, Arctic Ocean, cosmopolitan at 700-4800m
- R. discreta** Brady,1881
Koster Channel, Skagerrak, Oslofjord, Bergen, Greenland
- R. inaequalis** le Calvez,1935
= *Rhabdammina linearis* : Auctt., non Brady,1879
Koster Channel, Oslofjord, Skagerrak at 200-500 m., Mediterranean
- R. scabra** Höglund,1947
Oslofjord, Skagerrak at 200-500 m
- R. sp. I** : Gabel,1971
Fladen Ground
- R. sp. II** : Gabel,1971
Dogger Bank
- Rhizammina** Brady,1879
= *Testulosiphon* Avnimelech,1952 (TM) *Rhizammina indivisa* Brady,1884
- R. indivisa** Brady,1884
North Sea, Faeroe Channel
- R. algaeformis** Brady,1879 (TM)
Bergen, off W Scotland, off W Ireland, N & S Atlantic at 1260-5800m
- Dendrophryinae** Haeckel,1894
- Dendrophrya** T.S. Wright,1861
- D. erecta** T.S. Wright,1861 (TM)
Drøbak, off Scotland
- D. radiata** T.S. Wright,1861
Kieler Bucht, Scotland, Wales
- Psammatodendron** Norman, in Brady,1881
= *Dendrophryna* Vyavol,1966 (TM) *Psammatodendron dichotomicum* Neagu,1964
- P. arborescens** Norman, in Brady,1881 (TM)
@ tubes of *Haplops*
Öresund, W Norway, Britain, Iceland, Greenland, Novaja Zemlja, Franz Josefs Land, Arctic Ocean
- Halophyseminae** Loeblich & Tappan,1984
- Halophysema** Bowerbank,1862
- H. tumanowiczii** Bowerbank,1862 (TM)
= *Squamulina scopula* Carter,1870 (TM) of *Gastrophysema* Haeckel,1877
* According to Kiao,1960, a variety of *H. abyssorum* Goes exists at 540m in the North Sea, although rather uncommon.
Bergen, Wales, Ireland, Engl. Channel
- H. ramulosum** Bowerbank,1866
Kiel Canal, Engl. Channel, Ireland
- Nubeculariella** Averintsev,1911
- N. birulai** Averintsev,1911 (TM)
Arctic Ocean
- Hippocrepinellidae** Loeblich & Tappan,1984
- Hippocrepinella** Heron-Allen & Earland,1932
= *Hirudina* Vyavol,1966 (TM) *Hippocrepinella hirudiformis* Suzin, in Bogdanovich,1963
- H. alba** Heron-Allen & Earland,1932
Kieler Bucht, Kattegatt, Gullmarfjord, Koster Channel, Skagerrak, Oslofjord, Bergen
- H. ampulacea** Rhumbler,1935
Kieler Bucht, Drøbak
- H. hirudinea** Heron-Allen & Earland,1932 (TM)
Kieler Bucht, Kattegatt, Gullmarfjord, Koster Channel, Skagerrak, Oslofjord, Shetlands, off SW Ireland, Spitsbergen
- H. hirudinea** var. *crassa* Heron-Allen & Earland,1932
?Koster Channel at 200 m., ?Oslofjord, Antarctic
- Schizaminidae** Norvang,1961
- Schizammina** Heron-Allen & Earland,1929 (TM) *S. labyrinthica* Heron-Allen & Earland,1929 - off French Equatorial Africa
- Psammosphaeridae** Haeckel,1894
- Psammosphaerinae** Haeckel,1894
- Psammosphaera** Schulze,1875
= *Psammelia* Rhumbler,1935, non Lendenfeld,1887 (TM) *Psamella frankei* Rhumbler,1935
= *Arenosphaera* Schchedrina,1939 (TM) *Arenosphaera perforata* Schchedrina,1939
- P. fusca** Schulze,1875 (TM)
Kieler Bucht, Gullmarfjord, Oslofjord, Haugesund, Barents Sea, Spitsbergen, North Sea, Britain
- P. parva** Flint,1899
= *Psammosphaera fusca* Schulze,1875, var. *parva* Flint,1899
?Helgoland, Faeroe Channel, Hebrides, Wales, Brazil
- P. bowmani** Heron-Alen & Earland,1912
Gullmarfjord, Oslofjord at 10-60 m., NE & W Scotland, Ireland, Engl. Channel, Spitsbergen
- P. testacea** Flint,1899
= *Psammosphaera fusca* Schulze,1875, var. *testacea* Flint,1899
Drøbak, Britain, Gulf of Mexico
- P. rustica** Heron-Alen & Earland,1912
Scottish North Sea, W Scotland
- P. perforata** (Schchedrina,1939)
= *Arenosphaera perforata* Schchedrina,1939
Barents Sea
- Sorospaera** Brady,1879
- S. confusa** Brady,1879 (TM)
Drøbak, Faeroe Channel, Rockall Bank, Azores, cosmopolitan
- Spiculosiphon** Christiansen,1964
- S. radiata** Christiansen,1964 (TM)
N Norway at 100m.
- Storthosphaera** Schulze,1875
- S. albida** Schulze,1875 (TM)
Skagerrak at 626-700 m., Bergen, Bukkenfjord, Faeroe Channel, Biscay
- Saccamminidae** Brady,1884
- Saccammininae** Brady,1884
- Lagenammina** Rhumbler,1911
= *Proteonella* Lukina,1969 (TM) *Reophax diffugiformis* Brady,1879
- L. laguncula** Rhumbler,1911 (TM)
N Atlantic at 1524-2400 m.
- L. diffugiformis** (Brady,1879)
= *Reophax diffugiformis* Brady,1879
Bergen, Sognefjord, Lofoten, Barents Sea, Arctic Ocean
- L. arenulata** (Skinner,1961)
= *Reophax diffugiformis* Brady,1879 subsp. *arenulata* Skinner,1961
= *Proteonina diffugiformis* : Höglund,1947
Skagerrak at 215-220 m., Faeroe Channel, off Greenland, Novaja Zemlja, Wales
- L. cf. hancocki** (Cushman & McCulloch,1948)
= *Proteonina compressa* Cushman & McCulloch,1939, non ?,1???
- O. opaca** Dahlgren,1962 (TM)
Gullmar Fjord to 1 m.
- Saccammina** Carpenter,1869
= *Saccammina* M. Sars,1869 (nom. nud.)
- S. sphaericola** Brady,1871 (TM)
= *S. sphaericola* M. Sars,1869 (nom. nud.)
= *S. sphaericola* G.O. Sars,1871
Koster Channel, Skagerrak at 200-700 m., Oslofjord, Bergen, Faeroe Channel, around Britain, Biscay, Greenland, Franz Josefs Land, Arctic Ocean
- S. tubulata** Rhumbler, in Wiesner,1931
= *Proteonina* cf. *tubulata* : Höglund,1947

- ?Skagerrak at 215-515 m., Oslofjord
- S. socialis** Brady,1884
= *S. consociata* Flint,1899
Faeroe Channel, off S Rockall Bank
- S. minuta** Rhumbler,1911
off Hebrides at 2275 m.
- Technitella** Norman,1878
- T. legumen** Norman,1878 (TM)
Gullmarfjord, Koster Channel, Säcken, Skagerrak, Oslofjord, Bergen, Bukken area, Iceland, Scottish North Sea, Ireland
- T. melo** Norman,1878
off S Rockall Bank at 1215 fathoms, Biscay
- T. thompsoni** Heron-Alen & Earland,1909
NE Scotland, off E England
- T. teivyense** Haynes,1973
Wales
- Pilulininae** Brady,1884
- Pilulina** Carpenter,1870
- P. argentea** Höglund,1947
Skagerrak at 500-700 m.
- P. jeffreysii** Carpenter,1875 (TM)
off S Rockall Bank at 630-1476 fathoms.
- Thurammininae** A.D. Miklukho-Maklay,1963
- Astrammina** Rhumbler, in Wiesner,1931
- A. rara** Rhumbler, in Wiesner,1931 (TM)
= *Armarella sphaerica* Heron-Alen & Earland,1932 (TM of *Armarella* Heron-Alen & Earland,1932)
= *Pelosphaera cornuta* Heron-Alen & Earland,1932 (TM of *Pelosphaera* Heron-Alen & Earland,1932)
Kieler Bucht, Kattegatt, Gullmarfjord, Koster Channel, Skagerrak, Oslofjord, Spitsbergen
- Pseudothurammina** D.B. Scott, Medioli & M.A. Williamson,1981 (TM *Thurammina? limnetis* D.B. Scott & Medioli,1980)
- P. sp.**
Kieler Bucht
- Thurammina** Brady,1879
- T. papillata** Brady,1879 (TM)
off Lofoten, North Sea, around Britain, Faeroe Channel
- T. faerleensis** Höglund,19??
= *T. (?) sphaerica* Höglund,1947, non ?,1???
Gullmarfjord, Uddevalla
- T. compressa** Brady,1879
off S Rockall Bank at 630 fathoms
- Hemisphaeraminidae** Loeblich & Tappan,1961
- Hemisphaeramininae** Loeblich & Tappan,1961
- Ammopemphix** Loeblich,1952 (TM *Urnula quadrupla* Wiesner,1931)
= *Urnula* Wiesner,1931, non Claparède & Lachmann,1857
- A. arctica** (Cushman,1933)
= *Urnula arctica* Cushman,1933
NE Greenland
- Hemisphaeramina** Loeblich & Tappan,1957 (TM *H. batalleri* Loeblich & Tappan,1957)
= *Fairfieldia* Summers,1958 (TM *F. dicantha* Summers,1958)
= *Iridella* Shchedrina,1962 (TM *I. marisalbi* Shchedrina,1962)
= *Metamorphaea* Browne, in Browne & Schott,1963 (TM *Webbinella tholus* Moreman,1933)
- H. bradyi** Loeblich & Tappan,1957
= *Webbinia hemispherica* Brady,1884, non Parker, Jones & Brady,1866
* There are doubts whether this is a protozoan and not an egg capsule of a gastropod.
North Sea, Wales
- Iridia** Heron-Alen & Earland,1914
- I. diaphana** Heron-Alen & Earland,1914 (TM)
Plymouth on sandy sediment at 5 fathoms, W Scotland, SE Africa
- Tholosina** Rhumbler,1895
- T. bulla** (Brady,1881)
= *Placopsisina bulla* Brady,1881 (TM)
Skagerrak, Drøbak, Bergen, Lofoten, off NW Ireland, Barents Sea
- T. vesicularis** (Brady,1879)
= *Placopsisina vesicularis* Brady,1879
* On red algae & stones
Kieler Bucht, Belt Sea, Drøbak, Shetland-Faeroe Channel, between NW Ireland & Rockall Bank
- T. sp. : Nagy,1965**
Spitsbergen
- Crithionininae** Hofker,1972
- Crithionina** Goës,1894 (TM *C. mamilla* Goës,1894)
- C. granum** Goës,1894
Gullmarfjord, Singlefjord, Skagerrak, Oslofjord, Bergen
- C. mammilla** Goës,1894
Gullmarfjord, Koster Channel, Skagerrak, Oslofjord, Bergen, North Sea, Faeroe Channel, W Scotland, W Ireland
- C. pisum** Goës,1896
= *C. abyssorum* Kiær,1899
Skagerrak at >200 m, Oslofjord, Bergen, Stadt, Lofoten
- C. heincke** Rhumbler,1928
* On red algae & stones
W Baltic
- C. pisum** Goës,1896 var. **hispida** Flint,1899
= *C. abyssorum* Kiær,1899 (p.p.)
Skagerrak at 200-700 m., Oslofjord, Bergen, Faeroe Channel
- Pseudowebbinella** Shchedrina,1962
- P. goesi** (Höglund,1947)
= *Crithionina goesi* Höglund,1947 (TM)
= *Placopsisina bulla* : Goës,1894, non Brady,18??
Bohusl, Koster Channel, Skagerrak, Oslofjord, North Sea, Norway, White Sea, Arctic Ocean
- Komokioacea** Tendal & Hessler,1977
- Komokidida** Tendal & Hessler,1977
- Ipoa** Tendal & Hessler,1977 (TM *Ipoa fragila* Tendal & Hessler,1977)
- I. sp. : Tendal & Hessler,1977**
off Biscay in deep water
- Lana** Tendal & Hessler,1977 (TM *Lana neglecta* Tendal & Hessler,1977)
- L. sp. : Tendal & Hessler,1977**
Skagerrak at 400 m, off Bergen at 670 m.
- Normanina** Cushman,1928
- N. conferta** (Norman,1878)
= *Haliphysema confertum* Norman,1878 (TM)
* The Oslofjord specimen may be another species according to Tendal & Hessler,1977
?Oslofjord at 202 m., Davis Strait
- N. sp. : Tendal & Hessler,1977**
off Biscay in deep water
- Septuma** Tendal & Hessler,1977 (TM *Septuma ocoillo* Tendal & Hessler,1977)
- S. sp. : Tendal & Hessler,1977**
off Biscay in deep water
- Baculellidae** Tendal & Hessler,1977
- Baculella** Tendal & Hessler,1977 (TM *Baculella globofera* Tendal & Hessler,1977)
- B. sp. : Tendal & Hessler,1977**
off Biscay in deep water
- Edgertonia** Tendal & Hessler,1977 (TM *Edgertonia tolerans* Tendal & Hessler,1977)
- E. sp. : Tendal & Hessler,1977**
off Biscay in deep water
- Globipelorhiza** Cedhagen & Mattson,1991
- G. sublittoralis** Cedhagen & Mattson,1991 (TM)
Koster Channel, Bergen, N Norway
- Hippocrepinaceae** Rhumbler,1895
- Hippocrepinidae** Rhumbler,1895
- Hyperammininae** Eimer & Fickert,1899
- Botellina** Carpenter, Jeffreys & Thomson,1870
- B. labyrinthica** Brady,1881 (TM)
Koster, Bergen, Hammerfest, Faeroe Channel, NE & W Scotland, N Ireland, N Atlantic at 835-913 m.
- Hyperammina** Brady,1878
= *Bactrammina* Eimer & Fickert,1899 (TM *H. elongata* Brady,1878)
* What is *Hyperammina arboreascens* Norman, cited by Kiær,1900 from Bergen (Novaja Zemlya - S Africa, 40-700m)?
- H. elongata** Brady,1878 (TM)
Koster Channel, Skagerrak at 200-507 m., Oslofjord, Bergen, Faeroes, NE Greenland, Novaja Zemlya, Franz Joseph Land
- H. laevigata** Wright,1891
= *H. elongata* Brady,1878 var. *laevigata* Wright,1891
Gullmarfjord, Koster Channel, Skagerrak, Oslofjord, Bergen, off Ireland, Barents Sea
- H. friabilis** Brady,1884
Skagerrak at >200 m., Oslofjord, Bergen, around Britain
- H. fragilis** Höglund,1947
Gullmarfjord, Koster Channel, Skagerrak, Oslofjord, Greenland
- H. bradyi** Stschedrina,19??
Barents Sea
- H. sp. : Nørvang,1945**
Iceland
- Saccorhiza** Eimer & Fickert,1899
?= *Pseudoschizammina* Saidova,1975 (TM *P. edita* Saidova,1975)
- S. ramosa** (Brady,1879)
= *Hyperammina ramosa* Brady,1879 (TM)
Hardangerfjord, Bergen, Sognefjord, Stadt, Iceland, Barents Sea, Franz Josef Land, Arctic Ocean
- Hippocrepinidae** Rhumbler,1895
- Hippocrepina** Parker, in G.M. Dawson,1870
- H. acuta** (Höglund,1947)
= *Hippocrepinella acuta* Höglund,1947
Gullmarfjord
- H. cylindrica** Höglund,1947
Skagerrak at 500-510 m., Oslofjord
- H. indivisa** Parker, in G.M. Dawson,1870 (TM)
Norway, NE Scotland, W Ireland, W Greenland, Barents Sea, Novaja Zemlya, NW Atlantic, Arctic Ocean
- H. pusilla** Heron-Alen & Earland,1930
Kattegatt, Gullmarfjord, Koster Channel, Oslofjord, North Sea, Plymouth
- H. flexibilis** (Wiesner,1931)
= *Technitella flexibilis* Wiesner,1931
= *Hippocrepinella remanei* Rhumbler,1935
Stolpe Rinne, Kieler Bucht, Belt Sea
- Jaculella** Brady,1879
- J. acuta** Brady,1879 (TM)
off W Norway, Shetlands

- J. obtusa** Brady,1882
Koster Channel, Drøbak, Bergen, Faeroe Channel, Iceland, off W Scotland, W Ireland, Siberian Arctic
- Ammodiscacea** Reuss,1862
- Ammodiscidae** Reuss,1862
- Ammodiscinae** Reuss,1862
- Ammodiscus** Reuss,1862 (TM *Ammodiscus infimus* L.G. Bornemann,1874, non *Orbis infimus* Strickland,1846)
= *Grzybowskia* Myatlyuk,1970 (TM *Cornuspira angusta* Friedberg,1901)
- A. catinus** Höglund,1947
= *A. incertus* : Goës,1894 (p.p.), non d'Orbigny, in de la Sagra,1839
Gullmarfjord at 58-118 m., Koster Channel, Säcken, Skagerrak at 66-700m., Oslofjord
- A. gullmarense** Höglund,1948
= *A. planus* Höglund,1947, non Loeblich,1946
Gullmarfjord at 24-79 m., Oslofjord, Shetlands, Spitsbergen
- A. minimus** Höglund,1947
Skagerrak at 196-352 m.
- A. planorbis** Höglund,1947
Skagerrak at 199-700 m., ?off Portugal
- A. intermedius** Höglund,1947
= *A. tenuis* : Goës,1894, non Brady,1884
Gullmarfjord, Koster Channel, Oslofjord, ?Gulf of Mexico
- A. arenacea** (Williamson,1858)
= *Spirillina arenacea* Williamson,1858
Plymouth
- Arenoturrispirillina** Tairov,1956 (TM *A. aptica* Tairov,1956)
- A. sp.**
Gullmar Fjord at 67-200 m.
- Spirillinoides** Rhumbler,1938
- S. circumcinctus** Rhumbler,1938 (TM)
Helgoland
- Tolyppamminae** Cushman,1928
- Ammolagena** Eimer & Fickert,1899
- A. clavata** (Jones & Parker,1860)
= *Trochammina irregularis* (d'Orbigny) var. *clavata* Jones & Parker,1860 (TM)
= *Webbina clavata* : Brady,1884.; Goës,1894
* Kiae,1900 comments upon a *Webbina tenuicollis* M. Sars from Oslofjord & Lofoten, which greatly remind of *Webbina clavata*, but is yellowish white. Are they synonymous?
Gullmarfjord, Koster Channel, Skagerrak, Oslofjord, Bergen, Trondheim, Faeroe Channel, Greenland, off Ireland, Mediterranean
- Tolyppamina** Rhumbler,1895
= *Serpulella* Eimer & Fickert,1899
- T. vagans** (Brady,1879)
= *Hyperammina vagans* Brady,1879 (TM)
Drøbak, Bergen, Spitsbergen, around Britain
- T. schaudinii** Rhumbler,1903
off Hebrides at 1524 m.
- Ammovertelliniae** Saidova,1981
- Gloomsipira** Rzehak,1885
= *Gordiammina* Rhumbler,1895
- G. charoides** (Jones & Parker,1860)
= *Trochammina squamata* Jones & Parker, var. *charoides* Jones & Parker,1860
Gullmarfjord, Koster Channel, Säcken, Skagerrak at 204-700 m., Oslofjord, Bergen, around Britain, Faeroe Channel, Biscay, Mediterranean
- G. glomerata** Höglund,1947
Kattegatt, Gullmarfjord at 24-67 m., Oslofjord
- G. gordialis** (Jones & Parker,1860)
= *Trochammina squamata* Jones & Parker, var. *gordialis* Jones & Parker,1860 (TM)
Britain, France, Mediterranean, Arctic Ocean
- G. patelliformis** Kiae,1900
* Diffrers from *Ammodiscus incertus* only in the construction of the final convolution"
Drøbak at 100m, Bundeifjord, Frierfjord
- Usbekistaniiae** Vyavlov,1968
- Turritellella** Rhumbler,1904
- T. shoneana** (Siddall,1878)
= *Trochammina shoneana* Siddall,1878 (TM)
Gullmarfjord, North Sea, Britain, Ireland, N Greenland
- T. spectabilis** (Brady,1881)
= *Ammodiscus spectabilis* Brady,1881
off Britain at 358 fathoms
- Rzehakinacea** Cushman,1933
- Rzehakinidae** Cushman,1933
- Miliammina** Heron-Allen & Earland,1930
- M. fusca** (Brady,1870)
= *Quinqueloculina fusca* Brady,1870
= *Quinqueloculina agglutinans* : Brady,1867, non d'Orbigny, in de la Sagra,1839
= *Miliammina earlandi* Loeblich & Tappan,1955
off W Bornholm, Kielr Bucht, Öresund, Oslofjord, Netherlands, Wadden Sea, W North Sea, Wales, W Scotland, Ireland, Eng. Channel
- M. arenacea** (Chapman,1916) : Gabel,1971
= *Miliolina oblonga* (Montagu), var. *arenacea* Chapman,1916, non *Miliolina arenacea* Rhumbler,1906 (TM)
North Sea
- Silicosigmoilina** Cushman & Church,1929 (TM *Silicosigmoilina californica* Cushman & Church,1929)
- S. groenlandica** (Cushman,1933)
= *Quinqueloculina fusca* Brady,1870, var. *groenlandica* Cushman,1933
* Correct genus?
- ?Iceland, W & NE Greenland
- Hormosinacea** Haeckel,1894
- Aschemocellidae** Vyalov,1966
- Kalamopsis** de Folin,1883
- K. vaillanti** de Folin,1883 (TM)
Biscay
- Telaminidae** Loeblich & Tappan,1985
- Tellammina** Gooday & Haynes,1983
- T. fragilis** Gooday & Haynes,1983 (TM)
NE Atlantic
- Tumidotubus** Gooday & Haynes,1983
- T. albus** Gooday & Haynes,1983
NE Atlantic
- Hormosinidae** Haeckel,1894
- Reophacinae** Cushman,1910
- Hormosinella** Shchedrina,1969
- H. distans** (Brady,1881)
= *Reophax distans* Brady,1881 (TM)
Faeroe Channel
- Leptohalysis** Loeblich & Tappan,1984
- L. catella** (Höglund,1947)
= *Reophax catella* Höglund,1947 (TM)
Gullmarfjord at 20-79 m., Skagerrak at 66-305 m., Norway at 150-200 m., England at 20-46 m.
- L. catenata** (Höglund,1947)
= *Reophax catenata* Höglund,1947
* Correct genus?
Gullmarfjord, Malmönfjord, Oslofjord
- L. gracilis** (Kiae,1900)
= *Nodulina gracilis* Kiae,1900
= *Reophax gracilis* : Höglund,1947
* Correct genus?
Kattegatt, Gullmarfjord, Skagerrak, Oslofjord, Drammenfjord, Frierfjord
- L. scottii** (Chaster,1892)
= *Reophax scottii* Chaster,1892
* Correct genus?
Kattegatt, Gullmarfjord, Skagerrak, Drøbak, NE & W Scotland, W Ireland, Engl. Channel
- Nodulina** Rhumbler,1895
- N. dentaliniformis** (Brady,1881)
= *Reophax dentaliniformis* Brady,1881 (TM)
Kattegatt, Gullmarfjord, Koster Channel, Skagerrak, Oslofjord, Bergen, Barents Sea
- Reophax** de Montfort,1808
= *Proteonina* Williamson,1858
= *Lituolina* Goës,1881
- R. subfusciformis** Earland,1933
Oslofjord
- R. curtus** Cushman,1920
= *R. subfusciformis* : Höglund,1947 (p.p.)
= *R. scorpiurus* : Goës,1894 (p.p. - plate 5, fig. 160-163), non Montfort,1808
Kattegatt, Gullmarfjord, Koster Channel, Skagerrak, North Sea, W & NE Greenland, Barents Sea
- R. decorata** (Earland,1933)
= *Proteonina decorata* Earland,1933
* Correct genus?
Kieler Bucht, Drøbak
- R. fusiformis** (Williamson,1858)
= *Proteonina fusiformis* Williamson,1858 (TM of *Proteonina* Williamson,1858)
Kattegatt, Gullmarfjord, Skagerrak, Oslofjord, Engl. Channel, SW Britain, NW Scotland, Wales, Barents Sea, off Greenland, Spitsbergen, Novaja Zemlya
- R. nodulosa** Brady,1879
Koster Channel, North Sea, Novaja Zemlya, Franz Josefs Land
- R. guttifera** Brady,1881
Skagerrak, Oslofjord, Faeroe Channel, Spitsbergen
- R. nana** Rhumbler,1913
=? *R. communis* Lacroix,1930
Kattegatt, Gullmarfjord, Skagerrak, Oslofjord
- R. pilulifera** Brady,1884
= *R. subfusciformis* : Höglund,1947 (p.p. - plate 9, fig. 3.)
Skagerrak, Bergen, off W & SW Ireland
- R. agglutinatus** Cushman,1920
Bergen
- R. regularis** Höglund,1947
= *R. dentaliniforme* : Goës,1894, non Brady,1884
central - S & W Baltic, Gullmarfjord, Koster Channel, Skagerrak, Oslofjord, North Sea
- R. rostrata** Höglund,1947
Gullmarfjord, Koster Channel, Skagerrak, Oslofjord, off Shetlands
- R. scorpiurus** de Montfort,1808 (TM)
Gullmarfjord, Skagerrak, Drøbak, Bergen, Lofoten, N Norway, Engl. Channel, Arcachon, Iceland, NE Greenland, Barents Sea, Arctic Ocean
- R. sp. I** Höglund,1947
Gullmarfjord
- R. sp. II** Höglund,1947
Skagerrak
- R. moniliformis** Siddall,1886
Engl. Channel, Wales, W Scotland, W Ireland
- R. micacea** (Cushman,1918) : Gabel,1971
= *Proteonina micacea* Cushman,1918
North Sea

- R. amblystoma** (Rhumbler,1911)
= *Proteonina amblystoma* Rhumbler,1911
* Correct genus?
off Hebrides at 2275 m.
- R. bacillaris** Brady,1881
off S Rockall Bank, off SW Greenland
- R. sabulosa** Brady,1882
off W Trondheimsfjord at 911m, Faeroe Channel
- R. flexibilis** Schumberger,18??
Barents Sea
- R. cf. tubulosa** (Rhumbler,19??) : Gabel,1971
North Sea
- Scherochorella** Loeblich & Tappan,1984 (TM *Reophax minuta* Tappan,1940)
- Subreophax** Saidova,1975
- S. aduncus** (Brady, in Tizard & Murray,1882)
= *Reophax aduncus* Brady, in Tizard & Murray,1882 (TM)
SW Baltic, Kattegatt, Faeroe Channel
- Cuneatinae Loeblich & Tappan,1984
- Cuneata** K.V. Furstenko,1979
- C. arctica** (Brady,1881)
= *Reophax arctica* Brady,1881 (TM)
= *Bigenerina arctica* : Cushman,1948
NE & W Greenland, Spitsbergen, Novaja Zemlya, Wales, Engl. Channel
- Hormosininae** Haekel,1894
- Archimerismus** Loeblich & Tappan,1984 (TM *Hyperammina subnodosa* Brady,1884)
- A. subnodosa** (Brady,1884)
= *Hyperammina subnodosa* Brady,1884 : Gabel,1971
North Sea, Barents Sea
- Hormosina** Brady,1879
- H. globulifera** Brady,1879 (TM)
Skagerrak at 500-510 m., Barents Sea
- H. carpenteri** Brady,1881
Faeroe Channel, Rockall Bank, W Ireland
- H. normani** Brady,1881
off NW Ireland at 2524 m.
- Loeblichopsis** Hofker,1967
- L. cylindricus** (Brady,1884)
= *Reophax cylindrica* Brady,1884 (TM)
off SW Greenland at 3200 m.
- Reophanus** Saidova,1970
- R. ovicula** (Brady,1879)
= *Hormosina ovicula* Brady,1879 (TM)
Barents Sea
- Siliconodosarina** Colom,1963
- S. delicatula** Colom,1963 (TM)
Vigo Bay
- Thomasnellidae** Loeblich & Tappan,1984
- Protoschista** Eimer & Fickert,1899
- P. findens** (Parker, in Dawson,1870)
= *Lituola findens* Parker, in Dawson,1870) (TM)
= *Reophax findens* : Auctt.
NW, N & NE Greenland, off Ireland
- Lituolacea de Blainville,1827
- Haplophragmoididae** Maync,1952
- Buzasina** Loeblich & Tappan,1985 (TM *Trochammina ringens* Brady,1879)
= *Cystaminella Lukina,1980*, non Myatluik,1966
- Cribrostomoides** Cushman,1910
- C. subglobosa** (G.O. Sars,1872)
= *Lituola subglobosa* G.O. Sars,1872
= *Cribrostomoides bradyi* Cushman,1910 (TM)
= *Labrospira subglobosa* : Höglund,1947
Koster Channel, Skagerrak at 500-700 m., Oslofjord, Bergen, off Britain, Spitsbergen, Franz Josefs Land, Arctic Ocean
- C. nitida** (Goës,1896)
= *Haplophragmium nitidum* Goës,1896
= *Labrospira nitida* : Höglund,1947
Koster Channel, Skagerrak at 200-700 m., Oslofjord, Caribbean
- C. jeffreysi** (Williamson,1858)
= *Nontionina jeffreysi* Williamson,1858
= *Labrospira jeffreysi* : Höglund,1947
= *Alveophragmium jeffreysi* : Loeblich & Tappan,1953
?Arcona Basin, Gullmarfjord, Skagerrak at 68-305 m., Engl. Channel, Ireland, Barents Sea, NE & W Greenland
- C. kosterensis** (Höglund,1947)
= *Labrospira kosterensis* Höglund,1947
Gullmarfjord, Koster Channel, Oslofjord, North Sea, Bergen, Hardangerfjord, Greenland
- C. wiesneri** Parr,19?? : Gabel,1971
North Sea
- C. arctica** (Parker,1???) : Gabel,1971
North Sea
- Haplophragmoides** Cushman,1910
- = *Recurvoidella* Uchio,1960 (TM *R. parkerae* Uchio,1960)
= *Linguiferina* Alekseychik-Mitskevich,1973 (TM *Haplophragmoides subsphaeroides* Subbotina,1950)
= *Pauciculina* Alekseychik-Mitskevich,1973 (TM *Haplophragmoides Kirkii* Wickenden,1932)
= *Subtilina* Alekseychik-Mitskevich,1973 (TM *Haplophragmoides tenius* Cushman,1927)
= *Pseudohaplophragmoides* Saidova,1981 (TM *Haplophragmoides ultimus* Saidova,1975)
- H. bradyi** (Robertson,1891)
= *Trochammina robertsoni* Brady,1887, : Goës,1894
= *Trochammina bradyi* Robertson,1891
Kattegatt, Gullmarfjord, Skagerak, Drøbak, Bergen, Bukkenfjord, Vadsø, off W Britain, off SW Ireland
- H. fragile** Höglund,1947
Gullmarfjord at 15-80 m., Oslofjord
- H. membranaceum** Höglund,1947
Gullmarfjord at 24-79 m., Oslofjord
- H. pusillum** Höglund,1947
Gullmarfjord, Oslofjord
- H. wilberti** Anderson,1953
?= *Cribrostomoides jeffreysi* : Lutze,1965, non (Williamson,1858)
?SW Baltic, Wales
- H. scitulum** (Brady,1881)
= *Haplophragmium scitulum* Brady,1881
Faeroe Channel, Cape Verde
- H. canariense** (d'Orbigny,1839)
= *Nontionina canariense* d'Orbigny,1839 (TM)
Högloland, Barents Sea, Novaja Zemlya
- H. major** Cushman,1920
Iceland
- Labrospira** Höglund,1947
- L. crassimargo** (Norman,1892)
= *Haplophragmium crassimargo* Norman,1892 (TM)
Kattegatt, Gullmarfjord, Skagerrak at 66-300 m., Oslofjord, North Sea, N Norway, Spitsbergen, Barents Sea, W & NE Greenland
- Discamminidae** Mikhalevich,1980
- Ammoscalaria** Höglund,1947
- A. tenuimargo** (Brady, in Tizard & Murray,1882)
= *Haplophragmium tenuimargo* Brady, in Tizard & Murray,1882 (TM)
Koster Channel, Skagerrak at 200-700 m., Oslofjord, Faeroe Channel
- A. pseudospiralis** (Williamson,1858)
= *Proteonina pseudospiralis* Williamson,1858
= *Ammobaculites pseudospirale* : Cushman,1920
Arkona Basin, Kattegatt at 15-55 m., Gullmarfjord, Skagerrak at 68-106 m., Drøbak, Bergen, Iceland, W Scotland, Ireland, Cornwall, Arachon, Galicia
- A. runiana** (Heron-Allen & Earland,1916)
= *Haplophragmium runianum* Heron-Allen & Earland,1916
= *Ammodiscus rhumbleri* Rottgardt,1952
Belt Sea, Kattegatt, at 0-10 m., Gullmarfjord at 9-33 m., Drøbak, Plymouth, W Scotland
- Discammina** Lacroix,1932
- D. compressa** (Goës,1882) : Gabel,1971
= *Lituolina irregularis* var. *compressa* Goës,1882
= *Discammina fallax* Lacroix,1932 (TM)
North Sea, Mediterranean
- Lituotubidae** Loeblich & Tappan,1984
- Trochamminoides** Cushman,1910 (TM *Trochammina proteus* Karrer,1866)
- Lituolidae** de Blainville,1827
- Ammomarginulinidae** Podobina,1978
- Ammobaculites** Cushman,1910 (TM *Spirolina agglutinans* d'Orbigny,1846 - Miocene)
- A. balkwilli** Haynes,1973
= *Haplophragmium agglutinans* : Balkwill & Wright,1885, non *Spirolina agglutinans* d'Orbigny,1846
Irish Sea, Wales
- A. agglutinans** (d'Orbigny,1846)
= *Spirolina agglutinans* d'Orbigny,1846
* may have been misidentified *A. balkwilli* Haynes
Oslofjord
- Ammomarginulina** Wiesner,1931 (TM *A. ensis* Wiesner,1931)
- Ammotium** Loeblich & Tappan,1953
- A. cassis** (Parker, in Dawson,1870)
= *Lituola cassis* Parker, in Dawson,1870 (TM)
= *Ammobaculites cassis* : Auctt.
W Baltic, Kattegatt, Uddevalla, Oslofjord (dominating in the upper 10 m.), Spitsbergen, Barents Sea, NE & W Greenland
- Eratidus** Saidova,1975
- E. foliaceum** (Brady,1881)
= *Haplophragmium foliaceum* Brady,1881 (TM)
= *Ammobaculites foliaceus* : Auctt.
Iceland, central N Atlantic
- Flabellaminidae** Podobina,1978
- Triplasia** Reuss,1854 (TM *T. murchisoni* Reuss,1854)
- T. bicamerata** (Heron-Allen & Earland,1930)
= *Orthocerina bicamerata* Heron-Allen & Earland,1930
* Correct genus???
- Plymouth
- Lituolinae** de Blainville,1827
- Lituola** de Lamarck,1804 (TM *Lituolites nautiloidea* de Lamarck,1804)
= *Lituolites* de Lamarck,1804
- Placopsislinidae** Rhumbler,1913
- Placopsislinina** Rhumbler,1913
- Placopsisilina** d'Orbigny,1850
- P. cenomana** d'Orbigny,1850 (TM)
S Britain - southwards to tropical & subtropical seas
- P. confusa** Cushman,1920

- Gullmarfjord at 35–50 m.
- Haplophragmiacea Eimer & Fickert,1899
- Ammosphaeroidinidae Cushman,1927
- Ammosphaeroidininae Cushman,1927
- Adercotryma Loeblich & Tappan,1952
- A. glomeratum* (Brady,1878)
= *Lituola glomerata* Brady,1878 (TM)
= *Haplophragmoides glomeratum* : Auctt.
Öresund, Kattegatt, Gullmarfjord, Oslofjord, Scotland, Ireland, NW & NE Greenland, Novaja Zemlya, Franz Josef's Land, Japan
- Ammosphaeroidina* Cushman,1910 (TM) *Haplophragmium sphaeroidiniformae* Brady,1884
- Cystammina* Neumayr,1889 (TM) *Trochammina pauciloculata* Brady,1879
- Recurvoidinae Alekseychik-Mitskevich,1973
- Recurvoidea* Earland,1934 (TM) *R. contortus* Earland,1934
- R. laevigatum* Höglund,1947
Koster Channel, Skagerrak, Oslofjord
- R. trochanminiforme* Höglund,1947
Kattegatt, Gullmarfjord, Skagerrak, Oslofjord
- R. turbinatus* (Brady,1881)
= *Haplophragmium turbinatum* Brady,1881
= *Trochammina turbinata* : Cushman,1948
Spitsbergen, NE & NW Greenland
- Thalmannammina* Pokorný,1951 (TM) *Haplophragmium subturbinatum* Grzybowski,1898
= *Conglophragmium* Bermúdez & Rivero,1963 (TM) *Trochammina conglobata* Brady,1884
- Coscinophragmatacea Thalmann,1951
- Coscinophragmatidae Thalmann,1951
- Ammotrochoidea* Janin,1984 (TM) *A. bignoti* Janin,1984
- Loftusiacea Brady,1884
- Cyclamminidae Marie,1941
- Alveolophragmiinae Saidova,1981
- Alveolophragmium Shchedrina,1936
- A. orbiculatum* Shchedrina,1936 (TM)
= *Haplophragmoides subglobosum* : Colom,1952, non *Lituola subglobosa* G.O. Sars,1872
Kara Sea, Sea of Okhotsk, W approaches to the Engl. Channel, off Galicia
- A. wiesneri* (Parr,19??) : Gabel,1971
North Sea
- Cyclammininae Marie,1941
- Cyclammina* Brady,1879
- C. cancellata* Brady,1879 (TM)
W approaches to the Engl. Channel at >420 m., off Galicia
- Spirolectamminacea Cushman,1927
- Spirolectamminidae Cushman,1927
- Spirolectammininae Cushman,1927
- Spirolectammina* Cushman,1927
- S. biformis* (Parker & Jones,1865)
= *Textularia agglutinata* d'Orbigny var. *biformis* Parker & Jones,1865 (TM)
SW Baltic, Öresund, Kattegatt, Gullmarfjord, Skagerrak, Oslofjord, Spitsbergen, Barents Sea, Engl. Channel, Davis Strait, NE Greenland, Arctic Ocean
- S. earlandi* (Parker,1952)
= *Textularia elegans* Lacroix,1931, non *Pelecanium elegans* von Hantken,1876?
= *Textularia earlandi* Parker,1952
= *Textularia tenuissima* Earland,1933, non Hausler,1881
Kattegatt, Gullmarfjord, Skagerrak, Drøbak, Faeroes, SW Ireland, Cornwall, Wales, Engl. Channel, Mediterranean
- S. fusca* (Earland,1905)
= *Spirolecta fusca* Earland,1905
* Correct genus???:
around Britain
- S. sp. : Gabel,1971
Skagerrak
- Spirolectinella* Kisel'man,1972 (TM) *Spirolecta wrightii* Silvestri,1903
- S. wrightii* (Silvestri,1903)
= *Spirolecta sagittula* Wright,1891, non *Textularia sagittula* Defrance, in de Blainville,1824
= *Spirolecta wrightii* Silvestri,1903
= *Textularia sagittula* : Auct., non Defrance, in de Blainville,1824
= *Spirolectammina wrightii* : Auctt.
? = *Textularia williamseni* Goës,1894
Gullmarfjord, Drøbak, Skagerrak, North Sea, Belgium, Plymouth, W Britain, Arcachon, Galicia, Mediterranean
- Vulvulininae Saidova,1981
- Vulvulina* d'Orbigny,1826 (TM) *V. capreolus* d'Orbigny,1826 = *Nutilus (Orthoceras) pennatula* Batsch,1791
= *Schizophora* Reuss,1861 (TM) *S. neugeboreni* Reuss,1861
= *Venilina* Gümbel,1870 (TM) *V. nummulina* Gümbel,1870
* *Bigenerina sarsi* Kiaer,1900 from Lofoten, 200–1200m, is considered to "differ from *B. capreolus*, and *pennatula* by the more horizontally connected segments, their far greater number in the biserial series, and the elongated form, tapering towards the apex"
- Morulaepletina* Saidova,1981
- Morulaeplecta* Höglund,1947
- M. bulbosa* Höglund,1947 (TM)
Kattegatt, Gullmarfjord at 20–50 m., Skagerrak, Oslofjord
- Pseudobolivinidae Wiesner,1931
- Lacroixina Saidova,1981
- L. coeruleata* (Lacroix,1932)
= *Textularia coeruleata* Lacroix,1932 (TM)
* Only known from one Mediterranean & one Swedish specimen.
Gullmar Fjord at 32 m., Mediterranean
- Parvigenerina* Vella,1957 (TM) *Bifarina porrecta* (Brady) var. *arenacea* Heron-Allen & Earland,1922
- Nouriidae Chapman & Parr,1936
- Nouria* Heron-Allen & Earland,1914 (TM) *N. polymorphoides* Heron-Allen & Earland,1914
- Trochamminacea Schwager,1877
- Trochamminidae Schwager,1877
- Trochammininae Schwager,1877
- Ammoglobigerina* Eimer & Fickert,1899
= *Trochoporina* Soliman,1972 (TM) *Trochoporina praeglobigeriniformis* Soliman,1972
= *Globotrochamminopsis* Brönnimann & Whittaker, in Brönnimann & Annetti,1984 (TM) *Trochammina globulosa* Cushman,1920
- A. globigeriniformis* (Parker & Jones,1865)
= *Lituola nautiloidea* de Lamarck var. *globigeriniformis* Parker & Jones,1865
= *Ammoglobigerina bulboides* Eimer & Fickert,1899 (TM)
Bergen, North Sea, N Atlantic, Franz Josef's Land
- A. globigeriniformis* (Parker & Jones,1865) var. *pygmaea* (Höglund,1947)
= *Trochammina globigeriniformis* Parker & Jones,1865 var. *pygmaea* Höglund,1947
Skagerrak at 400–515 m., S & W Britain, Engl. Channel, SW Ireland
- Paratrochammina* Brönnimann,1979 (TM) *P. madeirae* Brönnimann,1979
- Tritaxis Schubert,1921
- T. fusca* (Williamson,1858)
= *Rotolina fusca* Williamson,1858 (TM)
= *Valvulina fusca* : Auctt.
* probably clinging to firm substrates.
Skagerrak at 400–626 m., Drøbak, Lofoten, Belgium, Scotland, Shetlands, Iceland, Barents Sea, W Scotland, W Ireland, Galicia, Canaries
- T. conica* (Parker & Jones,1865)
= *Valvulina conica* : Auctt.
= *Valvulina triangularis* d'Orbigny,1826 var. *conica* Parker & Jones,1865
Gullmarfjord, Skagerrak at 200–700 m., Oslofjord, Bergen, Engl. Channel, W Scotland, Faeroe Channel, Mediterranean
- Trochammina* Parker & Jones,1859
- T. inflata* (Montagu,1808)
= *Nautillus inflatus* Montagu,1808 (TM)
German Baltic, Wadden Sea, Devon - in brackish water, Wales, NW Scotland, Engl. Channel, Biscay, Mediterranean
- T. rotaliformis* Wright, in Heron-Allen & Earland,1911
* may cling to firm substrates.
Koster Channel, Skagerrak, Drøbak, Spitsbergen, Belgium, Engl. Channel, off Ireland, Arcachon, NE & W Greenland
- T. vesicularis* Goës,1894
* Correct genus?
Spitsbergen
- T. adaperata* Rhumbler,1938
= *Trochammina squamata* Jones & Parker var. *adaperata* Rhumbler,1938
Gullmarfjord, Koster Channel, Skagerrak, Oslofjord, Helgoland
- T. squamata* Jones & Parker,1860
Novaja Zemlya, Britain, Ireland
- T. squamata* Jones & Parker, var. *pluricubiculata* Rhumbler,1938
Helgoland
- T. squamata* Jones & Parker, var. *obtusa* Rhumbler,1938
Helgoland
- T. astrifica* Rhumbler,1938
= *Trochammina squamata astrifica* Rhumbler,1938
Kattegatt, Gullmarfjord, Skagerrak, Helgoland, Plymouth, Wales
- T. intermedia* Rhumbler,1938
= *T. squamata* Jones & Parker, var. *intermedia* Rhumbler,1938
Skagerrak at 66 m., Helgoland, Ireland, Wales
- T. labiosa* Höglund,1947
Gullmarfjord threshold, Skagerrak, Oslofjord
- T. stellata* Höglund,1947
Kattegatt, mouth of Gullmarfjord, Koster Channel, Skagerrak
- T. multiloculata* Höglund,1947
Koster Channel, Skagerrak
- T. nana* (Brady,1881)
= *Haplophragmium nanum* Brady,1881
North Sea, NE & NW Greenland, Novaja Zemlya, Franz Josef's Land, Arctic Ocean
- T. haynesi* Atkinson,1969
?= *T. sp.* : Lutze,1965
?SW Baltic, Wales
- T. karica* Stschedrina,19??
Barents Sea
- T. aff. laevigata* Cushman & Brönnimann,1948 : Murray,1965
Engl. Channel
- T. ?* sp. : Gabel,1971
Sylt
- Trochamminopsis* Brönnimann,1976
- T. quadriloba* (Höglund,1948)
= *Trochammina pusilla* Höglund,1947 (TM), non *Serpula pusilla* Geinitz,1848 (= *Trochammina pusilla* (Geinitz))
Jones, Parker & Kirby,1869
= *Trochammina quadriloba* Höglund,1948
Koster Channel, Skagerrak, Oslofjord, E Greenland
- Trochamminellinae* Brönnimann, Zaninetti & Whittaker,1983

- Pseudotrochammina** Frerichs,1969 (TM *P. triloba* Frerichs,1969)
= *Anmoglobigerinoides* Frerichs,1969 (TM *A. dehiscens* Frerichs,1969)
- P.** sp.
Skagerrak
- Trochamminella** Cushman,1943 (TM *T. siphonifera* Cushman,1943)
- T. bullata** Höglund,1947
Skagerrak at 500-700 m., Oslofjord
- T. atlantica** F. Parker,1952
NE & NW Greenland
- Jadammininae** Saidova,1981
- Jadammina** Bartenstein & Brand,1938
= *Borovina* Shmal'gausen,1950 (TM *B. zernovi* Shmal'gausen,1950)
= *Trochamminisca* Shchedrina,1955 (TM *T. cyclostoma* Shchedrina,1955)
- J. macrescens** (Brady,1870)
= *Trochammina inflata* (Montagu) var. *macrescens* H.B. Brady, in G.S. Brady, Robertson & H.B. Brady,1870
= *Jadammina polystoma* Bartenstein & Brand,1938 (TM)
Poland, SW Baltic, Oslofjord, German North Sea coast, Netherlands, Belgium, Engl. Channel, W Scotland, Ireland, Wales
- Arenoparrellinae** Saidova,1981
- Trochamminula** Shchedrina,1955
- T. fissuraperta** Shchedrina,1955 (TM)
Okhotsk Sea, Barents Sea
- Polystomammininae** Brönnimann & Beurlen,1977
- Deuterammina** Brönnimann,1976 (TM *Trochammina glabra* Heron-Allen & Earland,1932)
- D. (Centrodeuterammina)** Brönnimann & Whittaker,1983 **dublinensis** Brönnimann & Whittaker,1983
off Ireland
- Lepidodeuterammina** Brönnimann & Whittaker,1983
- L. ochracea** (Williamson,1858)
= *Rotalina ochracea* Williamson,1858 (TM)
= *Trochammina ochracea* : Auct.
* clinging to pebbles, sea-weeds etc. in current-swept areas
Gullmarfjord, Skagerrak, Oslofjord, Helgoland, Belgium, Shetlands, Barents Sea, NW Scotland, Ireland, Engl. Channel, Wales, Arcachon
- L. ochracea** (Williamson,1858) **heronearelandica** (Rhumbler,1938)
= *Trochammina ochracea* (Williamson,1858) *heronearelandica* Rhumbler,1938
* Correct genus?
Helgoland
- Polystomammina** Seiglie,1965
= *Arcoarella* Mikhalovich,1971 (TM *A. planulata* Mikhalovich,1971)
- P. nitida** (Brady,1884)
= *Trochammina nitida* Brady,1884 (TM)
SW Baltic, W Scotland, off SW Ireland, N Norway, Novaja Zemlya, Franz Josefs Land
- Remaneicidae** Loeblich & Tappan,1964
- Remaneicinae** Loeblich & Tappan,1964
- Remaneica** Rhumbler,1938
- R. helgolandica** Rhumbler,1938 (TM)
* The Skagerrak material may possibly be *R. plicata* (Terquem,1875)
?Skagerrak at 66-305 m., Helgoland, Wales
- R. plicata** *plicata* (Terquem,1875)
= *Patellina plicata* Terquem,1875
= *Trochammina plicata* : Auct.
Helgoland, W Scotland, W Ireland, W France
- R. plicata** (Terquem,1875) **lichenopsis** Rhumbler,1938
Helgoland, W Scotland
- Septotrochammina** S.Y. Zheng,1979 (TM *S. plicata* (Terquem) sensu S.Y. Zheng, 1979, non *Patellina plicata* Terquem,1876 (= *Remaneica gonzalezi* Seiglie,1964))
= *Remaneicella* Brönnimann, Zaninetti & Whittaker,1983 (TM *Remaneica gonzalezi* Seiglie,1964)
- Verneuilinaceae** Cushman,1911
- Prolixoplectidae** Loeblich & Tappan,1985
- Karrerulina** Finlay,1940 (TM *Gaudryina apicularis* Cushman,1911)
- Verneuiliniidae** Cushman,1911
- Verneuilininae** Cushman,1911
- Gaudryina** d'Orbigny,1839 (TM *Gaudryina rugosa* d'Orbigny,1840)
- G. rufa** Wright,1900
* clinging to firm substrates.
S & W Britain, Ireland, Biscay
- Verneuilina** d'Orbigny,1839 (TM *Verneuilina tricarinata* d'Orbigny,1840)
- V. media** Höglund,1947
Kattegatt, Gullmarfjord, Skagerrak, Oslofjord
- V. europea** Christiansen,1958
= *V. pusilla* : Heron-Allen & Earland,1913, non Goës,18??
= *V. advena* : Höglund,1947, non Cushman,1920
Kattegatt, Gullmarfjord, Skagerrak, Oslofjord
- V. polystropha** Reuss,1846
Novaja Zemlya, Faeroes, off W Ireland, W Scotland, Arctic Ocean
- V. minuta** Wiesner,19?? : Gabel,1971
North Sea
- V. propinqua** Brady,1884
warm Faeroe Channel
- Ataxophragmiacea** Schwager,1877
- Globotextulariidae** Cushman,1927
- Globotextulariinae** Cushman,1927
- Globotextularia** Eimer & Fickert,1899
- G. anceps** (Brady,1884)
= *Haplophragmium anceps* Brady,1884 (TM)
Southport, Irish Sea, W Ireland
- Liebusellinae** Saidova,1981
- Liebusella** Cushman,1933 (TM *Lituola nautiloidea* de Lamarck var. *soldanii* Jones & Parker,1860)
- L. goesi** Höglund,1947
= *Bigenerina digitata* : Goës,1894, non d'Orbigny,1826
= *Bigenerina cylindrica* : Nørvang,1941, non Cushman,1922
Kattegatt, Gullmarfjord, Koster Channel, Skagerrak, Oslofjord, Bergen
- Textulariellidae** Grönhagen & Lutherbacher,1966
- Textulariella** Cushman,1927 (TM *Textularia barrettii* Jones & Parker,1876)
= *Neocuneolina* Bermúdez & Rivero,1963 (TM *Cuneolina pavonia* d'Orbigny var. *angusta* Cushman,1919)
- Textulariacea** Ehrenberg,1838
- Eggerellidae** Cushman,1937
- Dorothiinae** Balakhmatova,1972
- Bannerella** Loeblich & Tappan,1985 (TM *Textularia gibbosa* d'Orbigny,1826)
- B. pseudoturris** (Cushman,1922)
= *Textularia pseudoturris* Cushman,1922
= *Dorothia pseudoturris* (Cushman,1937)
* Correct genus?
?Skagerrak, ?Oslofjord
- Eggerellinae** Cushman,1937
- Eggerella** Cushman,1935
- E. advena** (Cushman,1920)
= *Verneuilina advena* Cushman,1920
= *Eggerella arctica* Höglund,1947
non = *Verneuilina advena* : Höglund,1947
= *Verneuilina pygmaea* : Goës,1894, non (Egger,1857), nec Brady,1884
Spitsbergen, NE & W Greenland
- E. affixa** (Cushman,1911) : Gabel,1971
North Sea
- E. bradyi** (Cushman,1911)
= *Verneuilina bradyi* Cushman,1911 (TM)
= *Verneuilina pygmaea* : Brady,1884, non *Bulimina pygmaea* Egger,1857
W Ireland, Faeroe Channel, off SW Ireland at 1370-1866 m.
- Eggerelloides** Haynes,1973
- E. scabrum** (Williamson,1858)
= *Bulimina scabra* Williamson,1858 (TM)
= *Eggerella scabra* : Auct.
= *Verneuilina scabra* : Auct.
Arkona Sea, Belt Sea, Øresund, Kattegatt, Gullmarfjord, Skagerrak, Drøbak, Iceland, S North Sea, all around Britain, Spain
- Karrerellina** Cushman,1933 (TM *Gaudryina siphonella* Reuss,1851)
= *Valvotextularia* Hofker,1951 (TM *Textularia catenata* Cushman,1911)
- K. bradyi** (Cushman,1911)
= *Gaudryina bradyi* Cushman,1911
Celtic Sea, western approaches of the Engl. Channel
- Martinottiella** Cushman,1933
= *Listerella* Cushman,1933, non Jahn,1906 (TM *Clavulina primaeva* Cushman,1913)
- M. communis** (d'Orbigny,1826)
= *Clavulina communis* d'Orbigny,1826 (TM)
Gullmarfjord, Celtic Sea
- Meidamonella** Loeblich & Tappan,1986
- M. novangliae** (Cushman,1922)
= *Gaudryina bacata* Schwager,1866, var. *novangliae* Cushman,1922 (TM)
Faeroe Channel
- Textulariidae** Ehrenberg,1838
- Textulariinae** Ehrenberg,1838
- Bigenerina** d'Orbigny,1826
- B. nodosaria** d'Orbigny,1826 (TM)
Bergen, around Britain
- B. cylindrica** Cushman,1922
= *Bigenerina digitata* : Brady,1864, non *Gemmulina digitata* d'Orbigny,18??
?Drøbak, SW Ireland
- B. robusta** Brady,1881
off Shetlands
- Sahulia** Loeblich & Tappan,1985 (TM *Sahulia patelliformis* Loeblich & Tappan,1985 = *Textularia barkeri* Hofker,1978)
- Textularia** Defrance,1824 (TM *T. sagittula* Defrance, in de Blainville,1824)
= *Textilina* Nørvang,1966 (TM *Textularia stricta* Cushman,1911)
= *Valvulinema* Saidova,1975 (TM *Textularia milletti* Cushman,1911)
= *Textella* Mikhalovich,1979 (TM *Textularia foliacea* Heron-Allen & Earland var. *occidentalis* Cushman,1922)
= *Norvanginella* Mikhalovich,1981 (TM *Textilaria pseudorugosa* Lacroix subsp. *fistulosa* Mikhalovich,1973, non *Textularia sagittula* var. *fistulosa* Brady,1884 (= *Textularia valerieae* Loeblich & Tappan,1988))
= *Textilinata* Botvinnik,1983 (TM *Textilaria cognita* Botvinnik,1983 (= *Textularia sagittula* Defrance,1824))
- T. sagittula** Defrance, in de Blainville,1824
= *Spiroplectammina sagittula* (Defrance,1824) : Gabel,1971
North Sea
- T. sagittula** Defrance, in de Blainville,1824, var. *jugosa* Brady,18??
W Scotland
- T. bocki** Höglund,1947
= *Textilina bocki* : Nørvang,1966
= *Textularia agglutinans* : Goës,1894, non d'Orbigny,1839
= *Textularia gramine* : Balkwill & Wright, non *Valvulina gramine* d'Orbigny,1839
Kattegatt, Gullmarfjord, Malmönfjord, Skagerrak at 29-305 m., Oslofjord, Irish Sea, Celtic Sea, Biscay

- T. truncata* Höglund,1947
Gåsö Channel at ~15 m., Hållö at ~55-70 m, Oslofjord
- T. kategattensis* Höglund,1948
= *T. gracillima* Höglund,1947, non Andreæ, 1884
Öresund, Kattegatt, Gullmarfjord, Skagerrak, Oslofjord
- T. bigenerinoides* Lacroix,1932
Gullmarfjord, Oslofjord, ?Engl. Channel, Mediterranean
- T. contorta* Höglund,1947
Gullmarfjord
- T. skagerakensis* Höglund,1947
Skagerrak at 199-626 m., Oslofjord
- T. torquata* F. Parker,1952
Spitsbergen, W & NE Greenland
- T. agglutinans* d'Orbigny, in de la Sagra,1839
Cuba, ?Arctic Ocean, ?North Sea, ?Iceland
- T. conica* d'Orbigny, in de la Sagra,1839
off Britain
- T. fusiformis* Defrance,18??
W Scotland
- T. globulosa* Ehrenberg,18??
off Ireland
- T. aspera* Brady,18??
off SE Ireland at 3235 m.
- T. cuneiformis* d'Orbigny,1826
Iceland
- T. candeiana* d'Orbigny, in de la Sagra,1839
?Britain, Cuba
- Siphonotularinae Loeblich & Tappan,1985
- Siphonotularia Finlay,1939 (TM S. wairoana Finlay,1939)
- S. flintii* (Cushman,1911)
= *Textularia flintii* Cushman,1911
western approaches of the Engl. Channel at >420 m.
- S. rolshauseni* (Phleger & Parker,19??) : Gabel,1971
North Sea
- S. concava* (Karrer,18??)
= *Textularia concava* Karrer,18??
W Ireland, off SW Ireland
- S. concava* (Karrer,18??), var. *heterostoma* Formasini,19??
off W Scotland
- Planctostomatinae Loeblich & Tappan,1984
- Planctostoma* Loeblich & Tappan,1955 (TM *Textularia luculenta* Brady,1884)
- Pseudogaudryniidae Loeblich & Tappan,1985
- Pseudogaudryniinae Loeblich & Tappan,1985
- Migros* Finlay,1939 (TM *Gaudryina medwayensis* Parr,1935)
- Pseudogaudryina* Cushman,1936 (TM *Textularia atlantica* Bailey,1851)
- Valvulinidae Berthelin,1880
- Valvulininae Berthelin,1880
- Clavulina* d'Orbigny,1826 (TM *C. parisiensis* d'Orbigny,1826)
- C. obscura* Chaster,1892
S & W Britain, Celtic Sea, W Ireland
- C. sp.* : Gabel,1971
North Sea
- Goesella* Cushman,1933 (TM *Clavulina rotundata* Cushman,1913)
- Valvulina* d'Orbigny,1826
- V. triangularis* d'Orbigny,1826 (TM)
- Sussex
- Tritaxilininae Loeblich & Tappan,1986
- Tritaxilina* Cushman,1911 (TM *Clavulina caperata* Brady,1881)
- FUSULININA Wedekind,1937
* Not including any recent taxa.
- INVOLUTININA Hohenegger & Piller,1977
- Planispirillinidae Piller,1978
- Planispirillina* Bermúdez,1952 (TM *Spirillina limbata* Brady, var. *papillosa* Cushman,1915 = *Spirillina margaritifera* Terquem,1880, non Williamson,1858 = *Spirillina terquemi* A. Silvestri,1923)
- P. wrightii* (Heron-Allen & Earland,1930)
= *Spirillina wrightii* Heron-Allen & Earland,1930
= *Spirillina margaritifera* : Wright,1877, : Terquem,1875, : 1881, : Halkyard,1889, non Williamson,1858
Plymouth
- SPIRILLININA Hohenegger & Piller,1975
- Spirillinidae Reuss & Fritsch,1861
- Mychostomina* Berthelin,1881 (TM *Spirillina vivipara* Ehrenberg var. *revertens* Rhumbler,1906)
- Sejunctella* Loeblich & Tappan,1957
- S. late septata* (Terquem,1875)
= *Spirillina late septata* O. Terquem,1875
= *Sejunctella earlandi* Loeblich & Tappan,1957 (TM)
Faeroes, Belgium, Engl. Channel
- Spirillina* Ehrenberg,1843
- S. perforata* (Schultze,1854)
= *Cornuspirna perforata* : Schultze,1854
Netherlands, Belgium, Engl. Channel, Wales, off Mozambique
- S. vivipara* Ehrenberg,1843 (TM)
* clinging to firm substrates.
Bergen, N Norway, Belgium, Engl. Channel, Biscay
- S. vivipara* Ehrenberg,1843 *ruriana* Heron-Allen & Earland,1930
* clinging to firm substrates.
Belgium, Engl. Channel, Biscay, W Scotland
- S. obconica* Brady,1879
off Faeroes, W Ireland, Plymouth
- S. obconica* Brady,1879 var. *carinata* Halkyard,1889
= *S. vivipara* Ehrenberg,1843 var. *carinata* Halkyard,1889
* Correct genus?
Plymouth
- S. groomii* Chapman,1900
Plymouth, W Ireland
- S. limbata* Brady,1879, var. *denticulata* Brady,1884
= *Spirillina wrightii* Heron-Allen & Earland,1930
Belgium, Plymouth
- S. margaritifera* Williamson,1858
?Plymouth
- S. lucida* Sidebottom,1908
W Ireland, Mediterranean
- S. canaliculata* Terquem,18??
Engl. Channel
- S. nodosa* Terquem,18??
Engl. Channel
- S. arctica* Cushman,19??
Barents Sea
- S. sp. A* : Haynes,1973
Wales
- Turrispirillina* Cushman,1927 (TM *Spirillina conoidea* Paalzow,1917)
- T. sp. A* : Haynes,1973
Wales
- Patellinidae Rhumbler,1906
- Hergottiniae Loeblich & Tappan,1984
- Patellinoides Cushman,1933 (TM *Patellinoides conicus* Heron-Allen & Earland,1932)
- Patellininae Rhumbler,1906
- Patellina* Williamson,1858
- P. corrugata* Williamson,1858 (TM)
* clinging to firm substrates.
Oslofjord, Bergen, Vesterålen, Spitsbergen, Novaja Zemlya, German North Sea, Belgium, Faeroe Channel, Iceland, Shetlands, W Scotland, N Kent, Engl. Channel, Bristol Channel, Wales, Ireland, NE, N, NW & W Greenland, Novaja Zemlya, Barents Sea, Belgium, Arcachon
- CARTERININA Loeblich & Tappan,1981
* Recent species with tropical shallow water distribution.
- Miliolina Delage & Hérouard,1896
- Squamulinacea Reuss & Fritsch,1861
- Squamulinidae Reuss & Fritsch,1861
- Squamulininae Reuss & Fritsch,1861
- Squamulina* Schultze,1854
- S. laevis* Schultze,1854 (TM)
Baltic, North Sea, Britain, Mediterranean
- Cornspiraceae Schultze,1854
- Cornspiridae Schultze,1854
- Cornspirinae Schultze,1854 = *Cornspiroidinae* Saidova,1981
- Cornuspira* Schultze,1854
= *Cyclogrya* S.V. Wood,1842 (TM *Cyclogrya multiplex* S.V. Wood,1842 - Gen. & sp. suppressed for purposes of Law of Priority, not Law of Homonymy - ICZN Op. 1114)
= *Cornuspirella* Cushman,1928
= *Cornuspirodes* Cushman,1928
- C. foliacea* (Philippi,1844)
= *Orbis foliaceus* Philippi,1844 (TM)
= *Cornuspira diffusa* Heron-Allen & Earland,1913 (TM of *Cornuspirella* Cushman,1928)
= *Cornuspira striata* Brady, in Tizard & Murray,1882 (TM of *Cornuspirodes* Cushman,1928)
= *Cornuspira difformis* Cushman,1929
= *Cornuspira profundum* Schedlerina,1946
Bohus, Oslofjord, Bergen, Central North Sea, Barents Sea, Spitsbergen, N Siberia, Iceland, Greenland, Faeroes, Shetlands, Orkneys, off W & NW Scotland, N & W Ireland, Engl. Channel
- C. involvens* (Reuss,1850)
= *Operculina involvens* Reuss,1850
= *Cornuspira marginata* M. Sars,18??
* clinging to firm substrates.
Bergen, Lofoten, N Norway, Spitsbergen, Novaja Zemlya, Belgium, S Britain, NE & W Greenland
- C. selsevensis* Heron-Allen & Earland,1909
= *Cyclogrya selsevensis* : Auctt.
* Found attached to pleopods of *Nephrops norvegicus* (Linnaeus,1758)
North Sea, Engl. Channel, Wales, Isle of Man, Mediterranean
- C. crassisepta* Brady,1882
Faeroe Channel
- C. carinata* (Costa,1856)
= *Operculina carinata* Costa,1856
Drobak at 150-200m, Hardangerfjord, Bergen, Trondheim, Lofoten, Britain
- C. angigyrus* (Reuss,18??)
off W Scotland

- Meandrospirinae** Saidova,1981
- Meandrospira** Loeblich & Tappan,1946 (TM *M. washitensis* Loeblich & Tappan,1946)
= *Streblospira* Crespin & Belford,1957 (TM *S. meandrina* Crespin & Belford,1957)
= *Citella* Premoli Silva,1964 (TM *C. Julia* Premoli Silva,1964)
- Hemigordiopsidae** A. Nikitina,1969
- Hemigordiopsinae** A. Nikitina,1969
- Gordiospira** Heron-Allen & Earland,1932 (TM *G. fragilis* Heron-Allen & Earland,1932)
- G. arctica** Cushman,1933
Spitsbergen, NE, N & NW Greenland
- Fischerinidae** Millett,1898
- Fischerininae** Millett,1898
- Planispirina** Seguenza,1880
- P. communis** Seguenza,1880 (TM)
off Faeroes at 170 fathoms
- P. clairensis** Heron-Allen & Earland,1913
W Ireland
- Zoyaellinae** Saidova,1981
- Zoyaella** Loeblich & Tappan,1962 (TM *Ceratina trochamminoides* Goës,1894)
= *Ceratina* Goës,1894, non Latreille,1802, nec Menge,1868
- Glomulininae** Saidova,1981
- Glomulina** Rhumbler,1936
- G. fistulescens** Rhumbler,1936 (TM)
Baltic
- Nodobaculariellinae** Bogdanovich,1981
- Vertebralina** d'Orbigny,1826
- V. striata** d'Orbigny,1826 (TM)
Mediterranean, Shetlands
- Nubeculariidae** Jones,1875
- Nodobaculariinae** Cushman,1927
- Nodobacularia** Rhumbler,1895
- N. tibia** (Jones & Parker,1860)
= *Nubecularia tibia* Jones & Parker,1860 (TM)
NE Scotland, S England
- Nubeculinellinae** Avnimelech & Reiss,1954
- Webbina** d'Orbigny,1839 (TM *W. rugosa* d'Orbigny,1839)
- Nubecularinae** Jones,1875
- Gymnesina** Colom,1959 (TM *G. glomerosa* Colom,1959)
- Nubecularia** Defrance,1825
- N. lucifuga** Defrance,1825 (TM)
British region, e.g. Sussex
- N. inflata** Terquem,1876, non Brady,1884
France
- Ophthalmididae** Wiesner,1920
- Cornuloculina** Burbach,1886
- C. inconstans** (Brady,1879)
= *Hauerina inconstans* Brady,1879 (TM)
= *Hauerinella inconstans* : Auctt.
= *Ophthalmidium inconstans* : Auctt.
= *Ophthalmidium tumidulum* Brady,1884
Oslofjord, around Britain
- C. balkwilli** (Macfadyen,19??)
Britain
- C. carinatum** (Balkwill & Wright,1885)
= *Ophthalmidium carinatum* Balkwill & Wright,1885
* Correct genus???
around Britain
- Ophthalmina** Rhumbler,1936
- O. kilianensis** Rhumbler,1936 (TM)
* Om red algae & stones
W Baltic, N Germany, Belt Sea
- Spirothalmidium** Cushman,1927 (TM *Spiroloculina acutimargo* Brady,1884)
= *Spirothalmidium* : Cushman,1949
- S. acutimargo** (Brady,1884)
off W Scotland
- S. acutimargo** (Brady,1884) var. *emaciatum* Haynes,1973
Netherlands, Belgium, Wales, Ireland
- S. pusillum** (Earland,1934)
= *Spiroloculina pusilla* Earland,1934
Engl. Channel
- Discospirinidae** Wiesner,1931
- Discospirina** Munier-Chalmas,1902
- D. italica** (Costa,1856)
= *Pavonina italica* Costa,1856
= *Orbitolites tenuissima* W.B. Carpenter,1870 (TM)
off Britain, Mediterranean
- Miliolacea** Ehrenberg,1839
- Spiroloculinidae** Wiesner,1920
- Adelosina** d'Orbigny,1826 (TM *A. laevigata* d'Orbigny,1826)
= *Serpula (Retorta)* Walker & Boys,1784 (nomen rejectum, ICZN Op. 558)
= *Pollontes* Montfort,1808 (nomen rejectum, ICZN Op. 692) (TM *Pollontes vesicularis* de Montfort,1808)
- Inaequalina** Luczkowska,1971 (TM *Spiroloculina inaequalateralis* Schlumberger,1893)
= *Heterospiroloculina* McCulloch,1977 (TM *H. bikiniensis* McCulloch,1977)
- Nummulopyrga** Hofker,1983
= *Pseudopyrgo* Hofker,1976, non Rasheed,1971
- N. globulus** (Hofker,1976)
= *Pseudopyrgo globulus* Hofker,1976 (TM)
Biscay
- Planispirinoides** Parr,1950
? = *Pseudiopirilina* Saidova,1975 (TM *Pseudospirilina abyssalica* Saidova,1975)
- P. bucculenta** (Brady,1884)
= *Miliolina bucculenta* Brady,1884 (TM)
W Scotland, Barents Sea, Arctic Ocean
- Spiroloculina** d'Orbigny,1826
- S. depressa** d'Orbigny,1826 (TM)
Wales, NW Scotland, Engl. Channel, Galicia, Mediterranean
- S. excavata** d'Orbigny,1846
* clinging to firm substrates.
Engl. Channel, Arcachon, Biscay
- S. rotunda** d'Orbigny,1826
* clinging to firm substrates.
Engl. Channel
- S. henbesti** Petri,19?? : Gabel,1971
North Sea
- S. grataloupi** d'Orbigny,1826
? North Sea, Iceland, tropical Seas
- S. subimpressa** Parr,19?? : Gabel,1971
North Sea
- S. planulata** (Lamarck,1805)
= *Miliolites planulata* Lamarck,1805
around Britain, Mediterranean
- S. dorsata** Reuss,18??
Britain
- S. limbata** d'Orbigny,1826
Norway, Faeroes, Iceland, Britain, Mediterranean
- Hauerinidae** Schwager,1876
- Siphonapertinae** Saidova,1975
- Siphonaperta** Vella,1957 (TM *S. macbeathi* Vella,1957)
- Hauerininae** Schwager,1876
- Cycloforina** Luczkowska,1972
- C. contorta** (d'Orbigny,1846)
= *Quinqueloculina contorta* d'Orbigny,1846 (TM)
Bohusl., Norway (S of Lofoten), Iceland, off Scotland, off Ireland, France
- Hauerina** d'Orbigny,1839 (TM *H. compressa* d'Orbigny,1839)
- Lachlanella** Vella,1957 (TM *Quinqueloculina (Lachlanella) cooki* Vella,1957)
- Massilina** Schlumberger,1893
- M. secans** (d'Orbigny,1826)
= *Quinqueloculina secans* d'Orbigny,1826 (TM)
= *Miliolina secans* : Auctt.
* clinging to firm substrates. The commonest of all British Foraminifera, fide H.-A & E. Bohusl., Oslofjord, W? Norway, Belgium, W North Sea, Shetlands, Orkneys, Ireland, Scotland, Wales, S Britain, Engl. Channel, Arcachon, Mediterranean
- M. secans** (d'Orbigny,1826), var. *tenuistriata* Earland,1905
W Ireland, Sussex, Mediterranean
- M. carinata** (Fornasini,1905)
= *Spiroloculina niitida* d'Orbigny var. *carinata* Fornasini,1905
= *Spiroloculina milletti* Wiesner,1912
Wales, Japan, Argentina
- M. annexata** Schlumberger,1893
W Ireland, Mediterranean, Indo-Pacific
- Quinqueloculina** d'Orbigny,1826
= *Frumentarium* Fichtel & Moll,1798 (nomen rejectum, ICZN Op. 692)
= *Multiloculina* Abicht,1859 (TM *Serpula seminulum* Linnaeus,1758)
= *Trillina* Munier-Chalmas,1882 (TM *Triloculina strigillata* d'Orbigny,1850)
= *Rumanoloculina* Neagu,1984 (TM *Quinqueloculina robusta* Neagu,1968)
- Q. seminulum** (Linnaeus,1758)
= *Serpula seminulum* Linnaeus,1758 (TM)
* clinging to firm substrates.
Belt Sea, Oresund, Oslofjord, Bergen, Spitsbergen, Novaja Zemlya, Iceland, Faeroes, SE North Sea, Belgium, Shetlands, Orkneys, W Scotland, SW Ireland, Wales, Arcachon, Biscay, Galicia
- Q. aspera** d'Orbigny,1826
= *Q. agglutinata* : Auct., non Cushman,1917
Baltic, Oslofjord, Spitsbergen, NE & W Greenland, ?Zuideree, W Ireland, Wales, W Mediterranean
- Q. lamarckiana** d'Orbigny, in de la Sagra,1839
= *Q. auberiana* d'Orbigny, in de la Sagra,1839
Bergen, Britain
- Q. auberiana** d'Orbigny,1839, var. A : Haynes,1973
= *Q. vulgaris* : Colom,1952, non d'Orbigny,1826
= *Q. cuvieriana* : Brady,18??, non d'Orbigny,1839
Wales
- Q. auberiana** d'Orbigny,1839, var. B : Haynes,1973
Wales
- Q. arctica** Cushman,1933
N North Sea, Spitsbergen, NE & W Greenland

- Q. intricata*** Terquem,1878
 = *Q. semilunaris* Terquem,1880
 NW Scotland, Wales, Engl. Channel, Mediterranean
- Q. mediterranensis*** Le Calvez & Le Calvez,1958
 = *Miliolina bicornis* Sidebottom,1904, *non* (Walker & Jacob,1798)
 Wales, Mediterranean
- Q. stalkeri*** Loeblich & Tappan,1953
 = *Q. fusca* Cushman,1948, *non* Brady,1870
 Oslofjord, Spitsbergen, NE & W Greenland
- Q. bicornis*** (Walker & Jacob, in Kammacher,1798) var. *angulata* (Williamson,1858)
 = *Miliolina ferrusacii* (d'Orbigny,18??) : Heron-Allen & Earland,1916
 * clinging to firm substrates.
 Oslofjord, Belgium, Shetlands, NW Scotland, Wales, Celtic Sea, Engl. Channel, Arcachon, N Spain, W Mediterranean
- Q. clairensis*** (Heron-Allen & Earland,1930)
 = *Miliolina clairensis* Heron-Allen & Earland,1930
 * clinging to firm substrates.
 Belgium, W Scotland, Ireland, Bristol Channel, Engl. Channel, Arcachon, Mediterranean
- Q. clairensis*** (Heron-Allen & Earland,1930) var. A : Haynes,1973
 Wales
- Q. dimidiata*** Terquem,1876
 Engl. Channel
- Q. lata*** Terquem,1876
 * clinging to firm substrates.
 Belgium, NE Scotland, Engl. Channel, Wales, Arcachon
- Q. oblonga*** (Montagu,1803)
 = *Vermiculum oblongum* Montagu,1803
 = *Triloculina oblonga* : d'Orbigny,1826
 = *Miliolinella oblonga* : Murray,1965
 Bohusl., Drobak, Bergen, Lofoten, Spitsbergen, Iceland, W Scotland, Ireland, Engl. Channel, N Spain
- Q. candeiaria*** d'Orbigny, in de la Sagra,1839
 Helgoland, Cuba
- Q. cf. duthiersi*** (Schlumberger,1886)
 Wales
- Q. cf. rugosa*** d'Orbigny,1826
 Wales
- Q. lucida*** (Karrer,18??) : Gabel,1971
 North Sea
- Q. amygdaloidea*** (Brady,18??) : Gabel,1971
 North Sea
- Q. quadrata*** Norvang,1945
 = *Q. scleratica* : Cushman,1929, *non* Karrer,1868
 Iceland, around Britain
- Q. disparilis*** d'Orbigny,1826
 off Plymouth
- Q. seminuda*** Reuss,18??
 off W Scotland, W Ireland, Sussex
- Q. pygmaea*** Reuss,18??
 W Ireland
- Q. pulchella*** d'Orbigny,1826
 = *Miliolina pulchella* : Auctt.
 Bohusl., ?Norway, ?Britain, Iceland, Ireland
- Q. nitida*** Nørvang,1945
 Iceland
- Q. sp. aff. Q. stelligera*** Schlumberger,1893 : Gabel,1971
 North Sea
- Q. vulgaris*** d'Orbigny,1826
 Iceland, E Atlantic, Mediterranean
- Q. hauerina*** d'Orbigny,1846
 Iceland
- Q. implexa*** Terquem & Terquem,1886
 Iceland
- Q. proxima*** Terquem, ?in Terquem & Terquem,1886
 Iceland
- Q. sp. cf. Q. vulgaris*** d'Orbigny,1826 : Gabel,1971
 North Sea
- Q. sp. : Gabel,1971***
 Dogger Bank
- Miliolinellinae*** Vella,1957
- Affinetrina*** Luczkowska,1972 (TM *Triloculina planciana* d'Orbigny,1839)
- Biloculinella*** Wiesner,1931 (TM *Biloculina labiata* Schlumberger,1891)
- B. inflata*** (Wright,1902)
 = *Biloculina inflata* Wright,1902
 = *Pyrgo inflata* : Auctt.
 Bergen, Iceland, Britain
- B. sp. : Gabel,1971***
 North Sea
- Cruciloculina*** d'Orbigny,1839 (TM *C. triangularis* d'Orbigny,1839)
- Miliolinella*** Wiesner,1931
 = *Renoidea* Brown,1827 (TM *R. glabra* Brown,1827) (Suppressed, ICBN Op.1089)
 = *Pippinoidea* McCulloch,1977 (TM *Pippinoidea perplexa* McCulloch,1977)
 = *Neophthalmina* McCulloch,1977 (TM *Neophthalmina oregonensis* McCulloch,1977)
- M. subrotunda*** (Montagu,1803)
 = *Vermiculum subrotundum* Montagu,1803 (TM)
- = *Quinqueloculina subrotunda* (Montagu) *formahaauerinoides* Rhumbler,1936 (TM of *Pateoris* Loeblich & Tappan,1953)
 Poland, Kieler Bucht, Oslofjord, Bergen, Lofoten, Tromsø, Iceland, Spitsbergen, Belgium, W North Sea, Shetlands, W Scotland, SW Ireland, Wales, Engl. Channel, Arcachon, NE & W Greenland
- M. chukchiensis*** Loeblich & Tappan,1953
 N Greenland
- M. circularis*** (Bornemann,1855)
 = *Triloculina circularis* Bornemann,1855
 temperate, subtropical & tropical Atlantic, Faeroes, Iceland, Barents Sea, W Scotland, Ireland, England
- M. circularis*** (Bornemann,1855) var. *elongata* Kruit,1955
 Engl. Channel, Arcachon
- M. circularis*** (Bornemann,1855) var. *sublineata* Brady,18??
 W Ireland
- M. meridionalis*** (d'Orbigny,18??) : Gabel,1971
 North Sea
- M. pratense*** Meyer-Abich,19?? : Gabel,1971
 North Sea
- M. sp. : Gabel,1971***
 Braune Bank (North Sea)
- Pseudotriloculina*** Cherif,1970 (TM *Triloculina lecalvezae* Kaaschieter,1961)
- Pyrgo*** Defrance,1824 (TM *Pyrgo laevis* Defrance,1824)
 = *Biloculina* d'Orbigny,1826 (TM *Biloculina bulboides* d'Orbigny,1826)
 = *Pseudobiloculina* Cherif,1970 (TM *Biloculina oblonga* d'Orbigny,1839)
 = *Pyrgoides* Hofker,1976 (TM *Biloculina ringens* (de Lamarck) var. *denticulata* Brady,1884)
- P. bulboides*** (d'Orbigny,1826)
 = *Biloculina bulboides* d'Orbigny,1826
 Franz Josef Land
- P. ringens*** (de Lamarck,1804)
 = *Miltolites ringens* de Lamarck,1804
 Novaja Zemja, ?Iceland
- P. carinata*** (d'Orbigny,1839)
 = *Biloculina carinata* d'Orbigny,1839
 Belgium, NW Scotland, Wales, Cuba
- P. williamsoni*** (Silvestri,1923)
 = *Biloculina williamseni* Silvestri,1923
 = *Pyrgo elongata* : Cushman,1948, *non* (d'Orbigny,1826)
 Oslofjord, Spitsbergen, Engl. Channel, Wales, Iceland, NE & W Greenland
- P. depressa*** (d'Orbigny,1826)
 = *Biloculina depressa* d'Orbigny,1826
 Bergen, Belgium, Engl. Channel, Arcachon, Biscay, Galicia
- P. comata*** (Brady,1884)
 = *Biloculina comata* Brady,1884
 Gullmarfjord, Koster Channel, Oslofjord, Bergen, Faeroe Channel, Iceland
- P. elongata*** (d'Orbigny,1826)
 = *Biloculina elongata* d'Orbigny,1826
 Bergen, Barents Sea, Arctic Seas, Mediterranean
- P. cf. constricta*** Costa,1856
 Wales
- P. sp. A : Haynes,1973***
 Wales
- P. sp. B : Haynes,1973***
 Wales
- P. murryna*** (Schwager,1866) : Gabel,1971
 = *Biloculina murrayana* Schwager,1866
 North Sea, all deep ocean basins
- P. serrata*** (Bailey,1862)
 = *Biloculina serrata* Bailey,1862
 all deep ocean basins
- P. laevis*** Defrance,1824 : Gabel,1971
 North Sea, off Trondheimsfjord & off N Norway, Barents Sea
- P. lucernula*** (Schwager,1866)
 = *Biloculina lucernula* Schwager,1866
 ?North Sea, Iceland
- P. appendiculata*** (Reuss,1863) : Gabel,1971
 = *Biloculina appendiculata* Reuss,1863
 North Sea
- P. subglobulus*** Parr,19?? : Gabel,1971
 North Sea
- P. oblonga*** (d'Orbigny,1839) : Gabel,1971
 = *Biloculina oblonga* d'Orbigny,1839
 North Sea, Cuba
- P. anomala*** (Schlumberger,18??)
 Iceland, W Mediterranean
- P. vespertilio*** (Schlumberger,18??)
 Iceland, N Atlantic deep water
- P. inornata*** (d'Orbigny,1846)
 = *Biloculina inornata* d'Orbigny,1846
 Iceland
- P. sp. : Gabel,1971***
 E England
- Pyrgoella*** Cushman & E.M. White,1936
- P. sphaera*** (d'Orbigny,1839)
 = *Biloculina sphaera* d'Orbigny,1839 (TM)
 Oslofjord, Bergen, Gulf of Mexico
- P. globula*** (Bornemann,1855) : Gabel,1971
 = *Biloculina globulus* Bornemann,1855
 North Sea
- Triloculina*** d'Orbigny,1826
 = *Spidestomella* O.G. Costa,1856 (TM *S. globulifera* Costa,1856)

- T. trigonula** (de Lamarck,1804)
= *Mitolites trigonula* de Lamarck,1804 (TM)
Oslofjord-Vadsø, Iceland, NW Scotland, NW Ireland, Wales, Engl. Channel
- T. trihedra** Loeblich & Tappan,1953
Oslofjord, Wales, N & NW Greenland
- T. tricarinata** d'Orbigny,1826
Bergen, Iceland, Arctic Ocean
- T. agglutinans** (d'Orbigny,1839)
= *Quinqueloculina agglutinans* d'Orbigny, in de la Sagra,1839
= *Mitolina agglutinans* : Brady,1881
* Correct genus?
Novaja Zemlja, North Sea
- T. rotunda** d'Orbigny,1826 : Gabel,1971
North Sea
- T. anconensis** Schultze,1854
* Correct genus?
W Scotland, Adriatic
- T. brogniartii** d'Orbigny,1826
= *Mitolina brogniartii* : Auctt.
* Correct genus?
around Britain
- T. valvaria** Reuss,18??
Ostfjord, Lofoten, Faeroe Channel, W Ireland, Iceland
- T. suborbicularis** d'Orbigny,1826
= *Mitolina suborbicularis* : Auctt.
off W Scotland
- T. parvula** (Schlumberger,18???)
= *Mitolina parvula* (Schlumberger,18???)
* Correct genus???
Barents Sea
- T. pyriformis** (Schlumberger,18???)
= *Mitolina pyriformis* (Schlumberger,18???)
* Correct genus???
Barents Sea
- T. brevidentata** (Cushman,19???)
= *Mitolina brevidentata* (Cushman,19???)
* Correct genus???
Barents Sea
- Triloculinella** Riccio,1950 (TM *Triloculinella obliquinodus* Riccio,1950)
= *Scutoloris* Loeblich & Tappan,1953 (TM *Scutoloris tegminis* Loeblich & Tappan,1953)
- T. tegminis** (Loeblich & Tappan,1953)
Scutoloris tegminis Loeblich & Tappan,1953
NE & W Greenland
- T. sp. A** (Haynes,1973)
= *Scutoloris* sp. A Haynes,1973
Wales
- Triloculinoides** Shchedrina,1964 (TM *Triloculinoides magnus* Shchedrina,1964)
- Varidentella** Luczkowska,1972 (TM *Mitolina reussi* Bogdanovich,1947)
* Atlantic?
- Sigmoilinitinae** Luczkowska,1974
- Nummoloculina** Steinmann,1881
- N. contraria** (d'Orbigny,1846)
= *Biloculina contraria* d'Orbigny,1846 (TM)
Faeroe Channel, off Ireland, Mediterranean
- N. irregularis** (d'Orbigny,1839)
= *Biloculina irregularis* d'Orbigny,1839
Oslofjord
- Sigmoihauerina** S.Y. Zheng,1979 (TM *Hauerina bradyi* Cushman,1917)
- Sigmoilina** Schlumberger,1887
- S. sigmaidea** (Brady,1884)
= *Planispirina sigmaidea* Brady,1884 (TM)
off Britain
- Sigmoilinita** Seigle,1965 (TM *Quinqueloculina tenuis* Czjzek,1848)
- S. asperula** (Karrer,1868)
= *Spiroloculina asperula* Karrer,1868 (TM of *Spiroglutina* Mikhalevich,1983)
= *Ammonassimilina asperula* (Karrer,1868) : Gabel,1971
Faeroe Channel
- Sigmoilopsisinae** Vella,1957
- Sigmoilopsis** Finlay,1947
- S. schlumbergeri** (Silvestri,1904)
= *Sigmoilina schlumbergeri* Silvestri,1904 (TM)
Bergen, off SW Ireland
- S. moyi** Atkinson,1968
= *S. woodi* Atkinson,1968
= *S. schlumbergeri* : Atkinson,1968, non (Silvestri,1904)
= *S. wanganuiensis* : Atkinson,1968, non (Vella,1???)
Wales
- Tubinellinae** Rhumbler,1906
- Articulina** d'Orbigny,1826 (TM *Articulina nitida* d'Orbigny,1826)
= *Ceratospirulina* Ehrenberg,1858 (TM *C. sprattii* Ehrenberg,1858)
= *Ceratoloculina* Ehrenberg,1857 (Not available)
- A. sulcata** Reuss,1850
Atlantic, Mediterranean
- Parrina** Cushman,1931
= *Silvestria* Schubert,1921, non Verhoeff,1895, nec Brian,1902
- P. bradyi** Millett,1898
= *Nubecularia bradyi* Millett,1898 (TM)
= *Nubecularia inflata* Brady,1884, non Terquem,1876
- England, Mediterranean, Indo-Pacific
- Tubinella** Rhumbler,1906 (TM *Articulina finalis* var. *inornata* Brady,1884)
- Riveroinidae** Saidova,1981
- Pseudohauerina** Ponder,1972 (TM *Hauerina occidentalis* Cushman,1946)
- Alveolinacea** Ehrenberg,1839
- Alveolinidae** Ehrenberg,1839
- Borelis** de Montfort,1808 (TM *Borelis melonoidea* de Montfort,1808 = *Nautilus melo* Fichtel & Moll,1798)
= *Clausulus* de Montfort (TM *Clausulus indicator* de Montfort,1808 = *Nautilus melo* Fichtel & Moll,1798)
= *Melonites* de Lamarck,1812 (TM *Melonites sphaerica* de Lamarck,1812 = *Nautilus melo* Fichtel & Moll,1798)
= *Melonia* de Lamarck,1822 (TM *Melonites sphaerica* de Lamarck,1812 = *Nautilus melo* Fichtel & Moll,1798)
= *Neodalveolina* A. Silvestri,1928 (TM *A. bradyi* A. Silvestri,1927 = *Nautilus melo* Fichtel & Moll,1798)
- Soritacea** Ehrenberg,1839
- Peneropidae** Schultze,1854
- Dendritina** d'Orbigny,1826 (TM *Dendritina arbucula* d'Orbigny,1826)
= *Peloris* de Montfort,1808 (nom suppress. ICZN ZN(S) 2225/4 petition pending) (TM *Nautilus ambiguus* Fichtel & Moll,1798)
- Laevipeneroplis** Sulc,1936 (TM *Peneroplis karreri* Wiesner,1923 (nomen superfl.) = *Peneroplis planatus* (Fichtel & Moll,1798), var. *laevigatus* Karrer,1868, non *Peneroplis laevigata* Fornasini,1904)
= *Puteolus* Hofker,1950, non Montersato,1888 (*Mollusca*) (TM *Peneroplis proteus* d'Orbigny,1839)
= *Puteolina* Hofker,1952 (TM *Peneroplis proteus* d'Orbigny,1839)
= *Palaeopeneroplis* Hofker,1964 (TM *Palaeopeneroplis inornatus* Hofker,1964)
- Peneroplis** de Montfort,1808
- P. planatus** (Fichtel & Moll,1798)
= *Nautilus planatus* Fichtel & Moll,1798 (TM)
= *Cristellaria squamulosa* de Lamarck,1822 (TM of *Cristellaria* de Lamarck,1816)
Mediterranean, ?Shetlands
- P. hauerinus** (d'Orbigny,1846)
= *Cristellaria hauerina* d'Orbigny,1846
* Correct genus??
Plymouth
- P. gibba** (d'Orbigny,1839)
= *Cristellaria gibba* d'Orbigny,1839
= *Lenticulina gibba* : Gabel,1971
* Correct genus??
warm Faeroe Channel, W Scotland, W Ireland, Plymouth
- P. orbicularis** (d'Orbigny,1826)
= *Robulina orbicularis* d'Orbigny,1826
= *Cristellaria orbicularis* : Auctt.
= *Lenticulina orbicularis* : Gabel,1971
* Correct genus??
Plymouth
- P. albatrossi** (Cushman,1923)
= *Cristellaria albatrossi* Cushman,1923
= *Marginulina litus* : Parker & Jones,1865, non ?,!???
* Correct genus??
Norfolk, NE USA
- P. occidentalis** (Cushman,1923?), var. **glabrata** (Cushman,1923)
= *Cristellaria occidentalis* (Cushman,1923?), var. *glabrata* (Cushman,1923)
* Correct genus??
Iceland
- Soritidae** Ehrenberg,1839
- Archaiasinae** Cushman,1927
- Archaias** de Montfort,1808 (TM *Archais spirans* de Montfort,1808 = *Nautilus angulatus* Fichtel & Moll,1798)
= *Helenis* de Montfort,1808 (TM *Helenis spatulosus* de Montfort,1808 = *Nautilus aduncus* Fichtel & Moll,1798 = *Archais angulatus* (Fichtel & Moll,1798))
= *Iloetes* de Montfort,1808 (TM *Iloetes rotalitatus* de Montfort,1808 = *Nautilus orbiculus* Fichtel & Moll,1798, non Forskål,1775 = *Archais angulatus* (Fichtel & Moll,1798))
= *Orbiculina* de Lamarck,1816 (TM *Orbiculina nummularis* de Lamarck,1822 = *Nautilus orbiculos* Fichtel & Moll,1798)
- Cyclopoteolina** Seigle & Grove,1977 (TM *Peneroplis pertusus* (Forskål) var. *discoideus* Flint,1899)
- Soritinae** Ehrenberg,1839
- Amphisorus** Ehrenberg,1839 (TM *Nautilus orbiculos* Forskål,1775) (ICZN petition pending)
= *Taramellina* Munier-Chalmas,1902 (TM *Sorites dominicensis* Ehrenberg,1839)
- Sorites** Ehrenberg,1839 (TM *Nautilus orbiculos* Forskål,1775) (ICZN petition pending)
= *Taramellina* Munier-Chalmas,1902 (TM *Sorites dominicensis* Ehrenberg,1839)
- SILICOLOCULININA** Resig, Lowenstam, Echols & Weiner,1980
* Not represented in the area
- LAGENINA** Delage & Hérouard,1896
- Nodosariacea** Ehrenberg,1838
- Nodosariidae** Ehrenberg,1838
- Nodosariinae** Ehrenberg,1838
- Chrysalogonium** Schubert,1908 (TM *Nodosaria polystoma* Schwager,1866)
- Dentalina** Risso,1826 (TM *Nodosaria* (les *Dentalines*) *cuvieri* d'Orbigny,1826)
- D. subarcuata** (Montagu,1803)
= *Nautilus subarcuatus* Montagu,1803
Engl. Channel
- D. drammenensis** Feyling-Hanssen,1964
Oslofjord
- D. trondheimensis** Feyling-Hanssen,1964
= *Nodosaria communis* : Heron-Allen & Earland,1916, non d'Orbigny,1826
= *Dentalina subarcuata* : Williamson,1858, non Montagu,1803
Trondheim, ?Wales
- D. baggi** Galloway & Wissler,1927
= *Nodosaria pauperata* Baggi,1912, non *Dentalina pauperata* d'Orbigny,1846
= *Nodosaria callomorpha* Baggi,1912, non Reuss,1866
N & NW Greenland
- D. frobisherensis** Loeblich & Tappan,1953

- Spitsbergen, NE & W Greenland
D. ittai Loeblich & Tappan, 1953
 Spitsbergen, NE & W Greenland
- D. metvillensis* Loeblich & Tappan, 1953
 N Greenland
- D. obliqua* (Linnaeus, 1767)
 = *Nautilus obliquus* Linnaeus, 1767
 Gullmarfjord, Skagerrak, Iceland, Engl. Channel, Barents Sea
- D. pauperata* d'Orbigny, 1846
 Stavanger, Christiansand, Vardo, Iceland, NE, N & W Greenland, Barents Sea, Franz Josef's Land, around Britain
- D. mucronata* Neugeboren, 1856
 off W Scotland, off SW Ireland, Franz Josef's Land
- D. communis* (d'Orbigny, 1826)
 = *Nodosaria communis* d'Orbigny, 1826
 ?Bergen, Iceland, Engl. Channel
- D. consobrina* d'Orbigny, 1846
 = *Silostomella consobrina* (d'Orbigny, 1846) : Gabel, 1971
 * Correct genus??
 W Scotland, off SW Ireland
- D. consobrina* d'Orbigny, var. *emaciata* Reuss, 18??
 * Correct genus??
 ?warm Faeroe Channel, ?off SW Ireland
- D. subsoluta* (Cushman, 1923)
 = *Nodosaria subsoluta* Cushman, 1923
 = *Nodosaria soluta* : Brady, 1884, non *Dentalina soluta* Reuss, 1851
 off Bergen, Lofoten, Vadsø, warm Faeroe Channel, W Ireland, off SW Ireland, Iceland
- D. advena* (Cushman, 1923)
 = *Nodosaria advena* Cushman, 1923
 ?cold Faeroe Channel, off W Scotland
- D. inornata* d'Orbigny, var. *bradyensis* (Dervieux, 19??) : Gabel, 1971
 North Sea
- D. vertebralis* : Terquem & Terquem, 1886, hon (Batsch, 1???)
 Iceland
- D. sp. : Gabel, 1971
 German Bight
- Enantiodentalina* Marie, 1941 (TM *Enantiodentalina communis* Marie, 1941)
 = *Paradentalina* Uchio, 1960 (TM *Enantiodentalina muraii* Uchio, 1953)
- Grigelis* Mikhalevich, 1981
- G. monile* (Soldani, 1798)
 = *Orthoceras monile* Soldani, 1798
 = *Nodosaria pyrula* d'Orbigny, 1826 (TM)
 ?= *Dentalina guttifera* : Terquem & Terquem, 1886, non d'Orbigny, 1846
 Engl. Channel, ?Iceland
- Laevidentalina* Loeblich & Tappan, 1986 (TM *Laevidentalina aphelis* Loeblich & Tappan, 1986)
- Nodosaria* de Lamarck, 1812
 = *Glandulonodosaria* A. Silvestri, 1900 (TM *Nodosaria ambigua* Neugeboren, 1856)
- N. radicula* (Linnaeus, 1758)
 = *Nautilus radicula* Linnaeus, 1758 (TM)
 off Britain, Arctic Ocean
- N. calliomorpha* Reuss, 1866
 Engl. Channel, off SW Ireland
- N. lamnulifera* Boomgaart, 19?? : Gabel, 1971
 North Sea
- N. lagenifera* Neugeboren, 18?? : Gabel, 1971
 North Sea
- N. simplex* Silvestri, 1872
 warm Faeroe Channel, ?W Ireland
- N. subannulata* Cushman, 1923
 ?Nordland, off NE USA, Barents Sea
- N. farcimen* Soldani, 17??
 off Ireland
- N. filiformis* d'Orbigny, 1826
 Stavanger, Lofoten, Iceland, Britain, off Canaries
- N. aequalis* (Reuss, 1870)
 = *Glandulina laevigata* (d'Orbigny, 1826), var. *aqualis* Reuss, 1870
 Halland, off Norway, Faeroe Channel, Spitsbergen
- N. proxima* Silvestri, 1???
 W Scotland, S Cornwall
- N. raphanistrum* (Linnaeus, 1758)
 = *Nautilus raphanistrum* Linnaeus, 1758
 off W Scotland
- N. hispida* d'Orbigny, 18??
 Ireland, Isle of Man
- N. flintii* Cushman, 1923
 = *Nodosaria obliqua* : Brady, 1884 (p.p.) ; Goës, 1894 (p.p.), non (Linnaeus, 1758)
 ?around Britain, Caribbean
- N. plebeia* Reuss, 18??
 W Ireland
- N. setosa* Schwager, 18??
 Burwood Bank
- N. sp. : Gabel, 1971
 central North Sea
- Pandaglandulina* Loeblich & Tappan, 1955 (TM *Pandaglandulina dinopoli* Loeblich & Tappan, 1955)
- Pseudonodosaria* Boomgaart, 1949 (TM *Glandulina discreta* Reuss, 1850)
 = *Rectoglandulina* Loeblich & Tappan, 1955 (TM *Rectoglandulina appressa* Loeblich & Tappan, 1955)
- P. rotundata* (Reuss, 1849)
 = *Glandulina rotundata* Reuss, 1849
 = *Nodosaria rotundata* : Auctt.
 Oslofjord, Britain
- P. torrida* (Cushman, 1923)
 = *Nodosaria (Glandulina) laevigata* d'Orbigny, var. *torrida* Cushman, 1923
 Oslofjord, Gulf of Mexico
- Pyramidulina* Fornasini, 1894 (TM *Pyramidulina eptagona* Fornasini, 1894 = *Nautilus raphanus* Linnaeus, 1758)
 = *Pseudoglandulina* Cushman, 1929 (TM *Nautilus comatus* Batsch, 1791)
 = *Nodosariella* Wedekind, 1937 (TM *Nautilus raphanus* Linnaeus, 1758)
- Svenia* Brozen, 1937 (TM *Nodosaria laevigata* Nilsson, 1826)
- Lingulinina* Loeblich & Tappan, 1961
- Lingulina* d'Orbigny, 1826
- L. carinata* d'Orbigny, 1826 (TM)
 Engl. Channel
- L. bilobuli* Wright, 1910-11
 = *Lingulina carinata* var. *biloculi* Wright, 1910-11
 Engl. Channel, W Scotland, W Ireland
- L. quadrata* Heron-Allen & Earland, 1913
 NE Scotland, W Ireland
- L. bicarinata* Sidebottom, 1907
 = *Lingulina carinata* d'Orbigny, 1826, var. *bicarinata* Sidebottom, 1907
 NE & W Scotland, W Ireland, Mediterranean
- L. armata* Sidebottom, 1907
 NE Scotland, Orkneys, W Ireland, Mediterranean
- L. pellucida* Sidebottom, 1907
 NE Scotland, W Ireland, S Cornwall, Mediterranean
- Frondiculariinae* Reuss, 1860
- Frondicularia* Defrance, in d'Orbigny, 1826 (TM *Renulina complanata* Defrance, in de Blainville, 1824)
- F. (?) *translucens* Heron-Allen & Earland, 1913
 Ireland
- F. (?) *sidebottomi* Cushman, 1923
 = *Frondicularia spatulata* Sidebottom, 1907, non Williamson, 18??
 North Sea, N & W Scotland, W Ireland, Mediterranean
- F. (?) *pygmaea* Sidebottom, 1907
 NE Scotland, W Ireland, Mediterranean
- F. (?) *spatulata* Williamson, 18??
 Britain
- F. (?) *tenera* (Bornemann, 18???)
 ?W Scotland
- F. *affinis* Terquem & Terquem, 1886
 Iceland
- Plectofrondiculariinae* Cushman, 1927
- Amphimorphina* Neugeboren, 1850 (TM *Amphimorphina haueriana* Neugeboren, 1850)
 = *Amphimorphinella* Keyzer, 1953 (TM *Amphimorphinella butonensis* Keyzer, 1953)
- Vaginulinidae* Reuss, 1860
- Lenticulininae* Chapman, Parr & Collins, 1934
- Cribrorobulina* Thalmann, 1947 (TM *Robulina serpens* Seguenza, 1880)
 = *Cribrorobulina* Selli, 1941 (name not available)
- Lenticulina* de Lamarck, 1804 (TM *Lenticulites rotulatus* de Lamarck, 1804)
 = *Lenticulina* de Lamarck, 1804 (junior syn.)
 = *Phonemus* de Montfort, 1808 (TM *Nautilus vortex* Fichtel & Moll, 1798)
 = *Antenor* de Montfort, 1808 (TM *Antenor diaphaneus* Montfort, 1808)
 = *Oreas* de Montfort, 1808, non Hübner, 1807 (TM *Oreas subulata* de Montfort, 1808 = *Nautilus acutauricularis* Fichtel & Moll, 1798)
 = *Robulus* de Montfort, 1808 (TM *Robulus cultratus* de Montfort, 1808)
 = *Patrocles* de Montfort, 1808 (TM *Patrocles querelans* de Montfort, 1808 = *Nautilus vortex* Fichtel & Moll, 1798)
 = *Clysiophantes* de Montfort, 1808 (TM *Clysiophantes calcar* de Montfort, 1808)
 = *Herion* de Montfort, 1808 (TM *Herion rostratus* de Montfort, 1808)
 = *Rhinocurus* de Montfort, 1808 (TM *Rhinocurus araneosus* de Montfort, 1808)
 = *Lampus* de Montfort, 1808, non Meuschen, 1787 (TM *Lampus tritheimus* de Montfort, 1808)
 = *Soldania* d'Orbigny, 1826 (TM *Soldania carinata* d'Orbigny, 1826)
 = *Cochlea* Zalessky, 1926, non Costa, 1778, nec Martyn, 1784, nec Hitchcock, 1888 (TM *Cochlea sapracolli* Zalessky, 1926)
 = *Darbyella* Howe & Wallace, 1932 (TM *Darbyella danvillensis* Howe & Wallace, 1932)
 = *Perisiphinctina* Wedekind, 1937 (TM *Robulina depauperata* Reuss, 1851)
 = *Darbyella* Harris & Sutherland, 1954 (TM *Darbyella hempraeensis* Harris & Sutherland, 1954)
 = *Robidzhonia* Melnikova & Suleymanov, in Suleymanov, 1969 (TM *Robidzhonia prima* Melnikova & Suleymanov, in Suleymanov, 1969, non *Cristellaria prima* d'Orbigny, 1850 (= *Lenticulina suleymanovi* Loeblich & Tappan, 1988))
- L. rotulata* (de Lamarck, 1804)
 = *Lenticulites rotulatus* de Lamarck, 1804
 = *Cristellaria calcar* : Williamson, 1858, non (Linnaeus, 1767)
 ?Bergen, Engl. Channel
- L. peregrina* (Schwager, 1866)
 = *Cristellaria peregrina* Schwager, 1866
 ?North Sea, W Scotland, off SW Ireland, off the western approaches to the Engl. Channel, off Galicia
- L. cf. angulata* (Reuss, 1851)
 = *Robulina angulata* Reuss, 1851
 Oslofjord
- L. cultrata* (de Montfort, 1808)
 = *Nautilus cultratus* de Montfort, 1808
 = *Cristellaria cultrata* : Auctt.
 Plymouth
- L. acutauriculus* (Fichtel & Moll, 1798)
 = *Nautilus acutauricularis* Fichtel & Moll, 1798
 = *Oreas subulatus* de Montfort, 1808
 ?W Scotland, W Ireland
- L. thalmani* (Hessland, 19??) : Gabel, 1971
 North Sea

- L. convergens** (Bornemann,18??) : Gabel,1971
 = *Cristellaria convergens* Bornemann,18??
 off W Scotland, Barents Sea
- L. vortex** (Fichtel & Moll,1798)
 = *Nautilus vortex* Fichtel & Moll,1798
 ?W Scotland, off Ireland, S Cornwall
- L. sp.** : Gabel,1971
 German Bight
- Marginulinopsis** A. Silvestri,1904 (TM *Cristellaria bradyi* Goës,1894 = *Marginulina costata* (Batsch) : Brady,1884 = *Marginulinopsis denticostata* Thalmann,1937)
- Neolenticulina** McCulloch,1977 (TM *Neolenticulina chathamensis* McCulloch,1977)
- Percutazonaria** Loeblich & Tappan,1986 (TM *Cristellaria subculeata* Cushman,1923)
 = *Gladioria* Thalmann,1941, non Wick,1939 (TM *Cristellaria decorata* Reuss,1855)
 = *Gladioria* Wedekind,1937 (name not available)
- P. subculeata** (Cushman,1923)
 = *Cristellaria subculeata* Cushman,1923
 = *Cristellaria aculeata* : Brady,1884, non d'Orbigny,18
 ?Engl. Channel, tropical W Atlantic
- Saracenaria** Defrance,1824
 = *Enanthocristellaria* Marie,1941 (TM *Cristellaria navicula* d'Orbigny,1840)
- S. italica** Defrance,1824 (TM)
 = *Cristellaria italica* : Auctt.
 off SW Ireland, Wales, S Cornwall
- Marginulininae** Wedekind,1937
- Amphicoryna** Schlumberger, in *Cochlidion alexandreae* Zalessky,1926
 = *Plesiocorine* Schlumberger, in Milne-Edwards,1882 (TM *Plesiocorine edwardsi* Schlumberger, in Milne-Edwards,1882)
 = *Lagenomodosaria* A. Silvestri,1900 (TM *Nodosaria scalaris* (Batsch,1791) var. *separans* Brady,1884)
 = *Nodosariopsis* A. Silvestri,1902, non Rzehak,1895 (TM *Marginulina falk* Jones & Parker,1860)
- A. scalaris** (Batsch,1791)
 = *Nautilus (Orthoceras) scalaris* Batsch,1791
 Denmark, Oslofjord, Bergen, Vesteraalen, Iceland, Belgium, ?S & ?W Britain, ?Arcachon, ?Biscay
- A. separans** (Brady,1884)
 = *Nodosaria scalaris* (Batsch,1791) var. *separans* Brady,1884
 off Britain, Indo-Pacific
- A. hirsuta** (d'Orbigny,18??) : Gabel,1971
 = *Nodosaria hirsuta* : Auctt.
 North Sea
- A. sp.** : Murray,1965
 central North Sea
- Astacolus** de Montfort,1808
 = *Cochlidion Zalesskyi*,1926 (TM *Cochlidion alexandreae* Zalessky,1926)
 = *Polymorphinoides* Cushman & Hanzawa,1936 (TM *Polymorphinoides spiralis* Cushman & Hanzawa,1936)
 = *Sacculariella* Wedekind,1937 (TM *Sacculariella ensis* Wedekind,1937) (not available)
 = *Gladioria* Wick,1939, non Thalmann,1941 (TM *Cristellaria hermanni* Andea,1898)
 = *Enantio vaginalinula* Marie,1941 (TM *Cristellaria recta* d'Orbigny,1840)
- A. crepidulus** (Fichtel & Moll,1798)
 = *Nautilus crepidula* Fichtel & Moll,1798
 = *Astacolus crepidulatus* de Montfort,1808 (TM)
 = *Crepidulina astacolus* de Blainville,1824 (TM of *Crepidulina* de Blainville,1824)
 = *Cristellaria crepidula* : Auctt.
 Engl. Channel, Greenland
- A. hyalaculus** Loeblich & Tappan,1953
 NW Greenland
- Hemirobulina** Stache,1864 (TM *Cristellaria (Hemirobulina) arcuata* Stache,1864 = *Cristellaria (Hemicristellaria) procura* Stache,1864)
 = *Hemicristellaria* Stache,1864
 = *Enantomarginulina* Marie,1941 (TM *Enantomarginulina dorbignyi* Marie,1941)
 = *Pseudodimorphina* McCulloch,1977 (TM *Pseudodimorphina galapagensis* McCulloch,1977)
- Marginulina** d'Orbigny,1826 (TM *Marginulina raphanus* d'Orbigny,1826, non *Nautilus raphanus* Linnaeus,1758)
 = *Ellipsomarginulina* A. Silvestri,1923 (TM *Marginulina raphanus* d'Orbigny,1826)
 = *Marginulinella* Wedekind,1937 (TM *Nautilus (Orthoceras) costatus* Batsch,1791)
- M. costata** (Batsch,1791)
 = *Nautilus (Orthoceras) costatus* Batsch,1791
 off Britain
- M. glabra** d'Orbigny,1826
 around Britain
- M. glabra** d'Orbigny,1826, var. *obesa* Cushman,1923
 off Britain
- Vaginulinopsis** Silvestri,1904 (TM *Vaginulina soluta* Silvestri var. *carinata* Silvestri,1898)
Vaginulininae Reuss,1860
- Planularia** Defrance, in de Blainville,1826, non Nilsson,1826 (TM *Peneroplis auris* Defrance, in de Blainville,1824)
- Vaginulina** d'Orbigny,1826 (TM *Nautilus legumen* Linnaeus,1758)
- V. legumen** (Linnaeus,1758)
 = *Nautilus legumen* Linnaeus,1758 (TM)
 = *Dentalina legumen* : Auctt.
 around Britain, Plymouth
- V. linearis** (Montagu,1803)
 = *Nautilus linearis* Montagu,1803
 between Trondheim & North Cape, Shetlands, Hebrides, Plymouth, France
- V. spinigera** Brady,1881
 off Norway, warm Faeroe Channel, off SW Ireland
- Spiralingulina** Loeblich & Tappan,1986
- Spirilingulina** Sellier de Civrieux & Dessauvagie,1965 (TM *Lingulina polymorpha* O.G. Costa,1861 = *Lingulomopsis carlofortensis* Bornemann,1883)
 * Atlantic?
- Lagenidae** Reuss,1862
- Hyalinonetria** Patterson & Richardson,1987 (TM *Hyalinonetria sahlense* Patterson & Richardson,1987)
 = *Amphorina* d'Orbigny,1849, non de Quatrefages,1844 (TM *Amphorina gracilis* O.G. Costa,1856 = *Amphorina costai* Andersen,1961 non = *Lagena gracilis* Williamson,1848)
- Lagena** Walker & Jacob, in Kammacher,1798, non Bolten,1798, nec Schumacher,1817, nec Mörch,1852, nec Kramer,1881 (TM *Serpula (Lagena) sulcata* Walker & Jacob, in Kammacher,1798)
 = *Lagenulina* Terquem,1876 (TM *Lagenulina sulcata* Terquem,1876)
 = *Capitellina* Marsson,1878 (TM *Capitellina multistriata* Marsson,1878)
- L. sulcata** (Walker & Jacob, in Kammacher,1798) var. *torquiformis* Haynes,1973
 Wales, Irish Sea, Engl. Channel
- L. clavata** (d'Orbigny,1846)
 = *Oolina clavata* d'Orbigny,1846
 Bohusl, Skagerrak, Stavanger, Spitsbergen, Barents Sea, Iceland, North Sea, Belgium, Engl. Channel, Bristol Channel, Wales, W Scotland, off Ireland, Arcachon, Arctic Seas
- L. doveensis** Haynes,1973
 Wales, Oslofjord (late quaternary)
- L. filicosta** Reuss,18??
 = *L. striata* : Williamson,1858, : Cushman,1923, non (d'Orbigny,1839)
 Oslofjord, Britain
- L. acuticosta** Reuss,1862
 = *Oolina acuticosta* (Reuss,1862) : Gabel,1971
 Bergen, Iceland, Barents Sea, Britain
- L. hibernica** Haynes,1973
 Wales, Ireland
- L. interrupta** Williamson,1848
 = *Lagena striata* (Montagu) var. *alpha*, *interrupta* Williamson,1848
 Celtic Sea, Bristol Channel, western approaches to the Engl. Channel
- L. flatulenta** Loeblich & Tappan,1953
 = *L. laevis* : Auct., non (Montagu,1803)
 N & NW Greenland
- L. gracillima** (Seguenza,1862)
 = *Amphorina gracillima* Seguenza,1862
 Oslofjord, Bergen, Vestfjord, Iceland, Spitsbergen, Novaja Zemlja, N & NW Greenland, Engl. Channel
- L. laevis** (Montagu,1803)
 = *Vermiculum laeve* Montagu,1803
 = *Lagena vulgaris* Williamson,1858
 * Nørvang,1945 use Walker & Boys,1784 as author for *Serpula laevis*)
 Skagerrak, Bergen, North Sea, Friesian Islands, Spitsbergen, Barents Sea, Arctic Ocean, Belgium, Shetlands, Iceland, Irish Sea, S Britain, Engl. Channel, Biscay
- L. semistriata** Williamson,1848
 = *Lagena striata* (Walker) var. *beta*, *semistriata* Williamson,1848
 Skagerrak, Barents Sea, Novaja Zemlja, North Sea, Celtic Sea, off SW Ireland, Engl. Channel, Arcachon
- L. mollis** Cushman,1944
 = *L. distoma* Parker & Jones,1864
 N & NW Greenland
- L. distoma** Parker & Jones MS, in Brady,1864
 Oslofjord, Bergen, Vadsø, Iceland, Spitsbergen, Novaja Zemlja, W Scotland, Ireland, England
- L. perlucida** (Montagu,1803)
 = *Vermiculum perlucidum* Montagu,1803 (TM of *Vermiculum* Montagu,1803)
 Iceland, Engl. Channel, Wales, Kent, Ireland
- L. semilineata** Wright,1886
 = *L. caudata* : Cushman,1948, non (d'Orbigny,1839)
 off Scotland, off Ireland, N & W Greenland
- L. setigera** Millett,1901
 = *L. clavata* (d'Orbigny) var. *setigera* Millett,1901
 Oslofjord, NE & NW Greenland
- L. striata** (d'Orbigny,1839)
 = *Oolina striata* d'Orbigny,1839
 Oslofjord, Bergen, Vadsø, Iceland, Engl. Channel, ?Wales
- L. substriata** Williamson,1848
 Oslofjord, German North Sea coast, Kent, Engl. Channel, Wales, Bristol Channel, Cornwall, Arcachon
- L. tenuis** (Bornemann,1855)
 = *Ovalina tenuis* Bornemann,1855
 off Plymouth
- L. pacifica** Sidebottom,1912
 = *L. spinosa* : Heron-Alen & Earland,1916, non Millett,18??
 Bergen, Faeroe Channel, Wales, W Scotland, Mediterranean
- L. spicata** Cushman & McCulloch,1950
 = *L. sulcata* (Walker & Jacob,1798) var. *spicata* Cushman & McCulloch,1950
 Wales
- L. iyellii** (Seguenza,1862)
 = *Amphorina iyellii* Seguenza,1862
 Engl. Channel, off W Ireland, Mediterranean
- L. hispida** Reuss,1858
 Oslofjord, around Britain
- L. nebulosa** Cushman,1923
 = *L. laevis* (Montagu) var. *nebulosa* Cushman,1923
 Oslofjord
- L. protea** Chaster,1892
 around Britain
- L. pulchella** Brady,1866, var. *hexagona* Heron-Alen & Earland,1916
 off W Scotland
- L. apiopleura** Loeblich & Tappan,1953
 = *Oolina apiopleura* : Gabel,1971

- North Sea, Spitsbergen, NW Greenland
L. tricincta Gumbel,1868
 * Correct genus?
 Novaja Zemlya
- L. ornata* (Williamson,1858)
 = *Entosolenia marginata* (Walker & Boys,1784), var. *ornata* Williamson,1858
 * Correct genus?
 Engl. Channel
- L. annectens* Burrows & Holland, in Jones, Parker & Brady.,1866
 * Correct genus?
 W Scotland, Engl. Channel
- L. falcata* Chaster,1892
 Engl. Channel, NE & W Scotland, W Ireland
- L. stewartii* J. Wright,1910-11
 Engl. Channel, W Ireland
- L. crenata* Parker & Jones,1865
 Engl. Channel, off Australia
- L. elongata* (Ehrenberg,1844)
 = *Miliola elongata* Ehrenberg,1844
 = *Lagena gracilima* : Goës,1894 (p.p. - plate 13, figs 728 & 730), non *Amphorina gracillima* Seguenza,1862
 off Britain, Spitsbergen, off Greenland
- L. stelligera* Brady,1881
 ?Faeroe Channel, Sussex
- L. aspera* Reuss,1861
 off Britain
- L. botelliformis* Brady,1881
 W Scotland, off SW Ireland
- L. curvilineata* Balkwill & Wright,1882
 Britain
- L. faba* Balkwill & Millett,1884
 W Scotland, Ireland
- L. fimbriata* Brady,1881
 around Britain
- L. forcicula* Heron-Allen & Earland,1913
 NE Scotland
- L. lacunata* Burrows & Holland,18??
 off W Scotland
- L. ovum* (Ehrenberg,1843)
 = *Miliola ovum* Ehrenberg,1843
 * Correct genus?
 W Scotland, Ireland
- L. quadricostulata* Reuss,1870
 Faeroe Channel, Ireland, Sussex
- L. reniformis* Sidebottom,1913
 off Scotland
- L. reticulata* (Macgillivray,1843)
 = *Lagena reticulata* Macgillivray,1843
 around Britain, Iceland
- L. trigonomarginata* Parker & Jones,1865
 Yorkshire, Ireland
- L. trispida* Reuss,18??
 Stavanger, Iceland
- L. tricarinata* Gramb,1???
 Bergen, Iceland
- L. lisabellae* (d'Orbigny,18??)
 Iceland
- L. squamosa-alata* Brady,1884
 Iceland, off Ireland
- L. inornata* d'Orbigny,18??
 Iceland, Falkland Islands
- L. implexa* Terquem & Terquem,1886
 Iceland
- L.* sp. : Haynes,1973
 Wales
- Procerolagena* Puri,1954
- P. gracilis* (Williamson,1848)
 = *Lagena gracilis* Williamson,1848 (TM)
 NW Scotland, off SW Ireland, Celtic Sea, Engl. Channel, Iceland, Barents Sea, Novaja Zemlya
- Pygmaeoestron* Patterson & Richardson,1987
- P. hispidula* (Cushman,1913)
 = *Lagena hispidula* Cushman,1913 (TM)
 North Sea
- Reussoolina* Colom,1956
- R. apiculata* (Reuss,1851)
 = *Oolina apiculata* Reuss,1851 (TM)
 = *Fissurina apiculata* : Auctt.
 Oslofjord, Novaja Zemlya, Engl. Channel
- Polymorphinidae* d'Orbigny,1839
- Polymorphininae* d'Orbigny,1839
- Globulina* d'Orbigny,1839
- G. gibba* (d'Orbigny,1826)
 = *Polymorphina (les Globulines) gibba* d'Orbigny,1826 (TM)
 Belgium, Engl. Channel, Arcachon
- G. gibba* (d'Orbigny,1826) var. *myristiformis* (Williamson,1858)
 = *Polymorphina myristiformis* Williamson,1858
- Belgium, Engl. Channel, Arcachon, Mediterranean
G. cf. *inaequalis* Reuss,1850
 Wales
- G. amygdaloidea* Reuss,1851
 = *Polymorphina amygdaloidea* : Auctt.
 E Atlantic
- G. minuta* (Roemer,18??) : Gabel,1971
 North Sea
- G. spinosa* d'Orbigny,1846
 off Britain
- Guttulina* d'Orbigny,1839
 = *Sigmomorpha* Cushman & Ozawa,1928 (TM) *Sigmomorpha sadoensis* Cushman & Ozawa,1928
- G. communis* (d'Orbigny,1826)
 = *Polymorphina (les Guttulines) communis* d'Orbigny,1826 (TM)
 = *Guttulina problema* d'Orbigny,1826
 Plymouth, Iceland
- G. rotundata* Bornemann,1855
 = *Polymorphina rotundata* : Auctt.
 Plymouth
- G. (? Laryngosigma) harrisi* Haynes,1973
 NE Scotland, W North Sea, Ireland
- G. sororia* (Reuss,1863)
 = *Polymorphina (Guttulina) sororia* Reuss,1863
 * Correct genus?
 Plymouth
- G. sororia* (Reuss,1863), var. *cuspidata* Brady,1884
 * Correct genus?
 off W Ireland at 1477-2639 m.
- G. austriaca* d'Orbigny,18?? : Gabel,1971
 North Sea
- G. pacifica* (Cushman & Ozawa,1928?) : Gabel,1971
 North Sea
- Polymorphina* d'Orbigny,1826 (TM) *Polymorphina burdigalensis* d'Orbigny,1826
- P. compressa* d'Orbigny,1846
 Plymouth, Novaja Zemlya
- P. oblonga* d'Orbigny,1846
 Plymouth, Novaja Zemlya
- P. complexa* Sidebottom,1904
 = *Polymorphina (?) complexa* Sidebottom,1904
 * Correct genus?
 Plymouth
- P. ovata* d'Orbigny,1846 : Gabel,1971
 North Sea
- P. cylindroides* Roemer,1838
 off Britain
- P. angusta* Egger,1857
 Sussex
- P. complanata* d'Orbigny,1846
 around Britain
- P. regina* Brady, Parker & Jones,1870
 Sussex
- P. thouini* d'Orbigny,18??
 Wales
- P. acuta* (Roemer,1838)
 = *Globulina acuta* Roemer,1838
 * Correct genus?
 Faeroes, Spitsbergen
- Pseudopolymorphina* Cushman & Ozawa,1928 (TM) *Pseudopolymorphina hanzawai* Cushman & Ozawa,1928
- P. cf. novangliae* (Cushman,1923)
 Wales
- P. suboblonga* Cushman & Ozawa,1930
 Oslofjord, Wales
- P. ovalis* (Cushman & Ozawa,1928?) : Gabel,1971
 North Sea
- Pyrulina* d'Orbigny,1839 (TM) *Polymorphina (les Pyrulines) gutta* d'Orbigny,1826
 = *Pyrdinella* Cushman & Ozawa,1928 (TM) *Polymorphina lanceolata* Reuss,1851
- P. fusiformis* (Roemer,18??)
 = *Polymorphina fusiformis* Roemer,18??
 E Ireland
- P. lanceolata* (Reuss,1851)
 = *Polymorphina lanceolata* Reuss,1851
 E & W Atlantic
- Sigmomorphina* Cushman & Ozawa,1928 (TM) *Sigmomorpha (Sigmomorpha) yokoyamai* Cushman & Ozawa,1928
- S. undulosa* (Terquem,18??) : Gabel,1971
 North Sea
- S. semitecta* (Reuss,18??), var. *terquemiana* Cushman,19?? : Gabel,1971
 North Sea
- Webbinellinae* Rhumbler,1904
- Webbinella* Rhumbler,1904
 = *Lingulopyrulinoides* Saidova,1975 (TM) *Lingulopyrulinoides crassa* Saidova,1975
- W. hemisphaerica* (Jones, Parker & Brady,1866)
 = *Trochammina (Webbinella) irregularis* (d'Orbigny) var. *hemisphaerica* Jones, Parker & Brady,1866 (TM)

- = *Placopsisina bulla* : Goës,1894 (p.p. - plate 6, fig. 211 & 212), non Brady,1881
Koster Channel, Drobak, Engl. North Sea coast, W Ireland, Faeroe Channel
- Ramulininae** Brady,1884
- Ramulina** T.R. Jones, in J. Wright,1875 (TM) *Ramulina laevis* T.R. Jones, in J. Wright,1875
- R. globulifera** Brady,1879
Atlantic
- Sporadogenerina** Cushman,1927 (TM) *Sporadogenerina flintii* Cushman,1927 = *Ramulina proteiformis* Flint,1899
- Ellipsolagenidae** A. Silvestri,1923
- Oolininae** Loeblich & Tappan,1961
- Anturina** R.W. Jones,1984
- A. haynesi** R.W. Jones,1984 (TM)
= *Oolina laevigata* : Haynes,1973, non d'Orbigny,1839
Wales, Scilly Islands
- Buchnerina** R.W. Jones,1984 (TM) *Buchnerina iberica* R.W. Jones,1984
- Cushmanina** R.W. Jones,1984 (TM) *Lagena vulgaris* Williamson var. *desmorphora* F.W.O. Rymer Jones,1872
- Favulina** Patterson & Richardson,1987
- F. hexagona** (Williamson,1848)
= *Entosolenia squamosa* (Montagu) var. *hexagona* Williamson,1848 (TM)
= *Lagena hexagona* : Auctt.
= *Oolina hexagona* : Loeblich & Tappan,1953
= *Lagena favosa* Reuss,1863
Oslofjord, Bergen, North Cape, Faeroe Channel, Barents Sea, Spitsbergen, Iceland, Belgium, North Sea, Shetlands, Irish Sea, Celtic Sea, W Scotland, Wales, Engl. Channel, Arcachon
- Galwayella** Patterson & Pettis,1986 (TM) *Lagena trigonoelliptica* Balkwill & Millett,1884
- Heteromorpha** R.W. Jones,1984 (TM) *Oolina heteromorpha* Parr,1950
- Homalohedra** Patterson & Richardson,1987 (TM) *Lagena guntheri* Earland,1934
- Laculatina** Patterson & Richardson,1987 (TM) *Lagena quadrilatera* Earland, var. *striatula* Earland,1934
- Lagnea** Popescu,1983 (TM) *Fissurina radiata* Seguenza,1862
= *Solenina* R.W. Jones,1984 (TM) *Lagenosolenita?* *tenuistriatiformis* McCulloch,1977
- Oolina** d'Orbigny,1839 (TM) *O. laevigata* d'Orbigny,1839
= *Cenchridium* Ehrenberg,1845 (TM) *Cenchridium sphaerula* Ehrenberg,1845
= *Entosolenia* Williamson,1848 (TM) *Entosolenia lineata* Williamson,1848
- O. lineata** (Williamson,1848)
= *Entosolenia lineata* Williamson,1848, non : Cushman,1948
= *Lagena lineata* : Auctt.
North Sea, Shetlands, W Scotland, Irish Sea, Wales, Engl. Channel, Spitsbergen, NW Greenland
- O. melo** d'Orbigny,1839
= *Entosolenia squamosa* (Montagu) var. *catenulata* Williamson,1848
= *Entosolenia squamosa* (Montagu) var. *scalariformis* Williamson,1848
= *Lagena catenulata* : Cushman,1923
= *Lagena squamosa* : Brady,1884
Oslofjord, Spitsbergen, Barents Sea, Belgium, S & SW Britain, Engl. Channel, Iceland, N & NW Greenland, Biscay
- O. squamosa** (Montagu,1803)
= *Vermiculum squamosum* Montagu,1803
= *Entosolenia squamosa* : Auctt., non : Cushman,1941, nec : Cushman & Gray,1946, nec : Cushman & Todd,1947
= *Lagena squamosa* : Auctt., non : Cushman,1913
Bergen, Vadsø, Iceland, Spitsbergen, Novaja Zemlja, Franz Josefs Land, Barents Sea, Belgium, NE Greenland, W Scotland, Ireland, Engl. Channel, Wales, Arcachon
- O. squamosa** (Montagu) var. *montagui* (Alcock,1865)
= *Entosolenia montagui* Alcock,1865
Iceland, Engl. Channel, W Ireland, Barents Sea
- O. striatopunctata** (Parker & Jones,1865)
= *Lagena sulcata* (Walker & Jacob) var. *striatopunctata* (Parker & Jones,1865)
= *Lagena striatopunctata* : Auctt.
N & NW Greenland, Spitsbergen, Novaja Zemlja, Engl. Channel
- O. caudigera** (Wiesner,1931)
= *Lagena* (*Entosolenia*) *globosa* (Montagu) var. *caudigera* Wiesner,1931
= *Entosolenia lineata* : Cushman,1948, non Williamson,1848
Spitsbergen, NE, N & NW Greenland
- O. borealis** Loeblich & Tappan,1954
= *O. costata* (Williamson,1858, non (?))
= *Entosolenia costata* Williamson,1858
= *Lagena costata* : Auctt.
= *Oolina patonae* Haman,1966
Faeroes, NW & W Scotland, Ireland, Wales, Engl. Channel, N & NW Greenland
- O. globosa** (Montagu,1803)
= *Vermiculum globosum* Montagu,1803
= *Lagena globosa* : Auctt.
Drobak, Stavanger, Bergen, Vesteraalen, W Scotland, Ireland, Engl. Channel, Barents Sea, Spitsbergen, Novaja Zemlja, Iceland
- O. heronalleni** Haynes,1973
Wales, W Scotland
- O. williamsoni** (Alcock,1865)
= *Entosolenia williamsoni* Alcock,1865
= *Lagena williamsoni* : Auctt.
North Sea, Spitsbergen, Belgium, N Kent, Engl. Channel, Bristol Channel, Celtic Sea, Irish Sea, Wales, Arcachon, Biscay, Mediterranean
- O. trigonooblonga** Seguenza & Siddall,18?? : Gabel,1971
North Sea
- O. elegantissima** (Bornemann,18??) : Gabel,1971
- Vasicostella** Patterson & Richardson,1987 (TM) *Lagena vulgaris* Williamson var. *helephoromarginata* F.W.O. Rymer Jones,1872
- Ellipsolageninae** A. Silvestri,1923
- Fissurina** Reuss,1850
= *Ellipsolagenina* A. Silvestri,1923 (TM) *Lagena acutissima* Fornasini,1889
- = *Ellipsofissurina* A. Silvestri,1923 (TM) *Fissurina laevigata* Reuss,1850
- F. laevigata** Reuss,1850 (TM)
= *Lagena laevigata* : Nørvang,1941
Bergen, Iceland, Novaja Zemlja, Franz Josefs Land, around England, Engl. Channel
- F. cucurbitasema** Loeblich & Tappan,1953
Engl. Channel, Spitsbergen, NE & N Greenland
- F. elliptica** (Cushman,1923)
= *Lagena orbigniana* (Seguenza) var. *elliptica* Cushman,1923
Wales, Caribbean
- F. lagenoides** (Williamson,1858)
= *Entosolenia marginata* (Montagu) var. *lagenoides* Williamson,1858
Engl. Channel, Iceland, Barents Sea, Novaja Zemlja, Franz Josefs Land
- F. lagenoides** (Williamson,1858) var. *tenuistriata* (Brady,1881)
= *Lagena tubulifera* Brady, var. *tenuistriata* Brady,1881
Britain, Mediterranean
- F. lucida** (Williamson,1848)
= *Entosolenia marginata* (Montagu) var. *lucida* Williamson,1848
= *Entosolenia lucida* : Auctt., non : Cushman,1948
S North Sea, Engl. Channel, Bristol Channel, Wales, Greenland, Biscay, Mediterranean
- F. marginata** (Walker & Boys,1784)
= *Serpula* (*Lagena*) *marginata* Walker & Boys,1784
= *Vermiculum marginatum* Montagu,1803
= *Lagena marginata* : Auctt.
= *Entosolenia marginata* : Auctt.
Bergen, Iceland, Spitsbergen, Barents Sea, Celtic Sea, Wales, Engl. Channel, NE & NW Greenland
- F. marginata** var. *inaequilateralis* (Wright,1885-86)
= *Lagena marginata* var. *inaequilateralis* Wright,1885-86
* Correct genus?
Engl. Channel, W Scotland, SW Ireland
- F. marginata** var. *semicarinata* Sidebottom,1906
* Correct genus?
?W Ireland
- F. marginata** var. *semimarginata* Reuss,18??
* Correct genus?
W Scotland
- F. marginata** var. *marginata* Seguenza,1880
Engl. Channel, W Ireland, W Scotland
- F. orbigniana** Seguenza,1862
Norway, Barents Sea, Iceland, Celtic Sea, Engl. Channel, Arcachon, Biscay
- F. orbigniana** Seguenza,1862, var. *clathrata* (Brady,18???)
?Britain, Indo-Pacific
- F. orbigniana** Seguenza,1862, var. *variabilis* (Wright,1891)
= *Lagena orbigniana* Seguenza,1862, var. *variabilis* (Wright,1891)
off SW Ireland at 1370 m.
- F. orbigniana** Seguenza,1862, var. *walleriana* (Wright,1891)
= *Lagena orbigniana* Seguenza,1862, var. *walleriana* Wright,1891
W Scotland off SW Ireland
- F. quadrata** (Williamson,1858)
= *Entosolenia marginata* (Montagu) var. *quadrata* Williamson,1858
Engl. Channel
- F. serrata** (Schlumberger,1894)
= *Lagena serrata* Schlumberger,1894
= *Entosolenia serrata* : Cushman,1948
Oslofjord, NE & NW Greenland, Arctic Russia
- F. pseudoglobosa** (Buchner,1940)
= *Lagena pseudoglobosa* Buchner,1940
Oslofjord
- F. fasciata** (Egger,1857)
= *Lagena fasciata* Egger,1857
Oslofjord, off W Scotland, W Ireland
- F. rizae** Seguenza,1862
Engl. Channel
- F. bicarinata** Terquem,1882
Engl. Channel
- F. acuta** Reuss,1858
* Correct genus? Correct year (perhaps 1863)?
Engl. Channel, W Scotland, W Ireland
- F. schlichti** Silvestri,1902
* Correct genus?
Engl. Channel
- F. staphylearia** Schwager,1866
off SW Ireland in deep water
- F. danica** Madson,19?? : Gabel,1971
North Sea
- F. sp. A** : Haynes,1973
Wales
- F. sp. B** : Haynes,1973
Wales
- Lagenosolenia** McCulloch,1977 (TM) *Lagenosolenia soulei* McCulloch,1977
- Palliolatella** Patterson & Richardson,1987 (TM) *Palliolatella avita* Patterson & Richardson,1987
- Pseudoolina** R.W. Jones,1984 (TM) *Pseudoolina fissurinea* R.W. Jones,1984
- Parafissuriniae** R.W. Jones,1984
- Cursina** Patterson & Richardson,1987 (TM) *Cursina adorna* Patterson & Richardson,1987
- Parafissurina** Parr,1947 (TM) *Lagena ventricosa* A. Silvestri,1904
- P. malcomsonii** (Wright,1911)
= *Lagena laevigata* (Reuss), var. *malcomsonii* Wright,1911
Engl. Channel, W Ireland, W Scotland

- P. lateralis** (Cushman,1913)
= *Lagena lateralis* Cushman,1913
Oslofjord
- P. elliptica** (Seguenza,18??) : Gabel,1971
North Sea
- Pseudofissurina** R.W. Jones,1984 (TM *Pseudofissurina mccullochae* R.W. Jones,1984)
- Pseudosolenina** R.W. Jones,1984 (TM *Pseudosolenina borealis* R.W. Jones,1984)
- Ventrostoma** Schnitter,1970 (TM *Lagena favigera* Buchner,1940)
- V. depressa** (Chaster,1892)
= *Lagena depressa* Chaster,1892 (TM of *Arthurina* R.W. Jones,1984)
- V. cymbula** Heron-Allen & Earland,1913
W North Sea, NE Scotland, W Ireland
- Walterparria** R.W. Jones,1984
- W. milletti** (Chaster,1892)
= *Lagena milletti* Chaster,1892 (TM)
= *Parafissurina milletti* Murray,1971
Engl. Channel
- Wiesnerina** R.W. Jones,1984
- W. unguis** (Heron-Allen & Earland,1913)
= *Lagena unguis* Heron-Allen & Earland,1913 (TM)
North Sea, around Scotland, Cornwall
- Sipholageninae** Patterson & Richardson,1987
- Bifarilaminella** Patterson & Richardson,1987 (TM *Lagena advena* Cushman,1923)
- Sipholagena** Moncharmont Zei & Sgarella,1980 (TM *Lagena benevistita* Buchner,1940)
= *Buchneria* Moncharmont Zei & Sgarella,1977, non Börner,1952
- Glandulinidae** Reuss,1860
- Glandulininae** Reuss,1860
- Eosyrinx** Loeblich & Tappan,1953 (TM *Pseudopolymorpha curta* Cushman & Ozawa,1930 - Ungava Bay & W Greenland)
- Glandulina** d'Orbigny,1839
= *Psecadium* Neugeboren,1856 (TM *Psecadium ellipticum* Neugeboren,1856)
- G. laevigata** (d'Orbigny,1826)
= *Nodosaria* (les Glandulines) *laevigata* d'Orbigny,1826 (TM)
Bergen, Vesterålen, Iceland, N Greenland, Barents Sea, ?Shetlands, ?W Scotland, ?warm Faeroe Channel, ?off SW Ireland, ?Sussex
- G. laevigata** (d'Orbigny,1826) var. *occidentalis* (Cushman,1923)
= *Nodosaria laevigata* (d'Orbigny,1826) var. *occidentalis* Cushman,1923
*Correct genus?
Nordland, NE USA
- Laryngosigma** Loeblich & Tappan,1953
- L. hyalascidia** Loeblich & Tappan,1953 (TM)
SW Baltic, NE, N & NW Greenland, Wales
- L. lactea** (Walker & Jacob,1798)
= *Serpula lactea* Walker & Jacob,1798
= *Gutulina lactea* : Auct.
= *Polymorphina lactea* : Auctt.
Oslofjord, North Cape, ?Iceland, Engl. Channel, Wales, Novaja Zemlya
- L. lactea** (Walker & Jacob,1798) var. *concava* (Williamson,1858)
= *Polymorphina lactea* (Walker & Jacob) var. *concava* Williamson,1858
Wales, Engl. Channel, Mediterranean
- L. williamseni** (Terquem,1878)
= *Polymorphina williamseni* Terquem,1878
= *Polymorphina lactea* (Walker & Jacob) var. *oblonga* Williamson,1858, non *Polymorphina* (*Globulinea*) *oblonga* Roemer,1838, nec *Polymorphina oblonga* d'Orbigny,1846
Plymouth, NE Greenland
- Phlegeria** Loeblich & Tappan,1963 (TM *Phlegeria hyalina* Loeblich & Tappan,1963)
- Tappanella** Gudina & Saidova,1969 (TM *Tappanella arctica* Gudina & Saidova,1969 = *Glandulina laevigata* d'Orbigny : Gudina,1966, non *Glandulina laevigata* d'Orbigny)
- Entoligininae** Saidova,1981
- Entolingulina** Loeblich & Tappan,1961 (TM *Lingulina aselliformis* Buchner,1942)
Seabrookianae Cushman,1927
- Seabrookia** Brady,1890 (TM *Seabrookia pellucida* Brady,1890)
= *Miletia* J. Wright,1889, non Duncan,1889 (name not available)
= *Cerviciferina* Goddard & Jensen,1907 (TM *Cerviciferina hilli* Goddard & Jensen,1907)
- S. earlandi** (Wright,1889)
= *Miletia earlandi* Wright,1889
Norwegian fjords, North Sea, W & SW Ireland
- ROBERTININA** Loeblich & Tappan,1984
- Ceratobuliminaceae** Cushman,1927
- Ceratobuliminidae** Cushman,1927
- Ceratobulimininae** Cushman,1927
- Ceratobulimina** Toula,1915 (TM *Rotalina contraria* Reuss,1851)
- C. arctica** Green,1960
Oslofjord
- Lamareckina** Berthelin,1881 (TM *Pulvinulina erinacea* Karrer,1868)
= *Megalostomina* Rzehak,1891 (TM *Megalostomina fuchsii* Rzehak,1895 = *Discorbina fuchsii* Rzehak,1888)
- L. haliotidea** (Heron-Allen & Earland,1911)
= *Pulvinulina haliotidea* Heron-Allen & Earland,1911
* clinging to firm substrates.
Belgium, Engl. Channel, off Faeroes, W Scotland, W Ireland
- Rubratella** Grell,1956 (TM *Rubratella intermedia* Grell,1956 - France)
- Epistominidae** Wedekind,1937
- Epistomininae** Wedekind,1937
- Epistomina** Terquem,1883 (TM *Epistomina regularis* Terquem,1883)
= *Rotalia Hofker,1954* (TM *Rotalia spinulifera* Reuss,1863)
= *Voorthuyseana* Hofker,1954 (TM *Epistomina tenuicostata* Bartenstein & Brand,1951)
= *Hillermannia* Hofker,1954 (TM *Epistomina chapmani* ten Dam,1948)
= *Kaptarenkella* Antonova & G.K. Kasimova, in G.K. Kasimova & Antonova,1985 (TM *Lamarckella epistominoidea* Kaptarenko-Chernousova,1959)
- Hoeglundina** Brozen,1948 (TM *Rotalia elegans* d'Orbigny,1826)
- H. elegans** (d'Orbigny,1826)
= *Rotalia elegans* d'Orbigny,1826 (TM)
= *Rotalina partchiana* d'Orbigny,1846
boreal Atlantic, ?off Norway, Spitsbergen, off SW Ireland at 620-695 fathoms
- Robertinacea** Reuss,1850
- Robertinidae** Reuss,1850
- Alliatininae** McGowran,1966
- Alliatina** Troelsen,1954 (TM *Cushmanella excentrica* di Napoli Alliata,1952)
= *Pseudononionella* S.Y. Zheng, in S.Y. Zheng, Cheng, Wang & Fu,1978 (TM *Pseudononionella variabilis* S.Y. Zheng, in S.Y. Zheng, Cheng, Wang & Fu,1978)
- Alliatinella** D.J. Carter,1957 (TM *Alliatinella gedgravensis* D.J. Carter,1957)
= *Fawcettia* McCulloch,1977 (TM *Fawcettia panayensis* McCulloch,1977)
= *Subcushmanella* McCulloch,1977 (TM *Subcushmanella differens* McCulloch,1977)
=? *Subceroberolina* McCulloch,1977 (TM *Subceroberolina simplissima* McCulloch,1977)
- Robertininae** Reuss,1850
- Robertina** d'Orbigny,1846
- R. arctica** d'Orbigny,1846 (TM)
Iceland, Barents Sea, Spitsbergen, Greenland (Clausshavn), N Norway, Siberia
- Robertinoides** Höglund,1947
- R. normani** (Goës,1894)
= *Bulmina normani* Goës,1894 (TM)
Koster Channel, Säcken, Skagerrak
- R. sueicum** Höglund,1947
Gullmarfjord, Koster Channel, Skagerrak, Farsund, Börnestangen
- R. pumilum** Höglund,1947
Gullmarfjord, Skagerrak
- R. charlottenensis** (Cushman,1925)
= *Cassidulina charlottenensis* Cushman,1925
= *Roberina charlottenensis* : Auct.
= *Roberina californica* Cushman & F. Parker,1936
NE, N & NW Greenland
- GLOBIGERININA** Delage & Hérouard,1896
- Heterohelicacea** Cushman,1927
- Chiloguembelinidae** Reiss,1963
- Laterostomella** de Klasz & Réat,1962 (TM *Laterostomella guembeliniformis* de Klasz & Réat,1962)
= *Streptochilus* Brönnimann & Resig,1971 (TM *Bolivina tokelauae* Boersma, in Kierstead & al.,1969 = *Bolivina globulosa* Cushman,1933)
=? *Pseudoheterohelix* Aliyulla,1977 (TM *Guembelia pumilia* Subbotina,1950)
- Globorotaliacea** Cushman,1927
- Globorotaliidae** Cushman,1927
- Globorotalia** Cushman,1927
= *Planorotalia* Morozova,1957 (TM *Planulina membranacea* Ehrenberg,1854)
= *Obandyella* Haman & al.,1981 (TM *Rotalina hirsuta* d'Orbigny,1839)
- G. tumida** (Brady,1877)
= *Pulvinulina menardii* (d'Orbigny) var. *tumida* Brady,1877 (TM)
off SW Ireland
- G. hirsuta** (d'Orbigny, in Barker-Webb & Berthelot,1839)
= *Rotalina hirsuta* d'Orbigny, in Barker-Webb & Berthelot,1839
N Atlantic (subtropical), off W & NW Hebrides, off W Ireland
- G. inflata** (d'Orbigny, in Barker-Webb & Berthelot,1839)
= *Globigerina inflata* d'Orbigny, in Barker-Webb & Berthelot,1839
N Atlantic (cold-temperate), Wales
- G. scitula** (Brady,1882)
= *Pulvinulina scitula* Brady,1882
N Atlantic (subpolar to tropical, especially below 500 m.), off W Ireland, Engl. Channel, off W Hebrides
- G. menardii** (d'Orbigny,1826)
= *Rotalia menardii* d'Orbigny,1826
= *Pulvinulina menardii* : Auct.
* Correct genus?
Plymouth, pelagic between 55° 11' N & 51° 36' S
- G. menardii** (d'Orbigny,1826), var. *fimbriata* (Brady,1884)
Atlantic
- Neogloboquadrina** Bandy, Frerichs & Vincent,1967
- N. dutertrei** (d'Orbigny,1839)
= *Globigerina dutertrei* d'Orbigny,1839 (TM)
Plymouth, North Sea
- Truncorotalia** Cushman & Bermúdez,1949
- T. truncatulinoides** (d'Orbigny, in Barker-Webb & Berthelot,1839)
= *Rotalina truncatulinoides* d'Orbigny, in Barker-Webb & Berthelot,1839 (TM)
= *Globorotalia truncatulinoides* : Auctt.
N Atlantic (subtropical), off W Shetlands, off SW Ireland
- Pulleniatinidae** Cushman,1927
- Pulleniatina** Cushman,1927 (TM *Pullenia obliquiloculata* Parker & Jones,1865)
- Candeinidae** Cushman,1927
- Tenuitellinae** Banner,1982
- Tenuitella** Fleisher,1984 (TM *Globorotalia gemma* Jenkins,1966)

- Globigerinitinae Bermúdez,1961
- Tinophodella** Loeblich & Tappan,1957 (TM *Tinophodella ambitacrena* Loeblich & Tappan,1957)
- Candeina d'Orbigny, in de la Sagra,1839
- C. nitida* d'Orbigny, in de la Sagra,1839 (TM)
off S Iceland, Plymouth, West Indies
- Catapsydracidae Bolli, Loeblich & Tappan,1957
- Globoquadrina** Finlay,1947 (TM *Globorotalia dehiscens* Chapman, Parr & Collins,1934)
- G. hexagona** (Natland,1938)
= *Globigerina hexagona* Natland,1938
Wales
- Globigerinacea Carpenter, Parker & Jones,1862
- Globigerinidae Carpenter, Parker & Jones,1862
- Globigerininae Carpenter, Parker & Jones,1862
- Bella** Banner & Blow,1960 (TM *Globigerina digitata* Brady,1879)
- Globigerina** d'Orbigny,1826
- G. bulloides** d'Orbigny,1826 (TM)
* Absent from regions with strong coastal influence
W Skagerrak, N Atlantic (subpolar, cold temperate), N North Sea, Iceland
- G. falconensis** Blow,1959
N Atlantic (cold-temperate & subtropical)
- G. pachyderma** (Ehrenberg,1861)
= *Asterospira pachyderma* Ehrenberg,1861
* Bottom-living
N Atlantic (subpolar, cold temperate), North Sea, Faeroe Channel, Iceland, W Scotland, W Ireland, Barents Sea
- G. quinqueloba** Natland,1938
N Atlantic (cold-water, arctic to subarctic, cold temperate surface waters)
- G. glutinata** (Egger,1895)
= *Globigerina glutinata* Egger,1895
N Atlantic (tropical to subpolar)
- G. triloba** (Reuss,18??) : Gabel,1971
North Sea
- G. eggeri** Rhumbler,19?? : Gabel,1971
North Sea
- G. sp.** : Gabel,1971
North Sea
- Globigerinella** Cushman,1927 (TM *Globigerina aequilateralis* Brady,1879, non *Globigerina siphonifera* d'Orbigny,1839)
- G. aequilateralis** (Brady,1879)
= *Globigerina aequilateralis* Brady,1879
Gulf Stream waters, Caribbean
- Globigerinoides** Cushman,1927
- G. rubra** (d'Orbigny, in de la Sagra,1839)
= *Globigerina rubra* d'Orbigny, in de la Sagra,1839 (TM)
Gulf Stream waters - W coast of Europe, Caribbean, Gulf of Mexico
- G. rubra** var. **elevata** (d'Orbigny,1840)
= *Globigerina elevata* d'Orbigny,1840
* Correct genus?
Plymouth
- G. conglobata** (Brady,1879)
= *Globigerina conglobata* Brady,1879
Gulf Stream waters, Caribbean
- Globoturborotalita** Hofker,1976 (TM *Globigerina rubescens* Hofker,1956)
- Turborotalita** Blow & Banner,1962 (TM *Truncatulina humilis* Brady,1884)
- Orbulininae Schultze,1854
- Orbulina** d'Orbigny,1839
= *Coscinospaera* Stuart,1866 (TM *Coscinospaera ciliosa* Stuart,1866)
- O. universa** d'Orbigny, in Ramon de la Sagra,1839 (TM)
N Atlantic (tropical to subtropical), Wales, Iceland, Novaja Zemlya
- O. bilobata** (d'Orbigny,1846)
= *Globigerina bilobata* d'Orbigny,1846 (TM of *Biorbulina* Blow,1956)
= *Biorbulina bilobata* : Haynes,1973
Wales
- O. nitida** Terquem & Terquem,1886
Christiansand, Stavanger, Iceland
- Hastigerinidae Bolli, Loeblich & Tappan,1957
- Hastigerina** Thomson, in Murray,1876
- H. pelagica** (d'Orbigny,1839)
= *Nonionina pelagica* d'Orbigny,1839
= *Hastigerina murrayi* Thomson, in Murray,1876 (TM)
warm Faeroe Channel, off W Orkneys, off SW Ireland at 1370 m.
- Orcadia** Boltovskoy & Watanabe,1982 (TM *Hastigerinella riedeli* Rögl & Bolli,1973)
- ROTALINA Delage & Hérouard,1896
- Bolivinaceae Glaessner,1937
- Boliviniidae Glaessner,1937
- Bolivina** d'Orbigny,1839 (TM *Bolivina plicata* d'Orbigny,1839)
= *Grammostomum* Ehrenberg,1839 (TM *Grammostomum tenuis* Ehrenberg,1839)
= *Proroporus* Ehrenberg,1844 (TM *Proroporus lingua* Ehrenberg,1844)
= *Clidostomum* Ehrenberg,1845 (TM *Clidostomum polystigma* Ehrenberg,1845)
- B. pseudoplicata** Heron-Alen & Earland,1930
= *B. plicata* Goës,1894, non d'Orbigny,1839
- Kattegatt, Gullmarfjord, Skagerrak, Oslofjord, Bergen, Trondheim Fjord, Bukken, Belgium, Ireland, N Kent, Wales, S Britain, Arcachon, Biscay
- B. albatrossi** Cushman,1922
Gullmarfjord, Skagerrak, Oslofjord, Trondheim Fjord, Gulf of Mexico
- B. cf. robusta** Brady,1884
Gullmarfjord, Koster Channel, Skagerrak, Oslofjord, Bergen
- B. punctata** d'Orbigny,1839
* Correct genus?
along entire Norway, Iceland, Spitsbergen, Novaja Zemlya, around Britain
- B. gramen** (d'Orbigny, in de la Sagra,1839)
= *Valvulina gramen* d'Orbigny, in de la Sagra,1839
Gullmarfjord, Cuba
- B. variabilis** (Williamson,1858)
= *Textularia variabilis* Williamson,1858
= *Brizalina variabilis* : Murray,1971
S North Sea, Belgium, Engl. Channel, Bristol Channel, Celtic Sea, Wales, Mediterranean
- B. decussata** : Wright,1891, ?non Brady,1881
off SW Ireland at 1866 m.
- B. limbata** Brady,1881
W Ireland, Cape Verde
- B. nobilis** Hantken,1876
?Britain, ?Mediterranean
- B. durrandi** Millett,1900
?Sussex, Malay Archipelago
- B. costata** d'Orbigny,18??
?Britain
- B. (?) sp.** : Höglund,1947
Skagerrak
- Bolivinellina** Saidova,1975 (TM *Bolivinellina pescicula* Saidova,1975)
= *Bolivinellina* Burmistrova,1974 (name not available)
- Brizalina** O.G. Costa,1856 (TM *Brizalina aenariensis* O.G. Costa,1856)
- B. pseudopunctata** (Höglund,1947)
= *Bolivina pseudopunctata* Höglund,1947
Gullmarfjord, Koster Channel, Skagerrak, Oslofjord, off Bergen, Barents Sea, off Spitsbergen, Kent, Wales, Engl. Channel, Celtic Sea, N & NW Greenland
- B. difformis** (Williamson,1858)
= *Textularia variabilis* Williamson var. *difformis* Williamson,1858
= *Bolivina difformis* : Auctt.
Barents Sea, Ireland, Engl. Channel, Arcachon, Biscay
- B. spathulata** (Williamson,1858)
= *Textularia variabilis* Williamson var. *spathulata* Williamson,1858
= *Bolivina spathulata* : Auctt.
Bolivina dilatata : Brady,1884, : Goës,1894, non Reuss,1850
Gullmarfjord, Koster Channel, Skagerrak, Oslofjord, Shetlands, W Scotland, North Sea, Belgium, Barents Sea, Engl. Channel, Wales, Celtic Sea, Arcachon, Biscay, Galicia
- B. subaenariensis** (Cushman,1922)
= *Bolivina subaenariensis* Cushman,1922
Celtic Sea
- B. britannica** (Macfadyen,1942)
= *Bolivina britannica* Macfadyen,1942
= *Textularia variabilis* Williamson, var. *laevigata* Williamson,1858
= *Bolivina laevigata* : Auctt., non *Textularia laevigata* d'Orbigny,18??
North Sea, Faeroes, W Scotland, Wales
- B. inflata** (Heron-Alen & Earland,1913)
= *Bolivina inflata* Heron-Alen & Earland,1913
Wales, W Scotland, W Ireland, Engl. Channel
- B. minima** (Phleger & Parker,1951)
= *Bolivina minima* Phleger & Parker,1951
Wales, Gulf of Mexico
- B. ordinaria** (Phleger & Parker,1952)
= *Bolivina ordinaria* Phleger & Parker,1952
= *Bolivina simplex* Phleger & Parker,1951, non ?,1???
Wales, Gulf of Mexico
- B. striatula** (Cushman,1922)
= *Bolivina striatula* Cushman,1922
Gullmarfjord, Kent, Engl. Channel, Wales, Galicia, Florida
- B. superba** (Emiliani,1949)
= *Bolivina superba* Emiliani,1949
= *Bolivina beyrichii* : Heron-Alen & Earland,1916, : Cushman,1923, non Reuss,1851
= *Bolivina alata* : Cushman,1937, non *Vulvulina alata* Sequeira,1862
= *Bolivina barbata* Phleger & Parker,1951
= *Bolivina difformis* : Colom,1952, non (Williamson,1858)
Scotland, Ireland, Wales, Galicia
- B. textilaroides** (Reuss,18??)
= *Bolivina textilaroides* Reuss,18??
Helgoland, Britain
- B. cf. vadescens** (Cushman,1933)
= *Bolivina vadescens* Cushman,1933
Kattegatt, Gullmarfjord, Skagerrak, Wales
- B. alata** (Seguenza,18??) : Gabel,1971
North Sea
- B. earlandi** (Parr,1???) : Gabel,1971
North Sea
- B. subspinosa** (Cushman,1922)
= *Bolivina subspinosa* Cushman,1922
- Latibolivina** Srinivasan,1966 (TM *Bolivina anastomosa* Finlay,1939)
- Lugdunum** Saidova,1975 (TM *Bolivina hantkeniana* Brady,1881)
- Bolivinitacea Cushman,1927
- Bolivinitidae Cushman,1927
- Bolivinita** Cushman,1927

- B. quadrilatera** (Schwager,1866)
= *Textularia quadrilatera* Schwager,1866 (TM)
off SW Ireland, off Canaries, off Cape Verde
- Cassidulinacea** d'Orbigny,1839
- Cassidulinidae** d'Orbigny,1839
- Cassidulininae** d'Orbigny,1839
- Cassidulina** d'Orbigny,1826
= *Entrochus* Ehrenberg,1843 (TM) *Entrochus septatus* Ehrenberg,1843
= *Selenostomum* Ehrenberg,1858
= *Lernina* Saidova,1975 (TM) *Lernina micae* Saidova,1975
= *Lernina* Khushid,1973 (name not available)
- C. laevigata** d'Orbigny,1826 (TM)
* very common in W Norway
Bohusl., Oslofjord, Bergen, Vadso, Spitsbergen, Novaja Zemlya, Franz Josefs Land, Arctic Ocean, Greenland, Iceland, Shetlands, W Scotland, off W Ireland
- C. obtusa** Williamson,1858
= *C. crassa* : Auct., non d'Orbigny,1839
Oslofjord, Barents Sea, Spitsbergen, Novaja Zemlya, Iceland, Greenland, Arctic Ocean, Belgium, Engl. Channel, Celtic Sea, Arcachon, Biscay
- C. carinata** Silvestri,1896
= *Cassidulina laevigata* d'Orbigny var. *carinata* Silvestri,1896
Celtic Sea, S Cornwall, western approaches to the Engl. Channel, Biscay, Galicia
- C. reniforme** Nørvang,1945
= *Cassidulina crassa* d'Orbigny var. *reniforme* Nørvang,1945
Iceland, ?Wales
- C. curvata** Phleger & Parker,19?? : Gabel,1971
North Sea
- C. sp.**, aff. *C. pulchella* d'Orbigny,18?? : Gabel,1971
North Sea
- C. cf. translucens** Cushman & Hughes,1925 : Nørvang,1945
Iceland
- Cassidulinoides** Cushman,1927 (TM) *Cassidulina parkeriana* Brady,1881
- C. bradyi** (Norman, in Wright,1881)
= *Cassidulina bradyi* Norman, in Wright,1881
Østerfjord, Bergen, N North Sea, Faeroe Channel, Iceland, W Scotland, off W & SW Ireland, western approaches of the Engl. Channel at >420 m., Biscay
- Globocassidulina** Voloshinova,1960 (TM) *Cassidulina globosa* Hantken,1876
= *Cassilongina* Voloshinova,1960 (TM) *Cassidulina oblonga* Reuss,1850
= *Smyrnella* LA. Basov,1974 (name not available)
= *Sphaeroislandiella* Burmistrova,1974 (name not available)
= *Bryadynella* Burmistrova,1974 (name not available)
= *Cassidulinella* Saidova,1975 (TM) *Cassidulinella salebrosa* Saidova,1975
= *Smyrnella* Saidova,1975 (TM) *Cassidulina subglobosa* Brady var. *subcalifornica* Drooger,1953
= *Sphaeroislandiella* Saidova,1975 (TM) *Sphaeroislandiella notabilis* Saidova,1975
= *Bryadynella* Saidova,1975 (TM) *Cassidulina subglobosa* Brady,1881
- G. aff. G. subglobosa** (Brady,1881)
Barents Sea, Celtic Sea, western approaches to the Engl. Channel, Biscay
- G. oblonga** (Reuss,1850)
= *Cassidulina oblonga* Reuss,1850
Ireland
- G. nitidula** (Chaster,1892)
= *Pulvinulina nitidula* Chaster,1892
North Sea, Iceland, W Scotland, W Ireland, S Cornwall, Mediterranean
- Islandiella** Nørvang,1959 (TM) *Cassidulina islandica* Nørvang,1945
= *Cassilamellina* Voloshinova,1960 (TM) *Cassidulina californica* Cushman & Hughes,1925
= *Planocassidulina* Gudina,1966 (TM) *Cassidulina norcrossi* Cushman,1933
= *Cassandra* Gudina & Saidova, in Gudina & al.,1968 (TM) *Cassidulina inflata* Gudina,1966
= *Cassidulina* Sellier de Civrieux,1969 (TM) *Cassidulina norcrossi* Cushman,1933
? = *Rosaella* Saidova,1975 (TM) *Rosaella rosae* Saidova,1975
= *Discoislandiella* Saidova,1975 (TM) *Cassidulina smechovi* Voloshinova, in Voloshinova & Dain,1952
- I. islandica** (Nørvang,1945)
= *Cassidulina islandica* Nørvang,1945
= *Cassidulina islandica* Nørvang,1945 forma *minuta* Nørvang,1945, non *Cassidulina minuta* Cushman,1933
= *Cassidulina islandica* Nørvang,1945 var. *noervangi* Thalmann, in Phleger,1952
Iceland
- I. helenae** Feyling-Hansen & Buzas,1976
= *Cassidulina teretis* : Loeblich & Tappan,1953, non Tappan,1951
= *Cassidulina laevigata* : Parker & Jones,1865, : Cushman,1948, non d'Orbigny,1826
Spitsbergen, NE, N, NW & Greenland, Russian Arctic
- I. teretis** Tappan,1951
= *Cassidulina teretis* Tappan,1953 : Gabel,1971
North Sea
- I. norcrossi** (Cushman,1933)
= *Cassidulina norcrossi* Cushman,1933
North Sea, Iceland, Spitsbergen, NE & NW Greenland
- Ehrenbergininae Cushman,1927
- Burseolina** Seguenza,1880 (TM) *Burseolina calabra* Seguenza,1880
= *Cassisphearia* Saidova,1975 (TM) *Cassisphearia globula* Saidova,1975
= *Cushmanulla* Saidova,1975 (TM) *Cassidulina pacifica* Cushman,1925
- Ehrenbergina** Reuss,1850 (TM) *Ehrenbergina serrata* Reuss,1850
- Turillinae** Cushman,1927
- Stainforthiidae** Reiss,1963
- Cassidelina** Saidova,1975 (TM) *Cassidelina profunda* Saidova,1975
- Hopkinsina** Howe & Wallace,1932 (TM) *Hopkinsina danvillensis* Howe & Wallace,1932 = *Virgulina recta* Cushman var. *howei* Cushman,1936
- S. concava** Hofker,1956
- S. concava** (Höglund,1947)
= *Virgulina concava* Höglund,1947 (TM)
Kattegatt, Gulmarfjord, Koster Channel, Skagerrak, Oslofjord, Barents Sea
- S. concava** (Höggrund,1947) var. *loeblichii* (Feyling-Hansen,1954)
- = *Virgulina loeblichii* Feyling-Hansen,1954
Wales, Oslofjord (late quaternary)
- S. subdepressa** (Brady,18??) : Gabel,1971
North Sea
- Buliminacea** Jones,1875
- Siphogenerinoidae** Saidova,1981
- Siphogenerinoidinae** Saidova,1981
- Loxostomatina** Sellier de Civrieux,1969, non Saidova,1975 (TM) *Bolivina majori* Cushman,1922
= *Loxostomella* Saidova,1975
- Parabrizalina** Zweig-Strykowski & Reiss,1976
- P. porrecta** (Brady,1881)
= *Bolivina porrecta* Brady,1881 (TM)
= *Loxostomum porrectum* Höglund,1947
Skagerrak, Bergen, Færør Channel
- Rectobolivina** Cushman,1927
- R. bifrons** (Brady,1881) : Gabel,1971
= *Sagrina bifrons* Brady,1881 (TM)
North Sea
- Saidovina** Haman,1984 (TM) *Bolivina karreriana* Brady,1881
= *Loxostomina* Saidova,1975, non Sellier de Civrieux,1969
- Tubulogeneneriniae** Saidova,1981
- Rectuvigerina** Mathews,1945 (TM) *Siphogenerina multicostata* Cushman & Jarvis,1929
- Sagrina** d'Orbigny,1839 (TM) *Sagrina pulchella* d'Orbigny,1839
- Sagrinopsis** Sellier de Civrieux,1969 (TM) *Siphogenerina advena* Cushman,1922
- Siphogenerina** Schlumberger,1882 (TM) *Siphogenerina costata* Schlumberger,1883
= *Ellipsosiphogenerina* A. Silvestri,1902
= *Rectuvigerina* Saidova,1975 (TM) *Siphogenerina raphanus* (Parker & Jones) var. *tropica* Cushman,1926
- S. dimorpha** (Parker & Jones,1865)
= *Uvigerina* (*Sagrina*) *dimorpha* Parker & Jones,1865
= *Rectobolivina* (*dimorpha*) Gabel,1971
Koster Channel, Christianssand, Bergen, Bukken Fjord, Vestfjord, Iceland, off W Scotland, Canaries
- Buliminidae** Jones,1875
- Bulima** d'Orbigny,1826
- B. marginata** d'Orbigny,1826 (TM)
* Very common in W Norway
Öresund, Kattegatt, Gulmarfjord, Skagerrak, Oslofjord, Bergen, N North Sea, Iceland, Barents Sea, Spitsbergen, Belgium, Shetlands, W Scotland, Irish Sea, Wales, Arcachon, Biscay, Galicia
- B. exilis** Brady,1884
= *B. elegans* d'Orbigny, var. *exilis* Brady,1884
W Scotland, Sussex, Faeroe Channel, NE, NW & W Greenland
- B. cf. alazanensis** Cushman,1927
western approaches of the Engl. Channel at >420 m., Biscay, Galicia
- B. gibba** Fornasini,1902
Wales, N Atlantic, Mediterranean, Adriatic
- B. elongata** d'Orbigny,1846
= *B. elegans* : Brady,1884, non d'Orbigny,1826
W Scotland, Irish Sea, Celtic Sea, Engl. Channel, Wales, Biscay, Galicia, Adriatic
- B. elongata** d'Orbigny,1846 var. *lesleyae* Atkinson,1969
= *B. elongata* : Brady,1884, non d'Orbigny,1846
Wales, N Atlantic
- B. elongata** d'Orbigny,1846 var. *subulata* Cushman & Parker,1937
= *B. aculeata* : Cushman,1944, non d'Orbigny,1826
Wales, Engl. Channel, Galicia
- B. aculeata** d'Orbigny,1826
Plymouth, off W Scotland
- B. echinata** d'Orbigny,1826
off W Ireland
- B. inflata** Seguenza,1862
off European coast in fairly deep water
- B. minutissima** Wright,1892
W Scotland, W Ireland, Cornwall
- B. squammigera** d'Orbigny, in Barker, Webb & Berthelot, 1839
around Britain, Canaries
- B. buchiana** d'Orbigny,1846
off SW Ireland at 87-1370 m.
- B. cf. elegans** d'Orbigny,1826
Iceland, SW Ireland, Plymouth
- B. doliolum** Terquem & Terquem,1886
S & SW Norway, Iceland
- B. ovoides** Terquem & Terquem,1886
S & SW Norway, Iceland
- B. pygmaea** d'Orbigny,18??
Iceland
- B. spinulosa** d'Orbigny,18??
Iceland
- B. submarginata** Parr,19?? : Gabel,1971
N North Sea
- Globobulimina** Cushman,1927 (TM) *Globobulimina pacifica* Cushman,1927
- G. turgida** (Bailey,1851)
= *Bulimina turgida* Bailey,1851
Kattegatt, Gulmarfjord, Skagerrak, Oslofjord, Bergen, N Norway
- G. auriculata** (Bailey,1851?) forma *gullmarenensis* Höglund,1947

- Gullmarfjord, Skagerrak
- G. auriculata* (Bailey,1851?) *forma arctica* Höglund,1947
Spitsbergen, Greenland
- Praeglobobulimina* Hofker,1951
- P. pyrula* (d'Orbigny,18??)
= *Bulmina pyrula* d'Orbigny,18??
* Correct genus?
Iceland
- P. pyrula* (d'Orbigny,18??) var. *spinescens* (Brady,1884)
= *Bulmina pyrula* d'Orbigny var. *spinescens* Brady,1884 (TM)
Bergen, Iceland
- Protoglobobulimina* Hofker,1951
- P. pupoides* (d'Orbigny,1846)
= *Bulmina pupoides* d'Orbigny,1846 (TM)
= *Praeglobobulimina pupoides* (d'Orbigny,1846) : Gabel,1971
North Sea, Iceland
- Buliminellidae* Hofker,1951
- Buliminella* Cushman,1911 (TM) *Bulmina elegantissima* d'Orbigny,1839
- B. borealis* Haynes,1973
= *B. elegantissima* : Auct., non (d'Orbigny,1839)
Øresund, Gullmarfjord, Skagerrak, Oslofjord, German North Sea coast, Wadden Sea, Belgium, Barents Sea, Shetlands, Irish Sea, Engl. Channel, Wales, Barents Sea, Arctic Ocean
- B. convoluta* (Williamson,1858)
= *Bulmina pupoides* d'Orbigny, var. *convoluta* Williamson,1858
Bergen, Shetlands, W Scotland, W Ireland, Cornwall
- B. subterea* (Brady,1881)
= *Bulmina subterea* Brady,1881
Norway, W Scotland, W & N Ireland, Cornwall, Faeroe Channel, Greenland, Spitsbergen, Novaja Zemlja
- Uvigerinidae* Haackel,1894
- Uvigerininae* Haackel,1894
- Euvigerina* Thalmann,1952
= *Euvigerina* Hofker,1951 (name not available)
- E. aculeata* (d'Orbigny,1846)
= *Uvigerina aculeata* d'Orbigny,1846 (TM)
off SW Ireland at 1829-1866 m.
- Neuvigerina* Thalmann,1952
= *Neuvigerina* Hofker,1950 & Hofker,1951 (name not available)
- N. ampullacea* (Brady,1884)
= *Uvigerina asperula* var. *ampullacea* Brady,1884 (TM)
off S Ireland at 1326 m.
- Uvigerina* d'Orbigny,1826 (TM) *Uvigerina pygmaea* d'Orbigny,1826 - fossil
= *Noviuvia Vella*,1963 (TM) *Uvigerina peregrina* Cushman,1923
- U. peregrina* Cushman,1923
= *U. pygmaea* : Goës,1894, non d'Orbigny,1826
* One of the most common species in W Norway.
Kattegatt, Gullmarfjord, Skagerrak, Oslofjord, North Sea, entire Norwegian coast, Barents Sea, Iceland, Celtic Sea, Biscay, Galicia, Novaja Zemlja, Franz Josef's Land
- U. canariensis* d'Orbigny,1839
W & SW Ireland, Wales, Sussex, Mediterranean, Canaries, Azores, Cape Verde
- U. auberiana* d'Orbigny, in de la Sagra,1839
off St Rockall Bank at 1152 m., Cuba
- U. porrecta* Brady,1879
? Sussex, Indo-Pacific
- U. asperula* Czjzek,18??
Atlantic
- U. tenuistrigata* Reuss,18??
- U. cf. striata* d'Orbigny,18?? : Gabel,1971
North Sea
- Angulogerininae* Galloway,1933
- Angulogerina* Cushman,1927
- A. angulosa* (Williamson,1858)
= *Uvigerina angulosa* Williamson,1858 (TM)
= *Trifarina angulosa* : Gabel,1971, : Murray,1979
Kattegatt, Skagerrak, Oslofjord, along Norway, North Sea, Bergen, Barents Sea, Iceland, Shetlands, Belgium, Engl. Channel, Bristol Channel, Celtic Sea, Wales, Arcachon, Biscay
- A. fluens* Todd, in Cushman & Todd,1947
= *A. angulosa* : Cushman,1948, non (Williamson,1858)
North Sea, Spitsbergen, NE, N, NW & W Greenland
- Trifarina* Cushman,1923
- T. bradyi* Cushman,1923 (TM)
= *Rhabdogonium tricarinatum* : Brady,1884, non *Vaginulina tricarinata* d'Orbigny,18??
North Sea, western approaches of the Engl. Channel, off Ireland, Biscay
- T. pygmaea* Gabel,1971
S North Sea
- Reussellidae* Cushman,1933
- Reussella* Galloway,1933
= *Reussia* Schwager,1877, non McCoy,1854
- R. spinulosa* (Reuss,1850)
= *Verneuilina spinulosa* Reuss,1850 (TM)
around Britain, warm Oceans
- Trimosinidae* Saidova,1981
- Mimosina* Millett,1900 (TM) *Mimosina histrix* Millett,1900
- Pavoninidae* Eimer & Fickert,1899
- Paronina* d'Orbigny,1826 (TM) *Pavonina flabelliformis* d'Orbigny,1826
= *Valvopavonina* Hofker,1951
- Furcenkoinaceae* Loeblich & Tappan,1961
- Furcenkoinidae* Loeblich & Tappan,1961
- Coryphostoma* Loeblich & Tappan,1962 (TM) *Bolivina plaitum* Carsey,1926
= *Laterostoma* Hamaoui,1979 (TM) *Laterostoma neumanniae* Hamaoui,1979
= *Laterostoma* Hamaoui,1975 (name not available)
- Furcenkoina* Loeblich & Tappan,1961 (TM) *Virgulina squammosa* d'Orbigny,1826 (Nom. cons. ICZN Op. 1834)
= *Virgulina* d'Orbigny,1826, non Bory de St. Vincent,1823
= *Neobuliminoides* McCulloch,1977 (TM) *Neobuliminoides cedrosensis* McCulloch,1977
- F. skagerakensis* (Höglund,1947)
= *Virgulina skagerakensis* Höglund,1947
* Correct genus?
Gullmarfjord, Koster Channel, Skagerrak, Oslofjord
- F. fusiformis* (Williamson,1858)
= *Bulmina pupoides* d'Orbigny,1846 var. *fusiformis* Williamson,1858
= *Bulmina* "fusiformis" : Höglund,1947
= *Virgulina* "fusiformis" : Auct.
= *Stainforthia* "fusiformis" : Auct.
Öresund, Kattegatt, Gullmarfjord, Skagerrak, Oslofjord, S Norway, central North Sea, Iceland, Barents Sea, W Scotland, SW Ireland, Bristol Channel, Celtic Sea, Engl. Channel, Arcachon, Biscay
- F. obscura* (Goës,1894)
= *Virgulina obscura* Goës,1894
* Correct genus?
Norway, North Sea
- F. sp. A* (Haynes,1973)
= *Stainforthia* sp. A : Haynes,1973
= *Virgulina schreibersiana* : Auct. (e.g. Goës,1894) non Czjzek,1848
Bohusl, Oslofjord, Bergen, Spitsbergen, Wales, Belgium, Engl. Channel, Arctic Ocean
- F. compressa* (Bailey,1894)
= *Bulmina compressa* : Terquem & Terquem,1886
* Correct genus???
Iceland, Arctic Ocean
- Rutherfordoides* McCulloch,1981 (TM) *Rutherfordia rotundiformis* McCulloch,1977
= *Rutherfordia* McCulloch,1977, non Macgillivray,1921
= *Hasolina* Nomura,1983 (TM) *Virgulina mexicana* Cushman,1922
- Sigmavirgulina* Loeblich & Tappan,1957
- S. tortuosa* (Brady,1881)
= *Bolivina tortuosa* Brady,1881 (TM)
Britain, Mediterranean, Cape Verde
- Virgulinellidae* Loeblich & Tappan,1984
- Virgulinella* Cushman,1932 (TM) *Virgulina pertusa* Reuss,1861
- Delosinacea* Parr,1950
- Caucasinidae* N.K. Bykova,1959
- Caucasininae* N.K. Bykova,1959
- Francesita* Loeblich & Tappan,1963 (TM) *Virgulina?* *advena* Cushman,1922
- Pleurostomellaceae* Reuss,1860
- Pleurostomellidae* Reuss,1860
- Pleurostomellinae* Reuss,1860
- Ellipsoglandulina* A. Silvestri,1900 (TM) *Ellipsoglandulina laevigata* A. Silvestri,1900
- Nodosarella* Rzehak,1895 (TM) *Lingulina tuberosa* Gümbel,1870
= *Ellipsonodosaria* A. Silvestri,1900 (TM) *Lingulina rotundata* d'Orbigny,1846
- Pleurostomella* Reuss,1860
= *Pleurostomellina* Schubert,1911 (TM) *Pleurostomella barroisi* Berthelin,1880 = *Pleurostomella fusiformis* Reuss,1860
= *Ellipsonodosaria* (*Ellipsoglandulina*) Franke,1928 (TM) *Dentalina subnodosa* Reuss,1851
= *Clarella* Fuchs,1967 (TM) *Nodosarella articulata* Brotzen,1936
- P. subnodosata* (Reuss,1851)
= *Dentalina nodosa* d'Orbigny : Reuss,1846, non d'Orbigny,1840
= *Dentalina subnodosata* Reuss,1851 (TM)
warm Faeroe Channel
- Stilostomellacea* Finlay,1947
- Stilostomellidae* Finlay,1947
- Nodogenerina* Cushman,1927 (TM) *Nodogenerina bradyi* Cushman,1927
- Orthomorphina* Stainforth,1952 (TM) *Nodogenerina havanensis* Cushman & Bermúdez,1937
- Siphonodosaria* A. Silvestri,1924 (TM) *Nodosaria abyssorum* Brady,1881
= *Sagrimonodosaria* Jedlicksha,1931
- Strictocostella* Patterson,1987 (TM) *Ellipsonodosaria modesta* Bermúdez var. *prolata* Cushman & Bermúdez,1937
- Discorbacea* Ehrenberg,1838
- Placentulinidae* G.K. Kasimova, Poroshina & Geodakchan,1980
- Ashbrookiinae* Loeblich & Tappan,1984
- Patellinella* Cushman,1928 (TM) *Textularia inconspicua* Brady,1884
- Bagginidae* Cushman,1927
- Baggininae* Cushman,1927
- Baggina* Cushman,1926 (TM) *Baggina californica* Cushman,1926
- B. hauerii* (d'Orbigny,18??)
= *Pulvinulina hauerii* d'Orbigny,18??
?Sussex
- Cancris* de Montfort,1808
= *Rotalina* d'Orbigny,1839 (TM) *Rotalina sagra* d'Orbigny,1839

- C. auricula** (Fichtel & Moll,1798)
 = *Nutilus auricula* Fichtel & Moll,1798
 = *Cancris auriculatus* de Montfort,1808 (TM)
 Christianssand, Stavanger, Bergen, Lofoten, Barents Sea, Iceland, Celtic Sea, Bristol Channel, Engl. Channel, Arcachon, Biscay, Galicia, Canaries
- C. oblongus** (Williamson,1858)
 = *Rotalina oblonga* Williamson,1858
 = *Pulvinulina auricula* : Goës,1894 ; Flint,1899, non (Fichtel & Moll,1798)
 = *Cancris auricula* : Cushman,1931, non (Fichtel & Moll,1798)
 North Sea, Arctic Seas, NW Scotland, Wales, Ireland, Celtic Sea, Plymouth, Biscay
- Valvularia** Cushman,1926 (TM *Valvularia californica* Cushman,1926)
- Eponidiidae** Hofker,1951
- Eponidiace** Hofker,1951
- Eponides** de Montfort,1808
 = *Pulvinulus* de Lamarck,1816
 = *Pulvinulina* Parker & Jones, in Carpenter, Parker & Jones,1862
 = *Eponidopsis* Reiss,1960 (TM *Eponides lorenensis* Finlay,1939)
 = *Criboeponides* Shchedrina,1964 (TM *Poroeponides criborepandus* Asano & Uchio, in Asano,1951)
 = *Podoella* Pishvanova,1972 (TM *Eponides probatus* Krasheninnikov,1958)
- E. repandus** (Fichtel & Moll,1798)
 = *Nautius repandus* Fichtel & Moll,1798 (TM)
 = *Placentula pulvinata* Lamarck,1822 (TM *Placentula* de Lamarck,1822)
 NE Scotland, Orkneys, Mediterranean, Azores
- E. repandus** (Fichtel & Moll,1798) *concameratus* (Montagu,1808)
 = *Serpula concamerata* Montagu,1808
 Faeroe Channel, Iceland, Shetlands, Hebrides, W Scotland, Ireland, Wales, Barents Sea, S North Sea, Belgium, Engl. Channel, Arcachon, Biscay
- E. karsteni** (Reuss,1855)
 = *Rotalia karsteni* Reuss,1855
 = *Pulvinulina karsteni* : Brady,1881
 * Correct genus?
 Plymouth, Barents Sea, Novaja Zemlja, Franz Josefs Land
- E. nitidula** (Chaster,1892)
 = *Pulvinulina nitidula* Chaster,1892
 = *Prinella nitidula* : Murray,1965
 * Correct genus?
 Engl. Channel
- E. concentrica** (Parker & Jones MS, in Brady,1864)
 = *Pulvinulina concentrica* Parker & Jones MS, in Brady,1864
 * Correct genus??
 Bergen, Plymouth
- E. umbrata** (Reuss,18??)
 Iceland
- Ioanella** Saidova,1975 (TM *Truncatulina tumidula* H.B. Brady,1884)
 = *Ioanella* Khusid,1973 (name not available)
- Poroeponides** Cushman,1944 (TM *Rosalina lateralis* Terquem,1878)
- Sestronophorinae** Saidova,1981
- Planopulvinulina** Schubert,1921 (TM *Pulvinulina dispansa* Brady,1884)
- Mississippiidae** Saidova,1981
- Mississippiinae** Saidova,1981
- Mississippiina** Howe,1930 (TM *Mississippiina monsouri* Howe,1930)
- Discorbidae** Ehrenberg,1838
- Discorbe** de Lamarck,1804 (TM *Discorbites vesicularis* de Lamarck,1804)
 = *Discorbites* de Lamarck,1804
- D. vilardeboana** (d'Orbigny,1839)
 = *Rosalina vilardeboana* d'Orbigny,1839
 = *Discorbia vilardeboana* : Auctt.
 Bergen, Plymouth
- D. australis** Parr,19?? : Gabel,1971
 North Sea
- D. floridiana** Cushman,19?? : Gabel,1971
 ?= *D. floridiana* Cushman,1922
 North Sea, Florida, West India
- D. araucana** (d'Orbigny,18??)
 * Correct genus??
 ?Zuiderzee, ?W Ireland, Chile
- D. obtusa** (d'Orbigny,18??)
 * Correct genus??
 Oslofjord, Britain
- D. biconcava** (Parker & Jones,18??)
 * Correct genus?
 ?Wales
- D. peruviana** d'Orbigny,18??
 * Correct genus?
 ?Brittain, Peru
- D. polypraphes** (Reuss,18??)
 * Correct genus??
 ?Brittain, Peru
- Neopeponides** Reiss,1960 (TM *Rotalina schreibersii* d'Orbigny,1846)
 = *Cyclospira* Eimer & Fickert,1899, non Hall & Clark,1894 (TM *Rotalina schreibersii* d'Orbigny,1846)
 = *Rotobis* Sellier de Civrieux,1977 (TM *Rosalina auberti* d'Orbigny,1839 = *Discorbia mira* Cushman,1922)
- Trochulina** d'Orbigny, in Ehrenberg,1839 (TM *Rotalia turbo* d'Orbigny,1826)
 = *Discorbia* Parker & Jones, in Carpenter & al.,1862 (TM *Rotalia turbo* d'Orbigny,1826)
 = *Rotorbinella* Bandy,1944 (TM *Rotorbinella colliculus* Bandy,1944)
 = *Lamellocorbus* Bermúdez,1952 (TM *Discorbia dimidiata* Jones & Parker, in Carpenter & al.,1862)
- T. turbo** (d'Orbigny,1826)
 = *Rotalia turbo* d'Orbigny,1826
 = *Discorbia turbo* : Auctt.
 Plymouth
- T. chasteri** (Heron-Allen & Earland,1913)
- = *Discorbia minutissima* Chaster,1892, non Seguenza,18??
 = *Discorbia chasteri* Heron-Allen & Earland,1913
 * Correct genus??
 off Faeroes, Plymouth
- T. chasteri** (Heron-Allen & Earland,1913), var. *bispinosa* (Heron-Allen & Earland,1913)
 = *Discorbia chasteri* Heron-Allen & Earland,1913, var. *bispinosa* Heron-Allen & Earland,1913
 off Faeroes, W Scotland, W Ireland, S Cornwall
- T. tabernacularis** (Brady,1879) : Heron-Allen & Earland,1930
 = *Discorbia tabernacularis* Brady,1879
 * Correct genus??
 Plymouth, Cape Verde - southwards
- T. lauriei** Heron-Allen & Earland,1924
 = *Discorbia tabernacularis* : Sidebottom,1910, non Brady,1879
 * Correct genus??
 Plymouth, Mediterranean
- T. rosacea** (d'Orbigny,1826)
 = *Rotalina rosacea* d'Orbigny,1826
 = *Discorbia rosacea* : Auctt.
 * Correct genus??
 Plymouth
- T. plana** Egger,1893
 Helgoland
- Rosalinidae** Reiss,1963
- Gavelinopsis** Hofker,1951
- G. praegeri** (Heron-Allen & Earland,1913)
 = *Discorbia praegeri* Heron-Allen & Earland,1913 (TM)
 * probably clinging to pebbles in current-swept areas
 W Ireland, around Scotland, Celtic Sea, Bristol Channel, Wales, Engl. Channel, Biscay
- Hyrrokin** Cedhagen,1992
- H. sarcophaga** Cedhagen,1992
 = *Pulvinulina punctulata* : Auctt., non = *Rotalia punctulata* d'Orbigny,1826
 = *Eponides punctulata* : Auctt., non (d'Orbigny,1826)
 @ *Isops*, *Geodia*, *Stellata*, *Aestia*, *Lophelia* etc.
 Bohusl., Norway, North Sea, Faeroes, Britain, Biscay
- Neoconorbina** Hofker,1951
- N. terquemi** (Rzehak,1888)
 = *Discorbia terquemi* Rzehak,1888
 = *Rosalina orbicularis* Terquem,1876, non d'Orbigny,1850 (TM)
 Helgoland, Plymouth
- N. neapolitana** Hofker,1951
 = *Rosalina neapolitana* : Haynes,1973
 W Ireland, Wales, Mediterranean
- Rosalina** d'Orbigny,1826
 = *Turbulinula* d'Orbigny,1839
 = *Pararosalina* McCulloch,1977 (TM *Pararosalina densitiva* McCulloch,1977)
- R. globularis** d'Orbigny,1826 (TM)
 = *Discorbia globularis* : Auctt.
 * clinging to firm substrates.
 Sweden, Norway, Belgium, N Kent, Ireland, Celtic Sea, Bristol Channel, Engl. Channel, N Spain
- R. globularis** d'Orbigny,1826 forma *cyprinodon* (Rhumbler,1938)
 = *Discorbis globularis* (d'Orbigny,1826) forma *cyprinodon* Rhumbler,1938
 * Correct genus?
 Helgoland
- R. globularis** d'Orbigny,1826, var. *anglica* (Cushman,1931)
 = *Discorbs globularis* (d'Orbigny,1826), var. *anglica* Cushman,1931
 = *Discorbia irregularis* Heron-Allen & Earland,1913, non Rhumbler,1906
 NE Scotland
- R. anomala** Terquem,1875
 = *R. globularis* ; Voortuyn,1858 ; Haake,1962, non d'Orbigny,1826
 = *Discorbia globularis* : Goës,1894, non d'Orbigny,1826
 North Sea, Arctic, NE Scotland, Ireland, Wales, Engl. Channel
- R. columbiensis** (Cushman,1925)
 = *Discorbs columbiensis* Cushman,1925
 Oslofjord
- R. concinna** (Brady,1884)
 = *Discorbia concinna* Brady,1884
 ?Netherlands, Engl. Channel
- R. globulus** d'Orbigny,1826
 Engl. Channel
- R. williamsoni** (Chapman & Parr,1932)
 = *Discorbs williamsoni* Chapman & Parr,1932
 = *Rotalina nitida* Williamson,1858, non Reuss,1844
 = *Discorbs nitida* : Cushman,1931
 * Should perhaps be moved to *Neoconorbina* Hofker?
 Oslofjord, German North Sea, Netherlands, Belgium, Faeroes, Iceland, all around Britain, Galicia, Mediterranean
- R. irregularis** (Rhumbler,1906)
 = *Discorbia irregularis* Rhumbler,1906
 W Ireland, Engl. Channel
- R. cf. bradyi** (Cushman,1915) : Haynes,1973
 Wales
- R. cf. mediterranensis** d'Orbigny,1826 : Murray,1965
 Engl. Channel
- R. sp.** : Gabel,1971
 German Bight
- Rotorboidea** Sellier de Civrieux,1977 (TM *Discorbia valvulata* (d'Orbigny) var. *granulosa* Heron-Allen & Earland,1915)
- Tretomphalus** Möbius,1880 (TM *Rosalina bulboides* d'Orbigny,1839)
- Bronnimannia** Bermúdez,1952 (TM *Discorbs palmerae* Bermúdez,1935)
 = *Neobronnimannia* McCulloch,1977 (TM *Neobronnimannia compacta* McCulloch,1977)
- Rotaliellidae** Loeblich & Tappan,1964

- Metarotaliella*** Grell,1962
M. parva Grell,1962 (TM)
 France
- Rotaliella*** Grell,1954 (TM) *Rotaliella heterocaryotica* Grell,1954
- R. roscoffensis*** ?Grell,195?
 Roscoff
- Sphaeroidinidae** Cushman,1927
- Sphaeroidina*** d'Orbigny,1826
 = *Sexloculina* Czjzek,1848 (TM) *Sexloculina haueri* Czjzek,1848
- S. bulloides*** d'Orbigny,1826 (TM)
 Bergen
- Glabratellacea** Loeblich & Tappan,1964
- Glabratellidae** Loeblich & Tappan,1964
- Discorbinoidea*** Saidova,1975 (TM) *Discorbinoidea subpatelliformis* Saidova,1975
 = *Sabinoides* McCulloch,1977 (TM) *Sabinoides densiformis* McCulloch,1977
- Glabrarella*** Dorreen,1948 (TM) *Glabrarella crassa* Dorreen,1948
 = *Glabroralisina* McCulloch,1977 (TM) *Glabroralisina distincta* McCulloch,1977
- G. milletti*** (Wright,1911)
 = *Discorbina milletti* Wright,1911
 = *Rosalina milletti* : Haynes,1973
 = *Discorbis orbicularis* (Terquem) var. *selseyenensis* (Heron-Allen & Earland,1911) : Cushman,1949
 * clinging to firm substrates. Correct genus?
 German North Sea, W Scotland, W Ireland, Irish Sea, Belgium, Engl. Channel, Wales
- G. wrightii*** (Brady,1881)
 = *Discorbina parisiensis* : Wright,1877 (p.p. - plate iv, fig. 2), non (d'Orbigny,1826)
 = *Discorbina wrightii* Brady,1881
 = *Discorbina bacata* Heron-Allen & Earland,1913
 = *Discorbis (Glabrarella) wrightii* : Haynes,1973
 Novaja Zemlja, W Scotland, Ireland, Wales, Engl. Channel
- G. parisiensis*** (d'Orbigny,1826)
 = *Rosalina parisiensis* d'Orbigny,1826
 = *Discorbina parisiensis* : Wright,1877 (p.p. - plate iv, fig. 1)
 * Correct genus?
 Belgium, Plymouth
- Glabratellina*** Seiglie & Bermúdez,1965 (TM) *Glabratellina arcuata* Seiglie & Bermúdez,1965
 = *Crumia* McCulloch,1977 (TM) *Crumia albida* McCulloch,1977
 = *Sabina* McCulloch,1977 non Parona,1909 (TM) *Sabina turriformis* McCulloch,1977
 = *Sabina* McCulloch,1981 (TM) *Sabina turriformis* McCulloch,1977
- Heronallenidae Loeblich & Tappan,1986
- Heronallenia*** Chapman & Parr,1931 (TM) *Discorbina wilsoni* Heron-Allen & Earland,1922
 = *Metatheronallenia* McCulloch,1977 (TM) *Metatheronallenia rugosiformis* McCulloch,1977
 = *Neotheronallenia* McCulloch,1977 (TM) *Neotheronallenia craigi* McCulloch,1977
 = *Planoheronallenia* McCulloch,1977 (TM) *Planoheronallenia wenmanni* McCulloch,1977
 = *Subheronallenia* McCulloch,1977 (TM) *Subheronallenia crosbyi* McCulloch,1977
 = *Guyhotytina* McCulloch,1981 (TM) *Guyhotytina helena* McCulloch,1981
- Buliminoididae** Seiglie,1970
- Buliminoides*** Cushman,1911 (TM) *Bulimina williamsoniana* Brady,1881
- Siphoninacea** Cushman,1927
- Siphoninidae** Cushman,1927
- Siphonininae** Cushman,1927
- Siphonina*** Reuss,1850 (TM) *Siphonina fimbriata* Reuss,1850 = *S. reticulata* (Czjzek,1848)
- S. georgiana*** Haynes,1973
 = *Truncatulina reticulata* : Brady,1884 ; Chaster,1892, non (Czjzek,1848)
 = *Siphonina tubulosa* : Heron-Allen & Earland,1930 non Cushman,1927
 = *Siphonina reticulata?* : Cushman,1931, non (Czjzek,1848)
 Belgium, Wales, Engl. Channel
- Discorbinellacea** Sigal,1952
- Parreloididae** Hofker,1956
- Cibicidoides*** Thalmann,1939 (TM) *Truncatulina mundula* Brady, Parker & Jones,1890
 = *Dendrina* Costa MS, in Fornasini,1898, non Quenstedt,1848 (TM) *Dendrina succinea* Costa MS, in Fornasini,1898
- Pseudoparrellidae** Voloshinova,1952
- Pseudoparrellinae** Voloshinova,1952
- Alabaminoidea*** Gudina & Saidova,1967 (TM) *Alabamina mitis* Gudina,1966
- Epistominella*** Husezima & Maruhasi,1944 (TM) *Epistominella pulchella* Husezima & Maruhasi,1944
- E. vitrea*** Parker, in Parker, Phleger & Peirson,1953
 NW Scotland, Celtic Sea, Bristol Channel, Engl. Channel
- E. exigua*** (Brady,1884)
 = *Pulvinulina exigua* Brady,1884
 Oslofjord, Spitsbergen
- E. narrensis*** (Kuwano,1950)
 = *Pseudoparrella narrensis* Kuwano,1950
 = *Pulvinulina exigua* : Hofker,1951, non Brady,1884
 Wales, Japan
- Pseudoparrella*** Cushman & ten Dam,1948 (TM) *Pulvinulina subperuviana* Cushman,1926
 = *Pulvinulina* Cushman,1926, non Eimer & Fickert,1899
 = *Peschongia* McCulloch,1977 (TM) *Peschongia bikiniensis* McCulloch,1977
- Stetsonia*** Saidova,1981
- Stetsonia*** F.L. Parker,1954 (TM) *Stetsonia minuta* F.L. Parker,1954
- Discorbinellidae** Sigal,1952
- Discorbinellinae** Sigal,1952
- Discorbinella*** Cushman & Martin,1935 (TM) *Discorbinella montereyensis* Cushman & Martin,1935
 = *Discopulvinulina* Hofker,1951 (TM) *Rosalina bertheloti* d'Orbigny,1839
 = *Discorbinellopsis* McCulloch,1977 (TM) *Discorbinellopsis symmetrica* McCulloch,1977
- = *Discorbinoides* McCulloch,1977, non Saidova,1975 (TM) *Discorbinoides chincaensis* McCulloch,1977
 = *Neoplanoiscoris* McCulloch,1977 (TM) *Neoplanoiscoris galapagensis* McCulloch,1977
 = *Discorbinella* McCulloch,1981 (TM) *Discorbinoides chincaensis* McCulloch,1977
- D. berthelothiana*** (Parker & Jones,1865)
 off Ireland
- D. sp. A. aff bertheloti*** (d'Orbigny,1839) : Haynes,1973
 Wales
- D. bertheloti*** (d'Orbigny,1839)
 = *Rosalina bertheloti* d'Orbigny,1839
 = *Discorbinus berthelothi?* : Nørvang,1941
 Bergen, Vesteraalen, Shetlands, Iceland, W Scotland, Ireland, Novaja Zemlja, Canaries
- Discorbitina*** Sellié de Civrieux,1977
- D. pustulata*** (Heron-Allen & Earland,1913)
 = *Discorbitina pustulata* Heron-Allen & Earland,1913 (TM)
 Plymouth
- Laticarinina*** Galloway & Wissler,1927
 = *Carinina* Galloway & Wissler,1927, non Hubrecht,1887
 = *Paricarinina* Finlay,1940 (TM) *Truncatulina temuiargo* Brady var. *altocamerata* Heron-Allen & Earland,1922
 = *Mesocarinina* McCulloch,1977 (TM) *Mesocarinina velata* McCulloch,1977
 = *Metacarinina* McCulloch,1977 (TM) *Metacarinina inferillata* McCulloch,1977
- L. pauperata*** (Parker & Jones,1865)
 = *Pulvinulina repanda* (Fichtel & Moll) var. *menardi* d'Orbigny subvar. *pauperata* Parker & Jones,1865 (TM)
 N Atlantic (56° N - southwards)
- Planorbulinacea** Schwager,1877
- Planulinidae** Bermúdez,1952
- Hyalinea*** Hofker,1951
 = *Hofkerinella* Bermúdez,1952
- H. balthica*** (Schröter,1783)
 = *Nautilus balthicus* Schröter,1783 (TM)
 = *Anomalina balthica* : Cushman,1931
 = *Operculina ammonoides* : Auctt., non Gronovius,1???
 * very common in W Norway
 Baltic, Oslofjord, S Norway, Bergen, Vadsø, central North Sea, Iceland, Barents Sea, Shetlands, Orkneys, Hebrides, NW Scotland, Ireland, Celtic Sea, western approaches to the Engl. Channel, Biscay, Galicia, Canaries
- Planulina*** d'Orbigny,1826
- P. ariminensis*** d'Orbigny,1826 (TM)
 N Atlantic at 150-1600 fathoms, off Brazil
- P. sp.*** : Murray,1965
 Engl. Channel
- Cibicididae** Cushman,1927
- Cibicididae** Cushman,1927
- Cibicides*** de Montfort,1808 (TM) *Cibicides refulgens* de Montfort,1808
 = *Storilus* de Montfort,1808 (TM) *Storilus radiatus* de Montfort,1808
 = *Truncatulina* d'Orbigny,1826
- C. pseudoungerianus*** (Cushman,1922)
 = *Truncatulina pseudoungeriana* Cushman,1922
 = *Truncatulina ungeriana* : Brady,1884, non *Rotalina ungeriana* d'Orbigny,1826
 * clinging to firm substrates.
 * Is this not a *Lobatula* Fleming?
 Oslofjord, Bergen, Barents Sea, Shetlands, off W & SW Ireland, Belgium, Celtic Sea, Arcachon, Biscay, Galicia
- C. fletcheri*** Galloway & Wissler,1927
 * Correct genus?
 Iceland, Faeroes, Wales
- C. fletcheri*** Galloway & Wissler,1927 var. *sachalinica* Vasilenko,1954
 ?= *Planorbulina akneriana* : Goës,1894, non (d'Orbigny,1846)
 * Correct genus?
 ?Bohusl, ?S Norway, ?Hardanger, Wales
- C. advenum*** (d'Orbigny,1838)
 = *Truncatulina advena* d'Orbigny,1839
 Plymouth
- C. refulgens*** de Montfort,1808
 along Norway - Vadsø, Spitsbergen, Barents Sea, Arctic Ocean, off W Scotland, W Ireland, Plymouth, Iceland
- C. akneriana*** (d'Orbigny,1846)
 = *Rosalina akneriana* d'Orbigny,1846
 * Correct genus?
 Plymouth
- C. pygmaea*** (Hantken,1875)
 = *Truncatulina pygmaea* Hantken,1875
 = *Pulvinulina pygmaea* Hantken,1875
 * Correct genus?
 Plymouth
- C. ungerianus*** (d'Orbigny,1846)
 = *Rotalina ungeriana* d'Orbigny,1846
 * Correct genus?
 Plymouth
- C. perlucina*** Nuttal,1??? : Gabel,1971
 North Sea
- C. irregularis*** (Terquem & Terquem,1886)
 = *Anomalina irregularis* Terquem & Terquem,1886
 * Correct genus???
- C. hemisphaerica*** (Terquem,18???)
 = *Anomalina hemisphaerica* Terquem,18???
- C. limbata*** (Terquem,18???)
 = *Anomalina limbata* Terquem,18???
- C. nodulosa*** (Terquem,18???)
 = *Anomalina nodulosa* Terquem,18???
- * Correct genus???
- Dunkerque

- C. scutellata** (Terquem,18??)
= *Anomalina scutellata* Terquem,18??
* Correct genus???
Dunkerque
- C. simplex** (Terquem,18??)
= *Anomalina simplex* Terquem,18??
* Correct genus???
Dunkerque
- C. boueana** (d'Orbigny,1846)
= *Truncatulina boueana* d'Orbigny,1846
* Correct genus?
Iceland
- C. tuberculata** (d'Orbigny,1826)
= *Truncatulina tuberculata* d'Orbigny,1826
* Correct genus???
Iceland, European coasts, Mediterranean
- C. globulosa** (Terquem & Terquem,1886)
= *Truncatulina globulosa* Terquem & Terquem,1886
* Correct genus???
Iceland
- C. rotundatus** Shchedrina,19??
Barents sea
- C. shokalskii** Shchedrina,19??
Barents Sea
- Cibicidina** Bandy,1949 (TM *Cibicidina walli* Bandy,1949)
- Fontbotia** González-Donoso & Linares,1970
- F. wuellerstorfi** (Schwager,1866)
= *Anomalina wuellerstorfi* Schwager,1866 (TM)
= *Planulina wuellerstorfi* (Schwager,1866) : Gabel,1971
off Bergen, along N Norway, Iceland
- Lobatula** Fleming,1828
- L. lobatus** (Walker & Jacob, in Kanmacher,1798)
= *Nautilia lobatus* Walker & Jacob, in Kanmacher,1798
= *Serpula lobata* Montagu,1803
= *Lobatula vulgaris* Fleming,1828 (TM)
= *Cibicides lobatulus* : Auctt.
* clings to seaweeds, tunicates, dead bryozoan colonies & other firm substrates.
Belt Sea, Øresund, Bohusl., Oslofjord, entire Norway, Spitsbergen, Barents Sea, Novaja Zemlya, Belgium, Iceland, Faeroes, Ireland, Engl. Channel, Celtic Sea, Orkneys, Arcachon, Biscay, Galicia
- Stichocibicidina** Saidova,1981
- Dycocibicides** Cushman & Valentine,1930 (TM *Dycocibicides biserialis* Cushman & Valentine,1930)
= *Rectococbicidella* McLean,1956 (TM *Rectococbicidella robertsi* McLean,1956)
- D. biserialis** Cushman & Valentine,1930 : Gabel,1971
North Sea
- D. uniserialis** Thalmann,18?? : Gabel,1971
North Sea
- Annulocibicidinae** Saidova,1981
- Cyclocibicides** Cushman,1927
- C. vermiculata** (d'Orbigny,1826)
= *Planorbilina vermiculata* d'Orbigny,1826 (TM)
Sussex, W Ireland, Mediterranean
- Planorbulinidae** Schwager,1877
- Caribeannelina** Saidova,1981
- Caribeannella** Bermúdez,1952 (TM *Caribeannela polystoma* Bermúdez,1952)
= *Oinomikadoina* Matsunaga,1954 (TM *Oinomikadoina ogiensis* Matsunaga,1954)
= *Pseudococbicidoides* Ujiiie,1956 (TM *Pseudococbicidoides katasensis* Ujiiie,1956)
- Planorbulininae** Schwager,1877
- Planorbulina** d'Orbigny,1826 (TM *P. mediterranensis* d'Orbigny,1826)
= *Asterodiscus* Ehrenberg,1839 (TM *Asterodiscus forskali* Ehrenberg,1839 = *Planorbulina mediterranensis* d'Orbigny,1826)
= *Spirorboprys* Ehrenberg,1844 (TM *Spirorboprys aegaea* Ehrenberg,1844 = *Planorbulina mediterranensis* d'Orbigny,1826)
= *Cibicidella* Cushman,1927 (TM *Truncatulina variabilis* d'Orbigny,1826)
- P. distoma** Terquem,1876
= *P. mediterranensis* : Auctt., non d'Orbigny,1826 (TM)
= *Planorbulina vulgaris* : Williamson,1858, non d'Orbigny,1839
* clinging to firm substrates.
W Sweden, along Norway - Tromsø, Iceland, E & W North Sea, Belgium, Barents Sea, Orkneys, Engl. Channel, Bristol Channel, Wales, Ireland, Celtic Sea, Arcachon, Biscay, Galicia
- P. variabilis** (d'Orbigny,1826)
= *Truncatulina variabilis* d'Orbigny,1826
Plymouth, Iceland, Mediterranean, Canaries
- P. sp. A** : Haynes,1973
Wales
- Planorbulinella** Cushman,1927 (TM *Planorbulina vulgaris* d'Orbigny var. *larvata* Parker & Jones,1865)
- Cymbaloporidae** Cushman,1927
- Cymbaloporinae** Cushman,1927
- Cymbaloporella** Cushman,1928 (TM *Rosalina squamosa* d'Orbigny,1839)
= *Pseudotretomphalus* Hofker,1979 (TM *Tretomphalus bulloides* (d'Orbigny) var. *plana* Cushman,1924)
- Victoriellidae** Chapman & Crespin,1930
- Rupertininae** Loeblich & Tappan,1961
- Rupertina** Loeblich & Tappan,1961
= *Rupertia* Wallich,1877, non Gray,1865 (TM *Rupertia stabilis* Wallich,1877)
- R. stabilis** (Wallich,1877)
= *Rupertia stabilis* Wallich,1877 (TM)
Christiansund, Spitsbergen, Iceland, Rockall Bank, Biscay
- Acervulinacea** Schultze,1854
- Acervulinidae** Schultze,1854
- Acervulina** Schultze,1854
= *Aphrosina* H.J. Carter,1879 (TM *Aphrosina informis* H.J. Carter,1879)
- A. inhaerens** Schultze,1854 (TM)
= *Gypsinia inhaerens* : Auctt.
* encrusts bryozoan colonies.
Bohusl., ?Norway, Belgium, Engl. Channel, NE & W Scotland, Orkneys
- Gypsinia** H.J. Carter,1877 (TM *Polytrema planum* H.J. Carter,1876)
- G. vesicularis** (Parker & Jones,1860)
= *Orbitalina vesicularis* Parker & Jones,1860
?Norway, W Ireland
- G. vesicularis** (Parker & Jones,1860), var. *intermedia* Goës,1894
Bossl., Norway
- Sphaerogypsina** Galloway,1933 (TM *Ceriopora globulus* Reuss,1848 = *Chaetetes pygmaeus* Reuss,1848 = *Tinoporus fuchsii* Karrer,1877)
= *Excentrogypsina* Popescu,1975 (TM *Tinoporus fuchsii* Karrer,1877)
- Asterigerinacea** d'Orbigny,1839
- Epistomariidae** Hofker,1954
- Epistomariinae** Hofker,1954
- Pseudoeponides** Uchio, in Kawai & al.,1950 (TM *Pseudoeponides japonicus* Uchio, in Kawai & al.,1950)
- Nuttallidiinae** Saidova,1981
- Nuttallides** Finlay,1939 (TM *Eponides truempyi* Nuttall,1930)
= *Osangularia* Saidova,1975 (TM *Eponides bradyi* Earland,1934)
- Asterigerinatidae** Reiss,1963
- Asterigerinata** Bermúdez,1949 (TM *Asterigerinata dominicana* Bermúdez,1949)
- A. mamilla** (Williamson,1858)
= *Rotalina mamilla* Williamson,1858
= *Discorbina mamilla* : Auctt.
Wadden Sea, Shetlands, NW Scotland, Ireland, Bristol Channel, Wales, Engl. Channel, Arcachon, N Spain, Mediterranean
- Biasterigerina** Seiglie & Bermúdez,1965
- B. planorbis** (d'Orbigny,1846)
= *Asterigerina planorbis* d'Orbigny,1846 (TM)
Plymouth
- Eoeponiella** Wickenden,1949 (TM *Eoeponiella linki* Wickenden,1949)
= *Heminwayia* Bermúdez,1951 (TM *Discorbis multisectus* Galloway & Heminway,1941)
= *Astrellina* Anderson,1963 (*Pninaella?* *pulchella* F.L. Parker,1952)
- E. pulchella** (F.L. Parker,1952)
= *Pninaella?* *pulchella* F.L. Parker,1952
Belt Sea
- Asterigerinidae** d'Orbigny,1839
- Asterigerina** d'Orbigny,1839 (TM *Asterigerina carinata* d'Orbigny,1839)
- Amphisteginidae** Cushman,1927
- Amphistegina** d'Orbigny,1826 (TM *Amphistegina quoyii* d'Orbigny,1826 = *Nautilus radiatus* Fichtel & Moll,1798)
= *Omphalophacus* Ehrenberg,1839 (TM *Omphalophacus hemprichi* Ehrenberg,1840)
- "**A. radiata** Terquem,1881", non Fichtel & Moll,1798
Dunkerque
- A. vulgaris** d'Orbigny,18?? : Terquem,1881"
Dunkerque
- Nonionacea** Schultze,1854
- Nonionidae** Schultze,1854
- Nonioninae** Schultze,1854
- Haynesina** Banner & Culver,1978
- H. germanica** (Ehrenberg,1840)
= *Nonionina germanica* Ehrenberg,1840 (TM)
= *Nonion germanicum* : Auctt.
= *Protephilidium germanicum* : Auctt.
= *Nonion tisburyense* Butcher,1948
= *Nonion paralium* Tintant,1954
= *Protephilidium anglicum* Murray,1965
* Tolerates 1 ppt salinity. Intertidal.
Uddevalla, SE North Sea, Wadden Sea, Barents Sea, Engl. Channel, Wales, Arcachon, N Spain
- Nonion** de Montfort,1808 (TM *Nautilus faba* Fichtel & Moll,1798 - replacement type species: see ICZN ZN(S) 2225/1 - petition pending)
= *Azera* Khalilov,1958 (TM *Azera transversa* Khalilov,1958)
= *Neonomalina* McCulloch,1977 (TM *Neonomalina chincoensis* McCulloch,1977)
= *Abbottina* McCulloch,1977 (TM *Abbottina tannerbankensis* McCulloch,1977)
- N. faba** (Fichtel & Moll,1798)
= *Nautilus faba* Fichtel & Moll,1798
Plymouth
- N. depressulus** (Walker & Jacob, in Kanmacher,1798)
= *Nautilus depressulus* Walker & Jacob, in Kanmacher,1798
Wadden Sea, E Kent, NW Scotland, W. Ireland, Celtic Sea, Engl. Channel, Bristol Channel, Wales, Novaja Zemlya, Franz Josef's Land
- N. depressulum granosum** (d'Orbigny,18??)
= *Nonionina granosa* d'Orbigny,18??
* Correct genus???
Helgoland
- N. zaandamae** (van Vorthuysen,1952)
= *Anomalinoidea barleeanum* (Williamson) var. *zaandamae* van Vorthuysen,1952
= *Nonion pomplioides* : Cushman,1930, non (Fichtel & Moll,1798)
= *Nonion barleeanum* : Auctt., non *Nonionina barleeanum* Williamson,1858
= *Nonion barleeanum* (Williamson) var. *inflatum* van Vorthuysen,1950, non *Nonionina inflata* Alth,1850, nec *Anomalinoidea barleeanum* (Williamson) var. *inflatum* (van Vorthuysen) van Vorthuysen,1950
Bergen, NE, N & NW Greenland, Barents Sea

- N. (Florilus) pauperatum* (Balkwill & Wright,1885)
 = *Nonionina pauperata* Balkwill & Wright,1885
 German North Sea, Netherlands, Wales, Ireland, W Scotland, Engl. Channel
- N. ornatissimum* Cushman,19?? : Gabel,1971
 North Sea
- N. crassulum* (Walker & Jacob, in Kanmacher,1798)
 = *Nautilus crassulus* Walker & Jacob, in Kanmacher,1798
 * Correct genus??
 N Kent, ?Iceland
- Nonionella* Cushman,1926 (TM *Nonionella miocenica* Cushman,1926)
 = *Ziesenhenneia* McCulloch,1977 (TM *Ziesenhenneia simplex* McCulloch,1977)
- N. auricula* Heron-Allen & Earland,1930
 Oslofjord, N Kent, Netherlands, Plymouth, Wales, W & NW Greenland
- N. turgida* (Williamson,1858)
 = *Rotalina turgida* Williamson,1858
 Øresund, Oslofjord, Hardangerfjord, Bergen, Lofoten, Vesterålen, Novaja Zemlya, Iceland, E Greenland, North Sea, Faeroe Channel, Barents Sea, Shetlands, Orkneys, Hebrides, Irish Sea, NW Scotland, Engl. Channel, Celtic Sea, Arcachon, Biscay, Mediterranean
- N. scapha* (Fichtel & Moll,1798)
 = *Nautilus scapha* Fichtel & Moll,1798, non de Wulfen,1791
 = *Nontonia scapha* : Brady,1881
 North Sea, Iceland, Novaja Zemlya
- N. turgida* (Williamson,1858), var. *digitata* Nørvang,1945
 Iceland, Wales
- N. iridea* Heron-Allen & Earland,1932
 Oslofjord
- N. (?) Nonionellina* sp. A : Haynes,1973
 Wales
- N. sloani* (d'Orbigny,18??) : Gabel,1971
 North Sea
- N. warburgi* Brotzen,19?? : Gabel,1971
 North Sea
- N. atlantica* Cushman,19?? : Gabel,1971
 North Sea
- N. extensa* (Brotzen,19??) : Gabel,1971
 North Sea
- Nonionellina* Voloshinova,1958
- N. labradorica* (J.W. Dawson,1860)
Nonionina labradorica J.W. Dawson,1860 (TM)
 = *Nonion labradoricum* : Auctt.
 Øresund, Bohusl, Oslofjord, Bergen, Barents Sea, Spitsbergen, Iceland, NE, N & NW Greenland, Novaja Zemlya
- Nonionoides* Saidova,1975 (TM *Nonionina grataloupi* d'Orbigny,1826)
- Pseudononion* Asano,1936 (TM *Pseudononion japonicum* Asano,1936)
- Astrononionina* Saidova,1981
- Astrononion* Cushman & Edwards,1937 (TM *Nonionina stelligera* d'Orbigny,1839)
- A. gallowayi* Loeblich & Tappan,1953
 = *A. stellata* Cushman & Edwards,1937 non *Nonionina stellata* Terquem,1882
 = *A. stelligerum* : Cushman,1948, non d'Orbigny,1839
 Oslofjord, Barents Sea, Spitsbergen, Iceland, NE, N & NW Greenland, Plymouth, Arctic Ocean
- A. stelligerum* (d'Orbigny,1839)
 = *Nonionina stelligera* d'Orbigny,1839
 Plymouth, Canaries, Barents Sea
- Laminononion* Hornbrook,1964 (TM *Astrononion tumidum* Cushman & Edwards,1937)
- Pulleniinae* Schwager,1877
- Melonis* de Montfort,1808
- M. pomphiloides* (Fichtel & Moll,1798)
 = *Nautilus pomphiloides* Fichtel & Moll,1798
 = *Melonis etruscus* de Montfort,1808 (TM)
 = *Nonionina barleiana* Williamson,1858
 Oslofjord, S North Sea, Iceland, Barents Sea, Shetlands, NW Scotland, Celtic Sea, western approaches to the Engl. Channel, Biscay, Galicia
- M. umbilicatum* (Montagu,1803)
 = *Nautilus umbilicatum* Montagu,1803
 = *Nonionina umbilicatula* : Brady,1881
 Plymouth, ?Iceland, Novaja Zemlya, Arctic Ocean
- P. pullenia* Parker & Jones,1862
- P. sphaeroides* (d'Orbigny,1826)
 = *Nonionina sphaeroides* d'Orbigny,1826
 = *Nonionina bulboides* d'Orbigny,1846 (TM)
 * very common in W Norway
 North Sea, Oslofjord, Bergen, Vardo, Spitsbergen, Novaja Zemlya, Faeroe Channel, Britain
- P. osloensis* Feyling-Hanssen,1954
 = *P. quinqueloba minuta* Feyling-Hanssen,1954
 Oslofjord
- P. subcarinata* (d'Orbigny,1839)
 = *Nonionina subcarinata* d'Orbigny,1839
 = *Nonionina quinqueloba* Reuss,1851
 Oslofjord, Stavanger, Bergen, Vesterålen, Iceland, warm Faeroe Channel, Shetlands, off Ireland, Engl. Channel
- Chilostomellacea* Brady,1881
- Chilostomellidae* Brady,1881
- Chilostomellinae* Brady,1881
- Allomorphinidae* Reuss,1849
- A. trigona* Reuss,1850 (TM)
 off Spitsbergen, Indo-Pacific
- Chilostomella* Reuss,1849
- C. ovoidea* Reuss,1850 (TM)
 Østerfjord, Bergen, Iceland, off Ireland
- Quadrimorphinidae* Saidova,1981
- Quadrimorphinida* Finlay,1939 (TM *Valvulina allomorphinoides* Reuss,1860)
 = *Gyromorphina* Marie,1941
- Alabaminidae* Hofker,1951
- Alabamina* Toulmin,1941 (TM *Alabamina wilcoxensis* Toulmin,1941)
 = *Eponides* Brotzen,1942 (TM *Eponides dorsoplatus* Brotzen,1940)
- Svratkina* Pokorn'y,1956 (TM *Discorbis tuberculata* (Balkwill & Wright,1885) var. *australiensis* Chapman, Parr & Collins,1934)
- S. tuberculata* (Balkwill & Wright,1885)
 = *Discorbis tuberculata* (Balkwill & Wright,1885)
 = *Discorbina tuberculata* Balkwill & Wright,1885
 * Correct genus?
 Plymouth
- Osangulariidae* Loeblich & Tappan,1964
- Osangulararia* Brotzen,1940 (TM *Osangulararia lens* Brotzen,1940)
 = *Parrella* Finlay,1939, non Ginsburg,1938 (TM *Anomalina bengalensis* Schwager,1866)
- Oridorsalidae* Loeblich & Tappan,1984
- Oridorsalis* Andersen,1961 (TM *Oridorsalis westi* Andersen,1961)
- Heterolepididae* González-Donoso,1969
- Anomalinoidea* Brotzen,1942 (TM *Anomalinoidea plummerae* Brotzen,1942 = *Anomalina pinguis* P.H. Jennings,1936)
- Gemmellidae* Vasilenko,1954 (TM *Cibicides (Gemmellidae) orcinus* Vasilenko,1954)
- Heterolepa* Franzénau,1884 (TM *Heterolepa simplex* Franzénau,1884 = *Rotalina dutemplei* d'Orbigny,1846)
 = *Pseudotruncatulina* Andreæ,1884 (TM *Rotalina dutemplei* d'Orbigny,1846)
 = *Cibusoides* Saidova,1975 (TM *Cibusoides elegans* Saidova,1975)
 = *Crawfordoides* McCulloch,1981 (TM *Crawfordoides semisinuosa* McCulloch,1981)
- Gavelinellidae* Hofker,1956
- Gyroidinoididae* Saidova,1981
- Gyroidinoides* Brotzen,1942 (TM *Rotalina nitida* Reuss,1844)
- G. sp.* : Murray,1965
 Engl. Channel
- Gavelinellinae* Hofker,1956
- Dicranomalina* Asano,1951 (TM *Dicranomalina japonica* Asano,1951 = *Rotalina semipunctata* Bailey,1851)
 = *Paromalina* Loeblich & Tappan,1957 (TM *Paromalina bilateralis* Loeblich & Tappan,1957 = *Rotalina semipunctata* Bailey,1851)
 = *Pseudorosalinoides* McCulloch,1977 (TM *Pseudorosalinoides chathamensis* McCulloch,1977 = *Rotalina semipunctata* Bailey,1851)
- D. crassa* (Cushman,1931)
 = *Anomalina coronata* Parker & Jones,1857, var. *crassa* Cushman,1931
 = *Paromalina crassa* : Auctt.
 Shetlands, western approaches to the Engl. Channel
- D. coronata* (Parker & Jones,1857)
 = *Anomalina coronata* Parker & Jones,1857
 = *Paromalina coronata* (Parker & Jones,1857) : Gabel,1971
 W Norway, Vadsø, Barents Sea, Shetlands, Iceland, off Canaries
- Gyroidina* d'Orbigny,1826 (TM *Gyroidina orbicularis* d'Orbigny,1826)
- G. orbicularis* d'Orbigny,1826
 off SW Ireland, Mediterranean
- Hansenisca* Loeblich & Tappan,1988 (TM *Gyroidina soldanii* d'Orbigny,1826)
- H. neosoldanii* (Brotzen,1936)
 = *Gyroidina neosoldanii* Brotzen,1936
 = *Gyroidina soldanii* : Auctt., non d'Orbigny,1826
 * Correct genus?
 Bergen, Lofoten, Iceland, S North Sea, western approaches to the Engl. Channel, Galicia
- H. neosoldanii* (Brotzen,1936) var. *nitidula* (Schwager,1866)
 = *Rotalia nitidula* Schwager,1866
 * Correct genus??
 off W Ireland at 1000 fathoms
- Hanzawaia* Asano,1944 (TM *Hanzawaia nipponica* Asano,1944)
 = *Florilus* de Montfort,1808 (suppression suggested - ICZN ZN(S) 2225/2)
- H. asterizans* (Fichtel & Moll,1798)
 = *Nautilus asterizans* Fichtel & Moll,1798
 = *Florilus stellaris* de Montfort,1808 (TM of *Florilus* de Montfort,1808)
 Plymouth, Novaja Zemlya, Mediterranean
- H. boueana* (d'Orbigny,1846)
 = *Nonionina boueana* d'Orbigny,1846
 * Correct species??
 Plymouth
- Karreriidae* Saidova,1981
- Karreria* Rzehak,1891 (TM *Karreria fallax* Rzehak,1891)
 = *Vagocibicides* Finlay,1939 (TM *Vagocibicides maoria* Finlay,1939)
- Trichohyalidae* Saidova,1981
- Buccella* Andersen,1952 (TM *Eponides hannai* Phleger & F.L. Parker,1951)
 = *Mesorotalia* McCulloch,1977 (TM *Mesorotalia fastidiosa* McCulloch,1977)
- B. frigida* (Cushman,1922)
 = *Pulvinulina frigida* Cushman,1922
 = *Eponides frigida* : Auctt.
 = *Pulvinulina karstenii* : Brady,1864, non Reuss,1855
 Øresund, N North Sea, German North Sea, Spitsbergen, Barents Sea, Belgium, Engl. Channel, Wales, NE, NW & W Greenland, Iceland, Biscay
- B. tenerima* (Bandy,19??)
 = *B. inusitata* Andersen,1952
 = *Eponides frigidus* : Cushman & Todd,1947, non *Pulvinulina frigida* Cushman,1922
 Spitsbergen, NE, NW & W Greenland

- Trichohyalus** Loeblich & Tappan,1953 (TM *Discorbis bartletti* Cushman,1933)
- T. pustulata** Loeblich & Tappan,1953
Spitsbergen
- Rotaliacea** Ehrenberg,1839
- Rotalidae** Ehrenberg,1839
- Pararotaliinae** Reiss,1963
- Pararotalia** Y. le Calvez,1949 (TM *Rotalina inermis* Terquem,1882)
= *Neorotalia* Bermúdez,1952 (TM *Rotalia mexicana* Nuttall,1928)
- Ammoniinae** Saidova,1981
- Ammonia** Brönnich,1772
= *Hammonia* Soldani,1789
= *Streblus* Fisher de Waldheim,1817
= *Turbinula* Risso,1826 (TM *Turbinulina italica* = *Rotalia* (les Turbinulines) *italica* d'Orbigny,1826 = *Nautilus beccarii* Linnaeus,1758)
= *Rolshaesenia* Bermúdez,1952 (TM *Rotalia rolshaeseni* Cushman & Bermúdez,1946)
- A. beccarii** (Linnaeus,1758)
= *Nautilus beccarii* Linnaeus,1758 (TM)
Baltic entrance, Øresund, Uddevalla, S Norway, SE North Sea, Faeroes, Iceland, Wadden Sea, S Britain, NW Scotland, Arcachon, Biscay, Galicia
- A. perlucida** (Heron-Allen & Earland,1913)
= *Rotalia perlucida* Heron-Allen & Earland,1913
Plymouth, W Scotland, W Ireland
- A. aberdoveynensis** Haynes,1973
= A. beccarii var. *bavatus* : Adams & Haynes,1965, non Hofker,19??
Wales
- A. batavus** (Hofker,1951)
= *Streblus batavus* Hofker,1951
Oslofjord, North Sea, Engl. Channel, Wales
- A. limnetes** (Todd & Brönnimann,1957)
= *Streblus limnetes* Todd & Brönnimann,1957
Wales, Trindad
- A. tepida** (Cushman,1926)
= *Rotalia beccarii* (Linnaeus) var. *tepidia* Cushman,1926
Poland, Wales, Puerto Rico
- Elphidiidae** Galloway,1933
- Elphidiinae** Galloway,1933
- Criboelphidium** Cushman & Brönnimann,1948 (TM *Criboelphidium vadescens* Cushman & Brönnimann,1948)
= *Elphidionon Hofker,1951* (TM *Polystomella poeyana* d'Orbigny, in de la Sagra,1839)
= *Porosomina Putrya*, in Voloshinova,1958 (TM *Nonionina subgranosa* Egger,1857)
= *Canalifera (Criptoanalifera)* Krasheninnikov,1960 (TM *Canalifera (Criptoanalifera) clara* Krasheninnikov,1960)
= *Retroelphidium Voloshinova*, in Voloshinova & al.,1970 (TM *Elphidium longipontis* Shchedrina,1962)
= *Toddinella Gudina & Levchuk,1983* (TM *Elphidium? ustulatum* Todd,1957)
- C. longipontis** (Shchedrina,1962)
= *Elphidium longipontis* Shchedrina,1962
Barents Sea
- Elphidiella** Cushman,1936 (TM *Polystomella arctica* Parker & Jones, in Brady,1864)
= *Saidovella Voloshinova & Kuznetsova, in Voloshinova & al.,1970* (TM *Saidovella okhotica* Voloshinova & Kuznetsova, in Voloshinova & al.,1970)
= *Pseudelphidiella Voloshinova & Kuznetsova, in Voloshinova & al.,1970* (TM *Elphidium gorbunovi* Shchedrina,1946)
=? *Cryptolophidella* Feyling-Hansen, in Feyling-Hansen & al.,1971 (TM *Cryptolophidella itriensis* Feyling-Hansen, in Feyling-Hansen & al.,1971)
- E. arctica** (Parker & Jones, in Brady,1864)
= *Polystomella arctica* Parker & Jones, in Brady,1864
North Sea, E & W Scotland, W Ireland, Faeroe Channel, Shetlands, Spitsbergen, Novaja Zemlya, Franz Josef Land, Iceland, NE & NW Greenland, Arctic Ocean
- E. groenlandica** (Cushman,1933)
= *Elphidium groenlandicum* Cushman,1933
Iceland, NE Greenland
- E. clavatum** de Montfort,1808 (TM *Nautilus macellus* var. β Fichtel & Moll,1798)
= *Geophorus* de Montfort,1808 (TM *Nautilus macellus* var. α Fichtel & Moll,1798)
= *Cellanthus* de Montfort,1808 (TM *Nautilus craticulus* Fichtel & Moll,1798)
= *Vorticilla* de Lamarck,1812 (TM *Nautilus craticulus* Fichtel & Moll,1798)
= *Polystomella* de Lamarck,1822 (TM *Nautilus crispus* Linnaeus,1758)
= *Polystomatum* Ehrenberg,1839 (TM *Nautilus strigillatus* Fichtel & Moll,1798)
= *Cellula* Agassiz,1844
= *Helicoza Möbius,1880* (TM *Nautilus craticulus* Fichtel & Moll,1798)
= *Carpenterella* Krasheninnikov,1953, non Collenette,1953, nec Bermúdez,1949 (TM *Nautilus craticulus* Fichtel & Moll,1798)
= *Canalifera* Krasheninnikov,1953 (name not available)
= *Faujasinella* Voloshinova,1958 (TM *Elphidium semivolutum* Myatlyuk, in Dabagyan, Myatlyuk & Pischanova,1956)
= *Planoelphidium* Voloshinova,1958 (TM *Polystomella lamina* Terquem,1878)
= *Canalifera* Krasheninnikov,1960 (TM *Elphidium eichwaldi* Bogdanovich, in Serova,1955)
= *Perfectonion Voloshinova & Leonenko, in Voloshinova & al.,1970* (TM *Elphidium incertum* (Williamson) var. *obscurnum* Voloshinova, in Voloshinova & Petrov,1939)
- E. articulatum** (d'Orbigny,1839)
= *Polystomella articulata* d'Orbigny,1839
= *Criononion cf. alvarezianum* : Lutze,1965
= E. *excavatum* : Auct., non (Terquem,1875)
Kieler Bucht, Wadden Sea, Engl. Channel, N Wales, Arcachon
- E. crispum** (Linnaeus,1758)
= *Nautilus crispus* Linnaeus,1758
= *Themone rigatus* de Montfort,1808 (TM of *Themone* de Montfort,1808)
= *Polystomella crispa* (Linnaeus,1758)
Bergen, Iceland, Belgium, Shetlands, Orkneys, Engl. Channel, N Kent, Arcachon, Galicia - southwards
- E. macellum** (Fichtel & Moll,1798)
= *Nautilus macellus* Fichtel & Moll,1798
= *Nautilus strigillatus* var. α Fichtel & Moll,1798 (TM of *Andromedes* de Montfort,1808)
= *Nautilus strigillatus* var. β Fichtel & Moll,1798 (TM of *Sporilus* de Montfort,1808)
= *Elphidium crispum* (Linnaeus) subsp. *spinosum* Atkinson,1969
* May be a synonym of E. *crispum* (Linnaeus,1758)
Engl. Channel, Wales, Oslofjord (post glacial), Barents Sea
- E. margaritaceum** Cushman,1930
= *Elphidium advenum* (Cushman) var. *margaritaceum* Cushman,1930
- Osfjord, German North Sea, Netherlands, Wales
- E. advenum** (Cushman,1922)
= *Polystomella advena* Cushman,1922
Engl. Channel
- E. gerthi** Van Vorthuysen,1957
Baltic entrance, Bristol Channel, S Cornwall
- E. magellanicum** Heron-Allen & Earland,1932
German North Sea, Bristol Channel, Wales, Engl. Channel
- E. cuvillieri** Lévy,1966
Engl. Channel, Wales
- E. exoticum** Haynes,1973
Wales
- E. asterotuberculatum** (Voorhuisen,1958)
= *Nonion depressulus* (Walker & Jacob) var. *asterotuberculatum* Voorhuisen,1958
= *Criononion askundi* : Lutze,1965, non (Brotzen, in Hessland,1946)
SW Baltic, Oslofjord, German North Sea, Netherlands, Wales
- E. earlandi** Cushman,1936
Bristol Channel, S. Cornwall, SW Spain
- E. incertum** (Williamson,1858)
= *Polystomella umbilicatula* (Walker) var. *incerta* Williamson,1858
= E. *incertum* : Cushman,1948
= E. *voorthuyseni* Haake,1962
= E. *varium* Buzas,1965
= *Criononion incertum* : Auct.
SW Baltic, Uddevalla, Oslofjord, along entire Norway, Barents Sea, Iceland, W Scotland, Wales, Engl. Channel
- E. incertum** (Williamson,1858) subsp. A : Lutze,1965
S & W Baltic
- E. excavatum** (Terquem,1875)
= *Polystomella excavata* Terquem,1875
?Engl. Channel
- E. clavatum** Cushman,1930 *selseense* (Heron-Allen & Earland,1911)
= *Polystomella striatopunctata* (Fichtel & Moll) var. *selseensis* Heron-Allen & Earland,1911
= *Elphidium excavatum* (Terquem,1875) forma *selseense* : Feyling-Hanssen,1972 (p.p. - non plate 5 fig.1-4)
= *Elphidium selseense* : Haake,1962, : Haynes,1973
German North Sea, Belgium, E & W Scotland, Engl. Channel, Wales, W Ireland
- E. clavatum** Cushman,1930 *lidoense* Cushman,1936
= *Elphidium lidoense* Cushman,1936, : Haake,1977
= *Elphidium excavatum* (Terquem,1875) forma *lidoensis* : Feyling-Hanssen,1972
= *Elphidium clavatum* : Buzas,1965 (p.p. - plate 2, fig. 6-7) : Buzas,1966 (p.p. - plate 71, fig. 3-4) : Hansen,1965, : Murray,1969
= *Criononion excavatum* : Haake,1967 (p.p. - plate 1, fig. 3,6 & 8), non (Terquem,1876)
Øresund, S North Sea, W France, Mediterranean
- E. clavatum** Cushman,1930 *album* Feyling-Hansen,1972
= *Elphidium excavatum* (Terquem,1875) forma *alba* Feyling-Hansen,1972
= *Elphidium incertum* : Cushman,1930, : 1944, non (Williamson,1858)
S Baltic, all Norway, Iceland
- E. clavatum** Cushman,1930 *clavatum* Cushman,1930
= *Elphidium incertum* (Williamson) var. *clavatum* Cushman,1930
= *Criononion excavatum* : Lutze,1965, : Haake,1967 (p.p. - plate 1, fig. 7,10 & 11), non (Terquem,1876)
= *Elphidium excavatum* : Knudsen,1973, non (Terquem,1876)
S Baltic (relict population), N Norway, Barents Sea, N, NW & W Greenland, Spitsbergen, Jan Mayen
- E. clavatum** Cushman,1930 *nudum* Wilkinson,1979
= *Elphidium clavatum* : Loeblich & Tappan,1953 (p.p. - plate 19, fig. 9-10)
= *Elphidium clavatum* : Feyling-Hansen & al.,1971 (p.p. - plate 20, fig. 5-6)
Spitsbergen, Jan Mayen
- E. clavatum** Cushman,1930 *terminatum* Wilkinson,1979
= *Elphidium subclavatum* Gudina,1964 (p.p. - plate 1, fig. 6-8)
= *Elphidium excavatum* (Terquem,1875) forma *clavata* : Feyling-Hansen,1972 (p.p. - plate 1, fig. 1-5 & plate 2, fig. 4-8)
= *Elphidium excavatum* : Hanssen & Lykke-Andersen,1976
E Greenland, Davis Strait
- E. clavatum** Cushman,1930 *subclavatum* Gudina,1964
= *Elphidium subclavatum* Gudina,1964 (p.p. - plate 1, fig. 4-5)
= *Elphidium clavatum* : Loeblich & Tappan,1953 (p.p. - plate 9, fig. 8)
= *Elphidium clavatum* : Feyling-Hansen & al.,1971 (p.p. - plate 11, fig. 10-11)
Spitsbergen
- E. subarcticum** Cushman,1944
= *Nonion paucilocum* Cushman,1944
* *Nonion paucilocum* should have priority over E. *subarcticum*
Osfjord, Barents Sea, Spitsbergen, NE & W Greenland
- E. waddensis** Van Vorthuysen,1951
= *Elphidium gunteri* Cole var. *waddensis* Van Vorthuysen,1951
= *Elphidium oceanensis* (d'Orbigny,1826)
=? *Polystomella oceanensis* d'Orbigny,1826
German North Sea, Wadden Sea, Belgium, Engl. Channel, Arcachon
- E. williamsoni** Haynes,1973
= *Polystomella umbilicata* Williamson,1858, non *Nautilus umbilicatus* Walker & Jacob,1798
= *Elphidium excavatum* : Auct., non (Terquem,1875)
= *Elphidium articulatum* : Auct., non d'Orbigny,1839
= *Criononion cf. alvarezianum* : Lutze,1965, non (d'Orbigny,1839)
Poland, SW Baltic, Oslofjord, German North Sea, Belgium, Engl. Channel, Wales, Ireland
- E. bartletti** Cushman,1933
= *Nonion striatopunctata* Parker & Jones,1865, non *Nautilus striato-punctatus* Fichtel & Moll,1798
= *Criboelphidium arcicium* Tappan,1951
Spitsbergen, NE, N & NW Greenland, ?Novaja Zemlya
- E. frigidum** Cushman,1933
Iceland, Spitsbergen, NE, N & NW Greenland
- E. orbiculare** (Brady,1881)
= *Nonionia orbicularis* Brady,1881
= *Nonion orbiculare* : Cushman,1930, : 1939, : 1948
Spitsbergen, Novaja Zemlya, NE, N & NW Greenland, Arctic Ocean, off W Scotland, Faeroe Channel
- E. striatopunctata** (Fichtel & Moll,1798)
= *Nautilus striato-punctatus* Fichtel & Moll,1798

- * Correct genus?
Plymouth
- E. subnodosum** (Münster, *fide* Roemer, 1838)
= *Robulina subnodosa* Münster, *fide* Roemer, 1838
- * Correct genus?
Plymouth
- E. lessonii** (d'Orbigny, 18??) : Gabel, 1971
North Sea
- E. aculeatum** (d'Orbigny, 18??) : Gabel, 1971
North Sea
- E. goesi** Shchedrina, 19??
Barents Sea
- E. minutum** Reuss, 18??
= *Cibronion minutum* (Reuss, 18??) : Gabel, 1971
- * Correct genus?
North Sea
- E. sibiricum** (Goës, 1894)
= *Polystomella sibirica* Goës, 1894
- * Correct genus?
off Novaja Zemja
- E. sp.** : Murray, 1968
Engl. Channel
- Nummulitacea de Blainville, 1827
- Nummulitidae de Blainville, 1827
- Heterostegina** d'Orbigny, 1826 (TM *Heterostegina depressa* d'Orbigny, 1826)
= *Grzybowskia* Bieda, 1950 (TM *Grzybowskia multifida* Bieda, 1950)
- Foraminifer genera of uncertain status:
- Crateriola** Strand, 1943 (TM *Craterella albescens* Dons, 1942)
= *Craterella* Dons, 1942, non Schrammen, 1901 (Spongaria), nec Kofoid & Campbell, 1929 (Ciliophora)
- Earlmyseria** Rhumbler, 1938
- E. punctulata** (d'Orbigny) forma *tiliputana* Rhumbler, 1938 (TM)
Helgoland, Scotland, W Ireland
- ACTINOPODEA Calkins, 1902
- ACANTHAREA Haeckel, 1881
- HOLACANTHIDA Schewiakoff, 1???
- Acanthochiasmida Haeckel
- Acanthochasma** Krohn, in Haeckel, 1861
- A. fusiforme** Haeckel, 1861?
Skagerrak, North Sea, Mediterranean
- A. rubescens** (Krohn, 18??)
Mediterranean
- A. krohni** Haeckel, 1862
Skagerrak, Bergen
- Acanthocyrta** Schewiakoff, 1926
- A. haeckeli** Schewiakoff, 1926?
?= *A. decacanthum* Popofsky, 1904
?= *A. bicuspidatum* Schewiakoff, 1926
Mediterranean
- Acanthoplegmidae Schewiakoff
- Acanthoplegma** Schewiakoff, 1926?
* For this genus Rechetniak, 1981 created the new order Plegmacantha, a view confirmed by electron microscopical data on one of its species, but related genera should be investigated before acceptance of this new taxon
- Acanthospira** Reinsch, 1877
- Acanthocolla** Schewiakoff, 1926
- SYMPHYACANTHIDA Schewiakoff
- Astrolithiidae Haeckel, 18?? emend. Schewiakoff
- Astrolithium** Haeckel, 1861
- A. bifidum** Haeckel, 1862
= *Acanthometra astrodes* Hertwig, 1879
= *Acanthometron bifidum* : Auctt.
N Atlantic S of 60° N.
- Acantholithium** Haeckel, 1887
- Astrolonche** Haeckel, 1881
- Heliolithium** Schewiakoff, 1926
- Amphilithiidae Haeckel, 18?? emend. Schewiakoff
- Amphilithium** Haeckel, 1881
- Amphibelone** Haeckel, 1862
- Pseudolithiidae Schewiakoff
- Pseudolithium** Schewiakoff, 1926
- Dicranophora** Schewiakoff, 1926, non Macquart, 1834 (Diptera)
- Haliommatidae
- Haliommatidium** J. Müller, 1858
- CHAUNACANTHIDA Schewiakoff
- Conaconidae Schewiakoff
- Conacon** Schewiakoff, 1926
- Gigartaconidae Schewiakoff
- Gigartacon** Schewiakoff, 1926
= *Sphaerocapsa* Haeckel, 1881
- G. cruciata** (Haeckel, 1887)
= *Sphaerocapsa cruciata* Haeckel, 1887
Faeroe Channel surface water
- Heteracon** Schewiakoff, 1926
- Amphiaconidae Schewiakoff
- Stauraconidae** Schewiakoff
- Stauracon** Schewiakoff, 1926
- ARTHRACTHIDA Schewiakoff
- Radiosphaera** Jørgensen, in Nordgaard & Jørgensen, 1905
* Correct order?
- R. anacanthica** Jørgensen, in Nordgaard & Jørgensen, 1905
Lofoten, Tromsø, Balstad, Skjerstadfjord, Foldenfjord
- Zygacantha** J. Müller, 1858
* Correct order?
- Z. septentrionalis** Popofsky, 1904
60.2° N 22.7° W at 800-1000 m.
- Z. rotunda** Popofsky, 1904
59.2-60.2° N 11.8-22.7° W
- Acanthonia** Haeckel, 1881
* Correct order?
- A. tetracopa** (J. Müller, 1858) Haeckel, 1881?
= *Acanthomera tetracopa* J. Müller, 1858
Bergen, 76-80° N 10-12° E, Mediterranean
- A. muelleri** Haeckel, 1862
= *Litholophus decapristis* Haeckel, 1887
* Is this possibly *Acanthometron muelleri* Hckl : Ostenfeld, 1916 from Skagerrak and is this a synonym of *Lithopora muelleri* Haeckel (see below)?
North Sea, Orkneys, Bear Island, Mediterranean
- A. ligurina** (Haeckel, 1865)
= *Litholophus ligurinus* Haeckel, 1865
= *Litholophus arcticus* Aurivillius, 1899
Stadt, Iceland, Spitsbergen, W Greenland
- A. nordgaardi** (Jørgensen, 1899)
= *Acanthostaurus nordgaardi* Jørgensen, 1899
W Norway
- A. pusilla** Jørgensen, 1899
Bergen
- Acanthonidium** Haeckel, 1887
* Correct order?
- A. echinoides** (Claparède & Lachmann, 1858)
Acanthometra echinoides Claparède & Lachmann, 1858
= *Acanthonia echinoides* : Jørgensen, 1899
= *Zygacanthidium echinoide* : Ostenfeld, 1916
Kattegatt, Skagerrak, North Sea, Bergen, Bodø, Tromsø, Finnmark, Faeroe Channel
- A. pallidum** (Claparède & Lachmann, 1858)
= *Acanthometra pallida* Claparède & Lachmann, 1858
= *Zygacanthidium pallidum* : Ostenfeld, 1916
= *Acanthostaurus pallidus* (Claparède & Lachmann, 1858) Haeckel, 1862
Skagerrak, North Sea, Bergen, Bodø, E Scotland, Faeroe Channel
- A. pallidum** var. *subulatum* (Jørgensen, 1899)
= *Acanthostaurus pallidus* (Claparède & Lachmann, 1858-59) Haeckel, 1862 var. *subulata* Jørgensen, 1899
Bergen
- A. cuspidatum** (Haeckel, 1862)
= *Acanthometra cuspidata* Haeckel, 1862
50-57° N 10-34° W
- A. tetrapterum** (Haeckel, 1860)
= *Acanthometra tetraptera* Haeckel, 1860
= *Acanthometra messanensis* Haeckel, 1860
24° S - 40 (56)° N
- Astrocapsa** Haeckel, 1887
* Correct order?
- A. tritonis** Haeckel, 1887
Faeroe Channel surface water
- A. coronata** Haeckel, 1887
Faeroe Channel surface water down to 200 fathoms
- Porocapsa** Haeckel, 1887
* Correct order?
- P. murrayana** Haeckel, 1887
Faeroe Channel surface water down to 640 fathoms
- P. octodon** Haeckel, 1887
Greenland
- Cannocapsa** Haeckel, 1887
* Correct order?
- C. osculata** Haeckel, 1887
Faeroe Channel surface water
- SPHAENACANTHINA Schewiakoff
- Acanthometridae Haeckel, 18?? emend. Schewiakoff
- Acanthometra** J. Müller, 1855
= *Acanthometron* Haeckel, 1887
- A. pellucida** J. Müller, 1855
= *A. elastica* Haeckel, 1862
Kattegatt, Bohusl, Skagerrak, Bergen, Stadt, Murmansk, Faeroes, Orkneys, Scotland, Mediterranean

- Amphilonche** Haeckel,1860
- A. belonoides** (Haeckel,1860)
= *Acanthometra belonoides* Haeckel,1860
= *Amphilonche concreta* Haeckel,1887
= *Amphilonche aciculata* Haeckel,1887
49°56' N 20°48' W
- A. elongata** (J. Müller,1858)
= *Acanthometra elongata* J. Müller,1858
= *Amphilonche acicula* Haeckel,1887
= *Acantholonche amphipolaris* Haeckel,1887
54°56' N 31°45' W
- A. mira** var. 1 : Popofsky,1904
59°2-60,2° N 11,8-22,7° W
- Tetralonche** Schewiakoff,1926
- Lithopteridae Haeckel,18?? emend. Popofsky
- Lithoptera** J. Müller,1858
- L. muelleri** Haeckel,18??
- Dorataspidae Haeckel,18?? emend. Schewiakoff
- Dorataspis** Haeckel,1861
- D. prototypus** (Haeckel,1887)
= *Phractaspis prototypus* Haeckel,1887
= *Lithophyllum condylatum* Haeckel,1887
Mediterranean
- D. bipennis** Haeckel,1862
Mediterranean
- D. heteropora** Bernstein,1???
- Barents Sea
- Lychnaspis** Haeckel,1881
- L. giltschii** Haeckel,1887
= *Dodecaspis tricinata* Haeckel,1887
41° N 58° W
- Icosaspis** Haeckel,1881
- I. tabulata** Haeckel,1887
= *Xiphacantha emarginata* Haeckel,1887
= *Stauracantha orthostaura* Haeckel,1887
= *Phatnacantha icosaspis* Haeckel,1887
= *Tignisphaera tabulata* : Popofsky,1906
56° N 10° W
- Pleuraspis** Haeckel,1881
- Stauraspis** Haeckel,1881
- Hystrichaspis** Haeckel,1887
- Dictyaspis** Haeckel,1887
- Siphonaspis** Haeckel
- Coscinaspis** Haeckel
- Cranaspis** Haeckel
- Acothaspis** Haeckel
- Hexalaspidae Haeckel
- Coleaspis** Haeckel,1881
- C. obscura** Haeckel,1887
Greenland, Arctic Ocean
- Hexalaspis** Haeckel,1885
- H. heliodiscus** Haeckel,1887
= *H. helioestrum* Haeckel,1887
= *H. hexapleura* Haeckel,1887
39° N 21° W
- Hexaconus** Haeckel,1885
- Diploconidae Haeckel
- Diploconus** Haeckel,1860
- Phractopeltidae Haeckel
- Phractopelma** Haeckel,1881
= *Phractopelta* Haeckel,1887
- Nivalidae
- PHYLLACANTHINA** Schewiakoff
- Phyllostauridae Schewiakoff
- Phyllostaurus** Haeckel,1862
- P. quadrifolius** (Haeckel,1862)
= *Acanthometra quadrifolia* Haeckel,1862
Skagerrak, North Sea, Spitsbergen, Mediterranean
- P. heterobelus** (Jørgensen,1899)
= *Acanthonia (?) heterobelus* Jørgensen,1899
W Norway
- Amphistaurus** Schewiakoff,1926
- Acanthostaurus** Haeckel,1862
- Lonchostaurus** Haeckel,1862
- Zygostaurus** Haeckel,1887
- Stauracanthidae Haeckel
- Xiphacantha** Haeckel,1861
- X. quadridentata** (J. Müller,1858) Haeckel,1862
= *Acanthometra quadridentata* J. Müller,1858
Bergen (Puddefjord)
- Pristacantha** Haeckel,1887
- Stauracantha** Haeckel,1881
- Phatnacantha** Haeckel,1881
- Dictyacanthidae Schewiakoff
- Dictyacantha** Schewiakoff,1926
- POLYCYSTINEA** Ehrenberg,1838 emend. Riedel,1967
- SPUMELLARIDA Ehrenberg,1858
- Rhopalodictyum** Ehrenberg,1860
- R. subacutum** Ehrenberg,1861
Faeroe Channel, Greenland
- Spongodiscidae
- Spongodiscus** Ehrenberg,1854
- S. favus** Ehrenberg,1861
Gaukvarð, Røst, Faeroe Channel, Greenland
- S. resurgens** Ehrenberg,1854
54° N 22° W
- Spongotrochus** Haeckel,1860
- S. glacialis** (Popofsky,1908)
Bergen, off E Greenland
- Theoperidae Haeckel,1881, emend. Riedel,1967
- Dictyophimus** Ehrenberg,1847, emend. Nigrini,1968
- D. multispinus** Bernstein,1???
- Barents Sea
- SPHAEROCOLLINA
- Orosphaeridae Haeckel,1887
- Oroscena** Haeckel,1887
- O. regalis** Borgert,1901
60,3°N 27°W
- Thalassothamnidiae Haeckel
- Bathysphaeridae Hollande & Enjumet
- Sphaeroidae Haeckel = Sphaerozoidae
* Temporary family.
- Collozoum** Haeckel,1862
- C. ellipsoidea** Haeckel,1887
Faeroes, between Faeroes & Shetlands, Faeroe Channel
- C. brandti** Wolfenden,1902
Orkneys
- C. groenlandicum** Pouchet,1893
between Greenland, Iceland & Norway, Barents Sea
- C. inerme** ?,1???
- * 2 species of parasitical dinoflagellates are known from this species: *Merodinium brandti* Hovasse,1923 (intracapsulaire) & *Merodinium verna* Hovasse,1923 (extracapsulaire).
boreal Atlantic, arctic Atlantic
- C. sp.** : Wolfenden,1902
Shetlands, Faeroe Channel
- Sphaerozoum** Meyen,1834
- S. hexactinum** Haeckel,1887
Faeroe Channel
- S. punctatum** ?,1???
- * sometimes with cytoplasma parasite *Merodinium dolosum* Chatton,1923
boreal Atlantic
- Raphidozoum** Haeckel,1862
- Belonozoum** Haeckel,1887
- Collosphaeridae J. Müller,1858
* Temporary family.
- Collosphaerinae J. Müller,1858
- Collosphaera** J. Müller,1858
- C. huxleyi** J. Müller,18??
boreal Atlantic, SW Ireland
- Acrosphaerinae
- Acrosphaera** Haeckel,1881
- A. murrayana** (Haeckel,1887)
= *Choenocapsphaera murrayana* Haeckel,1887
W Norwegian fjords, North Sea, Shetlands, Faeroes, Orkneys, off W Ireland
- A. spinosa** (Haeckel,1862)
= *Collosphaera spinosa* Haeckel,1862
= *Choenocapsphaera nassiforma* Haeckel,1887
= *Choenocapsphaera flammabunda* Haeckel,1887
boreal Atlantic, ?Shetlands, Mediterranean
- Tribonosphaera** Haeckel,1881
- Siphonosphaerinae
- Siphonosphaera** J. Müller,1858

- Solenosphaera* Haeckel,1887
Buccinosphaera Haeckel,1887
Exoxoplastidiata Cachon & Cachon
Thalasicollidae Haeckel
Thalassicolla Huxley,1851
T. nucleata Huxley,18??
 Shetlands, Faeroe Channel, SW Ireland, Sargasso Sea
T. borealis Ludwig,1908
 Irminger Sea
T. irmingiana Ludwig,1908
 Irminger Sea
T. gelatinosa Brandt,1905
 NE North Sea, Mediterranean
T. porosa Ludwig,1908
 Egersund
Thalassisolen Wolfenden,1902
T. atlanticus Wolfenden,1902
 Shetlands, Faeroe Channel
Actissa Haeckel,1884
A. septentrionalis Ludwig,1908
 off S Iceland
Thalassoplancta Haeckel,1862
T. longispicula Haeckel,1887
 Faeroe Channel
Thalassoxanthium Haeckel,1881
T. ovodimare Haeckel,1887
 off W Hebrides
Lampoxanthium Haeckel,1887
L. murayananum Fowler,1898
 Faeroe Channel
Thalassophysidae Brandt
Thalassophysa Haeckel,1881
T. pelagica (Haeckel,1862)
 = *Thalassicolla pelagica* Haeckel,1862
 Shetlands, Faeroe Channel, Mediterranean
Polycyrtaria
Excryptoaxoplastidiata Cachon & Cachon
Physematidae Brandt
Centroaxoplastidiata Cachon & Cachon
Centrociliidae Cachon & Cachon
SPHAERELLARINA
Sphaerostylus Haeckel,1881, non Chaudoir,1848 (Coleoptera)
 * Should be transposed to another genus. Which family?
S. cottus Haeckel,1887
 Faeroe Channel
Prunulum Haeckel,1887
 * Which family?
P. triplex (Ehrenberg,1854)
 = *Haliomma triplex* Ehrenberg,1854
 deep N Atlantic
Prunocarpus Haeckel,1887
 * Which family?
P. datura Haeckel,1887
 Faeroe Channel
Trochodiscus Haeckel,1887
 * Which family?
T. echiniscus Haeckel,1887
 Faeroe Channel
T. helioides Cleve,1899
 off W Spitsbergen
Cryptoaxoplastidiata Hollande & Enjumet
Macrosphaeridae
Haplosphaera Hollande & Enjumet,1960
Acanthosphaera Ehrenberg,1858
A. arctica Haeckel,1887
 Greenland
A. haliphormis Ehrenberg,1861
 Deep Greenlandic waters
A. teneriformis Jørgensen,1900
 deep W Norwegian waters
Cladococcus J. Müller,1856
C. viminalis Haeckel,1856
 Bergen, W Norwegian fjords, Mediterranean
C. spinifer Haeckel,1862
 deep W Norwegian waters, Mediterranean
Elaphococcus Haeckel,1881
E. dichotomus Haeckel,1887
 Greenland
E. drymodes Haeckel,1887
 Iceland
Diplosphaera Haeckel,1860
D. reticulata Haeckel,1887
 Faeroe Channel
Haeckeliella Hollande & Enjumet,1960
Thalassoplegma Hollande & Enjumet,1960
Rhizoplegma Haeckel,1881
R. boreale (Cleve,1899) Jørgensen,1???
 = *Hexadonous borealis* Cleve,1899
 deep Skagerrak, Bergen, Tysfjord, Skroven, Helligvær, Vestfjord, Stamsund, Henningsvær, Gaukvær, Tromsø, Malangen, Tranøybælt, Ofoten, Rombaken, Øxsund, Sagfjord, Lofoten, Røst, Svolver, Skjerstadfjord, Foldenfjord, Saltenfjord
Lychnosphaera Haeckel,1881
Heliosphaera Haeckel,1861
 * Correct family?
H. minuta Cleve,1900
 between Faeroes & Rockall
H. tenera Jørgensen,1900
 Bergen
H. actinota Haeckel,1862
 deep Arctic Ocean, Mediterranean, Azores, Canaries
Drymyomma Jørgensen,1900
 * Correct family?
D. elegans Jørgensen,1900
 = *Drymyosphaera elegans* : Björklund,1974
 Bergen, Tysfjord, Vestfjord, Gaukværø, Tromsø, Skroven, W Norwegian fjords
Elatomma Haeckel,1887
 * Correct family?
E. penicillatus Haeckel,1887
 Bergen
E. irregulare Haeckel,1887
 Faeroe Channel
Leptosphaera Haeckel,1887
 * Correct family?
L. arachnoides Jørgensen,1900
 Bergen
Arachnosphaera Haeckel,1860
 * Correct family
A. dichotoma Jørgensen,1900
 = *Haliomma spinulosum* var. Cleve,1900
 W Norway
Centrocubidae Hollande & Enjumet,19??
Centrocubus Haeckel,1887
C. mediterranensis Hollande & Enjumet,1960
 Mediterranean
Octodendron Haeckel,1887
Centroaxoplastidiata Hollande & Enjumet
 ?Family?
Ethmosphaera Haeckel,1861
E. leptosiphonia Haeckel,1887
 Faeroe Channel
Carposphaera Haeckel,1881
C. belladonna Haeckel,1887
 Faeroe Channel
Spongphaeridae Haeckel,18
Spongphaerinae Haeckel,18
Spongphaera Ehrenberg,1847
Spongodendron Lendenfeld,1887
Hexadoras Haeckel,1881
Plegmosphaerinae Hollande & Enjumet
Plegmosphaera Haeckel,1881
P. coelopila Haeckel,1887
 Faeroe Channel
Spongoplegma Haeckel,1881
Spongodyminae Hollande & Enjumet
Diplosponges Mast,1910
Spongadrymus Haeckel,1881
Rhizosphaeridae Hollande & Enjumet
Rhizosphaera Haeckel,1860
Actinosphaera Hollande & Enjumet,1960

- = *Haliomma* Ehrenberg,1839 (p.p.)
- A.? boreale* (Haeckel,1887)
= *Haliomma boreale* Haeckel,1887
Greenland
- A.? echinosphaera* (Cleve,1900)
= *Haliomma echinosphaera* Cleve,1900
45° N 49° W
- A.? beroes* (Ehrenberg,1854)
Haliomma beroes Ehrenberg,1854
54°N 22° W
- A.? irregularare* (Cleve,1900)
= *Haliomma irregularare* Cleve,1900
45° N 49° W
- Thecosphaeridae Hollande & Enjumet,19??
- Hexacontium Haeckel,1882
- H. entacanthum* Jørgensen,1900
deep Skagerrak, Bergen, Tysfjord, Tromsø, Skroven, Henningsvær, Røst, Tranøybet, Foldenfjord, off N Shetlands, off N Faeroes
- H. pachydermum* Jørgensen,1900
deep Skagerrak, Bergen, Tysfjord, Helligvær, Moskenstrømmen, Stamsund, Tromsø, Skroven, Ofoten, Øxsund, Henningsvær, Svolver, Tranøybet, Foldenfjord, off N Shetlands, off N Faeroes
- H. hostile* Cleve,1900
Bergen, W Norwegian fjords, off S Iceland
- H. macracanthum* Jørgensen,1900
Bergen
- H. setosum* Haeckel,1887
off S Iceland, Mediterranean
- H. armatum* Cleve,1900
off S Iceland
- H. sp. C* : Bjørklund,1974
=? *H. sp.* : Swanberg & Bjørklund,1987
Bergen, ?W Norwegian fjords
- Actinomma* Haeckel,1862
= *Echinomma* Haeckel,1881
= *Pityomma* Haeckel,1881
- A. trinacrium* Haeckel,1862
= *Echinomma trinacrum* : Auctt.
Bergen, Tysfjord, Vestfjord, Skroven, Helligvær, Gaukværø, Tromsø, Ofoten, Sagfjord, Henningsvær, W Norwegian fjords, Mediterranean
- A. leptodermum* (Jørgensen,1900)
= *Echinomma leptodermum* Jørgensen,1900
deep Skagerrak, Bergen, Tysfjord, Skroven, Helligvær, Gaukværø, Tromsø, Malangen, Vestfjord, Rombaken, Ofoten, Tranøybet, Henningsvær, Lofoten, Øxsund, Røst, Skjerstadfjord, Foldenfjord, Saltenfjord, Barents Sea
- A. mediterraneans* Hollande & Enjumet,1960
Mediterranean
- A. sp. A* : Swanberg & Bjørklund,1987
W Norwegian fjords
- A. sp. B* : Swanberg & Bjørklund,1987
W Norwegian fjords
- A. sol* Cleve,1900
45°N 49°W
- Cromyechinus* Haeckel,1881
= *Chromyechinus* Jørgensen, in Nordgaard & Jørgenssen,1905
- C. borealis* (Cleve,1899)
= *Actinomma boreale* Cleve,1899
= *Chromyomma borealis* : Jørgensen,1900
= *Chromyechinus borealis* : Bjørklund,1974
deep Skagerrak, Bergen, Røst, Tysfjord, Vestfjord, Gaukværø, Tromsø, Malangen, Skroven, Ofoten, Tranøybet, Øxsund, Spitsbergen, W Norwegian fjords
- Thecosphaera* Haeckel,1881
- T. sp. A : Bjørklund,1974
Bergen
- Cromyomma* Haeckel,1862
* Correct family?
- C. zonaster* (Ehrenberg,1872)
= *Acanthosphaera zonaster* Ehrenberg,1872
Greenland
- Hexalonche* Haeckel,1881
* Correct family?
- H. philosophica* Haeckel,1887
Faeroe Channel, Iceland
- H. diplocaantha* Jørgensen,1900
Bergen
- Periaxoplastidiata Hollande & Enjumet
- Cenosphaera* Ehrenberg,1854
? Which family?
- C. favosa* Haeckel,1887
Faeroe Channel
- Heliaster* Hollande & Enjumet,1960
* Fam. Heliasteridae (which superfamily?)
- Tetrapetalon* Hollande & Enjumet,1960
* Fam. Tetrapetalidae (which superfamily?)
- Arachnostylus* Hollande & Enjumet,1960
* Fam. Arachnostyliidae (which superfamily?)
- Cyrtidospaeridae Cachon & Cachon
- Cyrtidospaera* Haeckel,1860, emend. Cachon & Cachon,19??
- Stigmosphaeridae Hollande & Enjumet
- Stigmosphaera* Haeckel,1887
- Stigmosphaerusa* Hollande & Enjumet,1960
- Hexastylus* Haeckel,1882
- H. sp. B* : Bjørklund,1974
Bergen
- Stigmatostylus* Hollande & Enjumet,1960
- Excentroconchidae Hollande & Enjumet
- Excentroconcha* Mast,1910
- Hollandosphaeridae Deflandre
- Heliosoma* Haeckel,1881
- Hollandosphaera* Deflandre,19??
- Prunoidea* Haeckel
* with 7 families & 50 genera
- Ellipsidae
- Cenellipsis* Haeckel,1887
- Sponguridae
- Spongoliva* Haeckel,1887
- Panartidae
- Panartus* Haeckel,1887
- Discoidea Haeckel
* with 6 families & 100 genera
- Phacodiscidae
- Astrophacus* Haeckel,1881
- A. saturnus* (Ehrenberg,1861)
= *Chilomma saturnus* Ehrenberg,1861
?Greenland
- Astroestrum* Haeckel,1881
- A. acanthastrum* Haeckel,1887
Faeroe Channel
- Sethodiscus* Haeckel,1881
- S. macroporus* Haeckel,1887
Faeroe Channel
- Porodiscidae
- Euchitonnia* Ehrenberg,1861
- E. sp. 1* : Bjørklund,1974
Bergen
- E. sp. 2* : Bjørklund,1974
Bergen
- Stylocidya* Ehrenberg,1847, emend. Koslova,1972 (TM) *Stylocidya gracilis* Ehrenberg,1854
- S. tenuispina* Jørgensen,1905
Gaukværø, Røst, W Norwegian fjords
- S. aculeata* Jørgensen,1905
Røst
- S. validispina* Jørgensen,1905
Røst, W Norwegian fjords, off E Greenland
- S. aspera* Jørgensen,1900
Bergen
- S. sp.* : Bjørklund,1974
Bergen
- Hagiastrum* Haeckel,1881
- H. cristif* Haeckel,1887
Faeroe Channel
- Rhopalastrum* Ehrenberg,1847
- R. arcticum* Haeckel,1887
Arctic Ocean
- Amphicraspedum* Haeckel,1881
- A. murrayanum* (Haeckel,1879)
= *Amphymentium murrayanum* Haeckel,1879
Faeroe Channel
- Pylodiscidae
- Triolena* Haeckel,1887
- Larcoidea* Haeckel
* with 9 families & 50 genera
- Larcaridae
- Spongolarcus* Haeckel,1887
- S. amphicentria* Haeckel,1887
= *Amphicentria salpa* Ehrenberg,1861
= *Spongurus salpa* : Haeckel,1862
Greenland
- Pylonidae* Haeckel,18??
- Phorticium* Haeckel,1882
* Correct family?

- P. clevei* (Jørgensen,1900)
 = *Tetrapylonum clevei* Jørgensen,1900
 = *Phorticium pylonum* : Cleve,1899, non Haeckel,1887
 deep Skagerrak, Tysfjord, Skroven, Vestfjord, Henningsvær, Tromsø, Ofoten, Øxsund, Lofoten, Røst, Skjerstadfjord, Spitsbergen
- Octopyle* Haeckel,1881
- O. octostyle* Haeckel,1887 forma *minor* Jørgensen,1900
 W Norway
- Monozonium* Haeckel,1887
- Tetrapyle* J. Müller,1858
- T. polyacantha* Jørgensen,1900
 W Norway
- T. quadriloba* (Ehrenberg,1860)
 = *Schizomma quadrilobum* Ehrenberg,1860
 Greenland
- Litheliidae
- Spireuma* Haeckel,1881
 = *Spirema* Haeckel,1887
- S. haliomma* (Ehrenberg,1861)
 = *Flustrella haliomma* Ehrenberg,1861
 = *Spirema flustrella* Haeckel,1887
 Bergen, Faeroe Channel, Greenland
- Lithelius* Haeckel,1860
- L. spiralis* Haeckel,1860
 Bergen, Tromsø, W Norwegian fjords, Mediterranean
- L. arborescens* Haeckel,1887
 W Norwegian fjords, Faeroe Channel
- L.* sp. : Björklund,1974
 Bergen
- Larcospira* Haeckel,1887
- L. minor* (Jørgensen,1900)
 = *Lithelius minor* Jørgensen,1900
 Bergen, Tysfjord, Vestfjord, Gaukværø, Tromsø, Skroven, Tranøybæt, Sagfjord, Henningsvær, Lofoten, Røst, Foldenfjord
- Strebliomidae Haeckel,1887
- Streblopyle* Haeckel,1887
- Streblacantha* Haeckel,1887
- S. circumtexta* Jørgensen,1905
 = *Sorolarcus circumtextus* Jørgensen,1900 (p.p.)
 Bergen, W Norwegian fjords
- Soreumidae
- Sorolarcus* Haeckel,1887
- S. circumtextus* Jørgensen,1900
 * Is this species and *Strebacantha circumtexta* Jørgensen,1905 not identical?
 Tysfjord, Skroven, Vestfjord, Henningsvær, Gaukværø
- NASSELLARIDA Ehrenberg,1875 emend Haeckel, emend Petrushevskaya,1971
- Gonosphaera* Jørgensen,1905
- G. primordialis* Jørgensen,1905
 Bergen, W Norwegian fjords
- Lampronitra* Haeckel,1882
- L. erosa?* Cleve,1901
 Bergen
- Ceratocystis* Bütschli,1881
- C. galeus* (Cleve,1899)
 = *Sethocomas galea* Cleve,1899
 W Norwegian fjords, between Spitsbergen & Greenland
- C. histricosus* (Jørgensen,1905)
 = *Dictyophimus histricosus* Jørgensen,1905
 Bergen, W Norwegian fjords
- Pseudodictyophimus* ?,19??
- P. gracilipes* (Bailey,1856)
 = *Dictyophimus gracilipes* Cleve,1899
 = *Dictyophimus clevei* Jørgensen,1900
 Bergen, W Norwegian fjords, Barents Sea, Spitsbergen, off E Greenland
- Sethopera* Haeckel,1881
- S. oceania* (Ehrenberg,1872)
 = *Lithopera oceania* Ehrenberg,1872
 North Atlantic north of 50°N, Azores
- Dictyocephalus* Ehrenberg,1860
- D. papillosum?* Ehrenberg,1860
 Bergen
- D. reticulum* (Ehrenberg,1861)
 = *Eucyrtidium reticulum* Ehrenberg,1861
 Iceland, Greenland
- D. ocellatus* (Ehrenberg,1861)
 = *Eucyrtidium ocellatum* Ehrenberg,1861
 Faeroe Channel, Greenland
- D.* sp. : Cleve,1899
 Deep Arctic Ocean
- Dictyoceras* Haeckel,1862
- D. acanthicum* Jørgensen,1900
- North Sea, Bergen, W Norwegian fjords
- D. xiphophorum* Jørgensen,1900
 W Norway
- Pterocodon* Ehrenberg,1847
- P. amblycephalus* (Jørgensen,1900)
 = *Pterocorys amblycephalis* Jørgensen,1900
 Bergen
- Clathrocyclas* Haeckel,1881
- C. danæs* Haeckel,1887
 56°N 30°W, Azores
- C. coscinodiscus* Haeckel,1887
 53-56°N 36-50°W
- Artobotrys* ?,19??
- A. borealis* (Cleve,1899)
 = *Theocoris borealis* Cleve,1899
 = *Theocrysis borealis* : Björklund,1974
 Bergen, W Norwegian fjords, Spitsbergen, off E Greenland
- Mitrocalpis* Haeckel,1881
- M. aranaefera* (Popofsky,19??)
 W Norwegian fjords
- Theocalyptra* Haeckel,1881
- T. cornuta* (Bailey,1856)
 = *Halicalyptra cornuta* Bailey,1856
 = *Lophophæna calyptra* : Haeckel,1862
 = *Carpocanium cornutum* : Ehrenberg,1872
 N Atlantic north of 41°N, Greenland, off W Spitsbergen
- Theoconus* Haeckel,1887
- T.* sp. : Cleve,1900
 off Stadt
- Theocyrta* Haeckel,1887
- T. aculeata* Cleve,1900
 56°N 17°W, 48°N 29°W
- PLECTELLARINA Haeckel = APOAXOPLASTIDIATA
- Plectoidea Haeckel
- Plagon(d)idae Haeckel,1882, emend. Cachon & Cachon,19??
- Phormacantha* Jørgensen,1905
- P. hystrix* (Jørgensen,1905)
 = *Peridium hystrix* Jørgensen,1900
 Bergen, W Norwegian fjords
- P.* sp. D : Björklund,1974
 ?= *Phormacantha?* sp. : Swanson & Björklund,1987
 Bergen, ?W Norwegian fjords
- Helotholus* Jørgensen,1905
- H. histricosa* Jørgensen,1905
 Bergen
- Acanthocorys* Haeckel,1882
- A. umbellifera* (Haeckel,1862)
 = *Arachnocorys umbellifera* Haeckel,1862
 Bergen, W Norwegian fjords, N Norway, N Ice Sea, Azores, Mediterranean
- Peridium* Haeckel,1882
- P. longispinum* Jørgensen,1900
 Bergen, W Norwegian fjords, Lofoten
- P. minutum* Cleve,1899
 N Norway
- P.* sp. : Swanson & Björklund,1987
 W Norwegian fjords
- P.* sp. : Meunier,1910
 Arctic Ocean, Kara Sea
- P.* sp. ? *Phormacantha* sp. ? : Meunier,1910
 Arctic Ocean, Kara Sea
- Plagonidium* Haeckel,1881
- Tetraplagia* Haeckel,1881
- Plagiocarpa* Haeckel,1881
- P. procryptellaria* Haeckel,1887
 Iceland
- Pentaplagia* ?,19??
- Triplagiacantha* Schröder,1911
- T. abietina* (Hertwig,1879)
 = *Plagiocantha abietina* Hertwig,1879
 Mediterranean, 50-53°N 44-50°W
- Plagiocantha* J. Müller,1856
- P. arachnoides* (Claparède,1855)
 = *Acanthomeira arachnoides* Claparède,1855 (TM)
 = *Plectophora arachnoides* : Auct.
 * unlike most "Radiolaria" but like Acantharia this species lives close to the water surface
 N Kattegatt, Skagerrak, North Sea, Bergen, W Norwegian fjords, Murmansk, White Sea, Spitsbergen, Iceland, between Faeroes & Shetlands, N Scotland
- P.(?) intricata* (Cleve,1899)
 = *Peridium (?) intricatum* Cleve,1899
 * Possibly a synonym of *Plectacantha oikikos* Jørgensen,1905
 between Spitsbergen & Greenland

- P.(?) laxa* (Cleve,1899)
= *Peridium (?) laxum* Cleve,1899
* Possibly a synonym of *Plectacantha oikiskos* Jørgensen,1905
Trømsø
- Campylacantha* Jørgensen,1905
- C. cladophora* Jørgensen,1905
Bergen, W Norwegian fjords, Osterfjord
- Polyplagia* Haeckel,1881
- P. novenaria* Haeckel,1887
Faeroe Channel
- Plectophora* Haeckel,1882
- P. tricantha* (Popofsky,1908)
Bergen, W Norwegian fjords
- P. novena* Haeckel,1887
Faeroe Channel
- Plectacantha* Jørgensen,1905
- P. oikiskos* Jørgensen,1905
Skagerrak, Bergen, W Norwegian fjords
- P. trichoides* Jørgensen,1905
Bergen, W Norwegian fjords
- Protoscenium* Jørgensen,1905
- P. simplex* (Cleve,1899)
= *Plectanum simplex* Cleve,1899
Bergen, Gaukvaerø, off N Shetlands
- Acanthocystium* Haeckel,18??
?= *Acanthocystis* Haeckel,1881
* Correct family?
- A. anthemis* Haeckel,1887
off Ålesund
- Stephoidea* Haeckel
- Stephanidae* Haeckel
- Semantidae* Haeckel,1887
- Semantis* Haeckel,1887
- S. distephanus* Haeckel,1887
Bergen, 56°N 23°W
- S. micropora?* Popofsky,1908
Bergen
- Dictyocircus* Jørgensen,1905
- D. clathratus* Jørgensen,1905
Bergen, W Norwegian fjords, Henningsvær
- Coronidae* Haeckel
- Tympanidae* Haeckel
- Spyroidea* Haeckel,
- Zygospyridae* Haeckel
- Ceratospyris* Ehrenberg,1847
- C. hyperborea* Jørgensen,1905
Bergen, W Norwegian fjords
- Tholospyridae* Haeckel
- Phormospyridae* Haeckel
- Androspyridae* Haeckel
- CYRELLARINA = CYRTINA = PROAXOPLASTIDIATA Cachon & Cachon
- Cyrtoidea* Haeckel,1862, emend. Petrushevskaja,1971
- ?Family?
- Cornutella* Ehrenberg,1838
- C. profunda* Ehrenberg,1854
Bergen, W Norwegian fjords, Mediterranean, off E Greenland
- C. annulata* Ehrenberg,1861
Greenland
- Monocyrtoidea* Haeckel
- Euscenium* Haeckel,1887
- E. corynephorum* Jørgensen,1900
W Norwegian fjords
- Cladoscenium* Haeckel,1882
- C. tricolpium* (Haeckel,1881)
= *Archiscenium tricolpium* Haeckel,1881
= *Euscenium tricolpium* : Auctt.
Bergen, W Norwegian fjords, Lofoten, Finnmark, Spitsbergen, between Norway & Shetlands
- C. limbatum* Jørgensen,1905
Bergen, Gaukvaerø
- C. sp.* : Björklund,1974
Bergen
- Pteroscenium* Haeckel,1881
- P. arcuatum* Haeckel,1887
off Biscay
- Dicyrtoidae* Haeckel
- Litharachnium* Haeckel,1860
- L. tentorium* Haeckel,1862
Bergen, between Norway & Iceland, Mediterranean
- L. arachnodiscus* (Ehrenberg,1861)
= *Carpocanium arachnodiscus* Ehrenberg,1861
between Iceland & Greenland
- Sethopilium* Haeckel,1881
- S. meunieri* Schröder,1911
Arctic Ocean, Kara Sea
- Lithomelissa* Ehrenberg,1847
- L. setosa* Jørgensen,1900
Belt Sea, Kattegatt, Skagerrak, Bergen, W Norwegian fjords
- L. laticeps* Jørgensen,1905
Bergen, between Shetlands & Faeroes
- L. bicornis* Ehrenberg,1861
Bergen, Davis Strait
- L. thoracites* Haeckel,1862
Bergen, W Norwegian fjords, Mediterranean
- L. hystrix* Jørgensen,1900
Bergen, W Norwegian fjords
- L. sp. : Swanberg & Björklund,1987*
W Norwegian fjords
- Tricyrtoidae* Haeckel
- Theopilium* Haeckel,1881
- T. tricostatum* Haeckel,1887
56°N 10-30°W
- Corocalyptra* Haeckel,1887
- C. craspedota* (Jørgensen,1900)
= *Theocalyptra craspedota* Jørgensen,1900
North Sea, Bergen, W Norwegian fjords, Lofoten
- C. elisabethae* Haeckel,1887
56°N 23°W
- Pterocorydidae* Haeckel,1881, emend. Riedel,1967
- Pterocorys* Haeckel,1881
- P. irregularis* Cleve,1899
Spitsbergen, off Ålesund, 56°N 23°W
- Androcyclas* Jørgensen, in Nordgaard & Jørgensen,1905
- A. gamphonycha* (Jørgensen,1900)
= *Pterocorys gamphonychos* Jørgensen,1900
?= *Pterocorys theoconus* Jørgensen,1900
Bergen, W Norwegian fjords
- Lamprocyclas* Haeckel,1881
- L. aegles* (Ehrenberg,1854)
= *Podocyrts aegles* Ehrenberg,1854
off Stadt
- Stichocystoidae* Haeckel
- Stichocorys* Haeckel,1882
- S. seriata* (Jørgensen, in Gran,1902)
= *Eucyrtidium seriatum* Jørgensen, in Gran,1902
North Sea, Bergen, W Norwegian fjords
- Eucyrtidium* Ehrenberg,1847
- * Correct family?
- C. davisianna* Ehrenberg,1872
= *Pterocodon davisianna* : Ehrenberg,1872
= *Eucyrtidium davisiannum* : Haeckel,1862
= *Pterocanum davisiannum* : Haeckel,1862
= *Stichopilum davisiannum* : Auctt.
W Norwegian fjords, Spitsbergen, off E Greenland
- C. tabulata* Ehrenberg,1872
= *Sethocomas tabulatus* : Auctt.
?Arctic Ocean
- Lithostrobus* Haeckel,1887
- * Correct family?
- L. cuspidatus* (Ehrenberg,1872)
= *Eucyrtidium cuspidatum* Ehrenberg,1872
Greenland
- L. cornutella* (Ehrenberg,1872)
= *Eucyrtidium cornutella* Ehrenberg,1872
Arctic Ocean, Mediterranean
- L. botryocystis* Haeckel,1887
50°N 31°W, 55°N 44°W
- Lithomitra* Haeckel,1887
- L. lineata* (Ehrenberg,1838)
= *Lithocampe lineata* Ehrenberg,1838
= *Eucyrtidium lineatum* : Ehrenberg,1847
= *Dictyonitra lineata* : Stöhr,1880
N. Ice Sea, off E Greenland, Mediterranean
- L. hyperborea* (Bailey,1856)
= *Eucyrtidium hyperboreum* Bailey,1856
= *Lithocampe hyperborea* : Haeckel,1862

- Greenland
- L.* sp. : (Cleve,1899)
= *L. australis* : Cleve,1899
Spitsbergen
- Botryostrobus** Haeckel,1887
- B. platycephalus** (Ehrenberg,1872)
= *Eucyrtidium platycephalum* Ehrenberg,1872
= *Lithocampe platycephala* : Auctt.
W Norwegian fjords, N Atlantic
- Lithocampe** Ehrenberg,1839
- L. eupora** (Ehrenberg,1872)
= *Eucyrtidium euporum* Ehrenberg,1872
N Atlantic
- L. hispida** (Ehrenberg,1861)
= *Eucyrtidium hispidum* Ehrenberg,1861
N Ice Sea
- L. tumidula** (Bailey,1856)
= *Eucyrtidium tumidulum* Bailey,1856
= *Eucyrtidium increscens* Ehrenberg,1861
N Atlantic, N Ice Sea
- L. quadrarticulata** (Ehrenberg,1861)
= *Eucyrtidium quadrarticulatum* Ehrenberg,1861
Greenland
- Artostrobus** Haeckel,1887
- A. annulatus** (Bailey,1856)
= *Cornuella annulata* Bailey,1856
= *Eucyrtidium annulatum* : Haeckel,1862
between Greenland and Jan Mayen
- A. joergensei** Petrushevskaya,1967?
off E Greenland
- Botryoidea** Haeckel
- Cannabotryidae** Haeckel,1881, emend. Riedel,1967
- Botryocytis** Ehrenberg,1860
- B. platycephala** (Petrushevskaya,1967)
off E Greenland
- Amphimelissa** Jørgensen,1905
- A. setosa** (Cleve,1899)
= *Botryopyle setosa* Cleve,1899 (TM)
Skagerrak, Hardangerfjord, Bergen, Sognefjord, Tromsø, Norw. Sea, Barents Sea, Kara Sea, off SE Iceland, between Iceland & Greenland
- A. stenostoma** (Meunier,1910)
= *Botryopyle stenostoma* Meunier,1910
Barents Sea, Kara Sea, Arctic Ocean
- Lithobotryidae** Haeckel
- Botryocella** Haeckel,1881
- B. borealis** (Ehrenberg,1872)
= *Lithobothrys borealis* Ehrenberg,1872
Greenland
- Pyllobotryidae** Haeckel
- Botryocampe** Ehrenberg,1860
- B. inflata** (Bailey,1856)
= *Lithobothrys inflata* Bailey,1856
Greenland, Mediterranean
- PHAEODAREA** Haeckel,1879 emend. Haecker, Reschetnjak
= **TRIPYLEA** Hertwig,1879
= **CANNOPYLEA** Haeckel,1881
= **PANSOLENA** Haeckel,1878
- Lirella** Ehrenberg,1872
- L. melo** (Cleve,1899)
= *Beroetta melo* Cleve,1899
Bergen, Rost, W Spitsbergen, Greenland Sea
- L. bullata** (Stadum & Ling,19???)
W Norwegian fjords
- Borgerella** ?19??
- B. caudata** (Wallich,18???)
W Norwegian fjords
- PHAEOGYMNOCELLIDA** Cachon & Cachon,1982
- Phaeosphaeridae** Cachon-Enjumet
- Phaeosphaera** Cachon-Enjumet
- Phaeodinidae** Cachon-Enjumet
* Monogenic
- Phaeodina**
- Atlanticellidae** Cachon-Enjumet
* 5 genera
- Miracella** Borgert
- Gymnocella** Cachon-Enjumet
- Planktonetta** Borgert
- Halocella** Borgert
- Lobocella** Borgert
- PHAEOCYSTIDA**
- Aulacanthidae** Haeckel
* 11 genera
- Aulacantha** Haeckel,1861
- A. scolymantha** Haeckel,1862
Bergen, Faeroe Channel, off W Hebrides, W Greenland, Mediterranean
- A. laevissima** Haeckel,1887
Faeroe Channel, Greenland, Spitsbergen
- A. tenuis** Haeckel,18???
?off W Hebrides
- Aulographidae** Haeckel,1884
= *Aulographium* Haeckel,1879
- A. flosculus** Haeckel,1887
Faeroe Channel, N Gulf Stream branch
- A. tetraneustra** Haeckel,1887
50.8-60.3°N 27-47.3°W
- A. pulvinata** Haeckel,1887
Faeroe Channel, N Gulf Stream branch
- A. zetesios** (Fowler,1898)
= *Aulocoryne zetesios* Fowler,1898
Faeroe Channel, N Gulf Stream branch
- Auloceros** Haeckel,1887
- A. trigeminus** Haeckel,1887, var. Fowler,1898
Faeroe Channel, N Gulf Stream branch
- Aulospathis** Haeckel,1887
- A. bifurca** Haeckel,1887
60.3°N 27°W
- Aulodendron** Haeckel,1887
- A. antarcticum** Haeckel,1887
Spitsbergen, Greenland Sea
- Astracanthidae** Haeckel
* 1 genus
- Astracantha** Haeckel
- PHAEOSPHAERIDA**
- Aulosphaeridae** Haeckel
* 4 genera
- Aulosphaera** Haeckel,1860
- A. flexuosa** Haeckel,1887
Faeroe Channel, N Gulf Stream branch
- A. labradoriensis** Borgert,1901
Irminger Sea, W Greenland
- A. multifurca** Haeckel,1887
Greenland, Arctic Ocean
- A. neglecta** Borgert,1901
Mediterranean, Irminger Sea
- Auloscena** Haeckel,1887
- A. spectabilis** Haeckel,1887
W Spitsbergen, Greenland Sea
- A. verticillata** Haeckel,1887
Irminger Sea
- Aulatractus** Haeckel,1887
= *Aulotractus* Hensen,1887
- A. fusulus** Haeckel,1887
between Hebrides & Rockall, Faeroe Channel, N Gulf Stream branch
- A. ovulum** Borgert,1901
Irminger Sea
- A. septentrionalis** Borgert,1901
Irminger Sea
- A. fusiformis** Haeckel,1887
Irminger Sea
- A. proteus** Borgert,1901
Labrador Current
- Aulastrum** Haeckel,1887
- A. spinosum** Borgert,1901
Labrador Current
- A. mirabile** Borgert,1901
Irminger Sea
- Cannosphaeridae** Haeckel
* one genus
- Coelacanthatha** Hertwig,1879
= *Cannosphaera* Haeckel,1879
- C. lepta** (Jørgensen,1900)
= *Cannosphaera lepta* Jørgensen,1900
Hardangerfjord, Bergen, Foldenfjord, Skjerstadfjord, Svolver, Skroven, Henningsvær, Balstad, Tranøybet
- C. geometrica** (Borgert,1892)
= *Cannosphaera geometrica* Borgert,1892
Skroven, Vestfjord, Kvenangen, Glaukvær, Irminger Sea
- C. antarctica** (Haeckel,1887)
= *Cannosphaera antarctica* Haeckel,1887
Bergen, Irminger Sea, W Greenland
- Sagospheridae** Haeckel

- * 3 genera
- Sagoscena** Haeckel,1887
- S. castra* Haeckel,1887
Irminger Sea
- S. militaris* Borgert,1901
Irminger Sea
- S. elegans* Borgert,1891
Irminger Sea
- Sagenoscena* Haeckel,1???
- S. penicillata* Haeckel,1887
W Spitsbergen, Greenland Sea
- S. irmingeriana* Borgert,1901
Irminger Sea
- S. spathillata* Haeckel,1887
Greenland, Arctic Ocean
- Sagenoarium* Borgert,1891
- S. chuni* Borgert,1891
Irminger Sea
- S. norvegicum* Brock,1906
= *Sagenoarium* (?) *norvegicum* Brock,1906
62°20'N 0°0'E
- S. sp.* : Jørgensen,1900
Bergen
- PHAEOCALPIDA**
- Castanellidae** Haeckel
* 7 genera
- Castanidium* Haeckel,1879
- C. variabile* Borgert,1891
Mediterranean, Irminger Sea
- C. apstenei* Borgert,1901
Irminger Sea
- C. brandti* Borgert,1901
between Faeroes & Rockall, N Gulf Stream branch
- Castanarium* Haeckel
- Castanella* Haeckel
- Castanissa* Haeckel
- Castanura* Haeckel
- Castanea* Haeckel
- Circocastanea* Haeckel
- Circoporidae** Haeckel
* 2 genera
- Haeckeliana* Haeckel,1887
- H. labradoriana* Borgert,1901
Labrador Current
- Circospathis* Haeckel
- Tuscaroridae** Haeckel
* 5 genera
- Tuscarora* Murray,1885
- T. nationalis* Borgert,1892
Irminger Sea
- Tuscareta* Haeckel
- Tuscarilla* Haeckel
- Porospathidae** Borgert
* one genus
- Porospathis* Haeckel,1879
= *Polyptetta* Haeckel,1887
- P. holostoma* (Cleve,1899)
= *Polyptetta holostoma* Cleve,1899
W Spitsbergen, Greenland Sea
- Polypyramidae** Reschetyjak
* one genus
- Polypyramis* Reschetyjak
- PHAEOGROMIDA**
- Challengeridae** Murray,1876
* Hensen,1887 has a note about a probably new species of the family from Skagerrak.
* 3 genera
- Cadiumella* Strand,1928
= *Codium* Bailey,1856, non *Link*,1807 (Mollusca)
- C. marinum* (Bailey,1856)
= *Codium marinum* Bailey,1856
Labrador Current, Kamtschatka, Arctic Ocean
- Challengeron* Haeckel,1887
- C. diodon* Haeckel,1887
= *C. nathorstii* Cleve,1899
= *C. heterocanthum* Jørgensen,1900
Gullmarn, Skagerrak, Bergen, Foldenfjord, Skroven, Henningsvær, Balstad, Tysfjord, Lofoten, Sagfjord, Øksund, Raftsund, Tranøybet, Ofoten, Skjomen, Vestfjord, Raftsundet, Tromsø, Gaukvær, Stamsund, Helligvær, Bear Island, Iceland, Greenland Sea, Barents Sea, W Spitsbergen, Mediterranean
- C. armatum** Borgert,1901
Bergen, Gaukvær, between Hebrides & Rockall, N Gulf Stream branch
- C. neptuni* Borgert,1892
Labrador Current
- C. triodon* Haeckel,1887
off W Hebrides
- C. balfouri* (Murray,1885)
= *Challengeria balfouri* Murray,1885
off W Hebrides
- C. golfense* Haeckel,1887
Faeroe Channel, N Gulf Stream branch
- C. channeri* (Murray,1885)
= *Challengeria channeri* Murray,1885
between Faeroes & Rockall
- C. johannis* Haeckel,1887
Faeroe Channel, N Gulf Stream branch
- Lithogromia* Haeckel,1879
* This genus is not a member of the family, if it not is a synonym of one of the three.
- L. silicea* Haeckel,1887
Faeroe Channel, N Gulf Stream branch
- Protocystis** Wallich,1862
- P. tridens* (Haeckel,1887) Borgert,1901
Skagerrak, Hardangerfjord, Bergen, Foldenfjord, Skjerstadfjord, Svolver, Skroven, Henningsvær, Balstad, Tranøybet, Rost, Øksund, Ofoten, Lyngen, Kvænangen, Helligvær, Vestfjord, Gaukvær, between, Hebrides & Rockall, N Gulf Stream branch, E & W Greenland, Barents Sea, North Ice Sea
- P. xiphodon* (Haeckel,1887) Borgert,1901
= *Challengeria xiphodon* Haeckel,1887
Bergen, Skroven, Tysfjord, Rost, Raftsund, Øksund, Sagfjord, Henningsvær, Tranøybet, Ofoten, Vestfjord, Raftsundet, Tromsø, Kvænangen, Helligvær, Glaukvær, between Hebrides & Rockall, N Gulf Stream branch, E & W Greenland, Mediterranean
- P. harstoni* (Murray,1885) Borgert,1901
= *Protocystis aurita* Wallich,1869
= *Challengeria harstoni* Murray,1885
= *Challengeria thomsoni* Hensen,1887
Bergen, Sognefjord, Skroven, Rost, Tysfjord, Øksund, Tranøybet, Ofoten, Skjomen, Glaukvær, off W Hebrides, N Gulf Stream branch, S & W Spitsbergen, Greenland Sea
- P. tritonis* (Haeckel,1887)
= *Challengeria tritonis* Haeckel,1887
Faeroe Channel, N Gulf Stream branch
- P. bicornis* Borgert,1901
Labrador Current
- Entocannula** Haeckel,1879
* This genus is not a member of the family, if it not is a synonym of one of the three.
- E. hirsuta* Haeckel,1887
Faeroe Channel, N Gulf Stream branch
- Pharyngella** Haeckel,1887
* This genus is not a member of the family, if it not is a synonym of one of the three.
- P. gastrula* Haeckel,1887
Faeroe Channel, N Gulf Stream branch
- Medusettidae** Haeckel,1887
* 6 genera
- Medusetta** Haeckel,1887
- M. arcifera* Jørgensen,1900
Hardangerfjord, Bergen, Sognefjord, Reine, Ofoten, Tromsø, Henningsvær
- M. tiara* Haeckel,1887
Faeroe Channel, N Gulf Stream branch
- M. armata* Borgert,1901
Labrador Current
- Gazelletta* Haeckel,1887, ex Murray MS
- G. pentapodium* Jørgensen,1900
Bergen, Henningsvær, Rost, Vestfjord, Tromsø, Moskenstrømmen
- G. hexanema* Haeckel,1887
off S Orkneys
- G. pectinata* Haeckel,1887
Faeroe Channel, between Hebrides & Rockall, N Gulf Stream branch
- G. atlantica* Borgert,1901
between Faeroes & Rockall, Irminger Sea
- Euphysetta** Haeckel,1887
- E. nathorsti* Cleve,1899
Bergen, W Spitsbergen, Greenland Sea
- E. lucani* Borgert,1892
= *E. mediterranea* Lohmann,1899
Mediterranean
- Cortinetta** Haeckel
* 3 genera
- Gorgonetta** Haeckel
- Polyptetta** Haeckel
- PHAEOCONCHIDA**
- Concharidae** Haeckel
* 9 genera. Not recorded north of 50° N in the Atlantic
- PHAEODENDRIDIDA**
- Coelodendridiae* Haeckel,1862
* Many genera
- Coelodendrum* Haeckel,1860

- C. ramosissimum* Haeckel,1860
Bergen, Faeroe Channel, N Gulf Stream branch, Mediterranean, Irminger Sea
- Coeloplemma* Haeckel,1887
- C. murrayanum* Haeckel,1887
Faeroe Channel, off W Hebrides, N Gulf Stream branch
- C. tritonis* Haeckel,1887
Faeroe Channel, N Gulf Stream branch
- "ACTINELIDA" Popofsky,1904 emend. Mielk & Schewiakoff
* Incertae sedis. Possibly transitional between Acantharia & Heliozoa.
- ACTINELINA Haeckel,18?? emend. Popofsky, Mielk & Schroeder
- Astrolophidae Haeckel
- Actinelius* Haeckel,1865
- Astrolophus* Haeckel,1881
- Podactinelius* Schröder,1907
- ACTINASTRINA Popofsky
- Trizonidae Popofsky
- Trizone* Popofsky,1904
- Actinastridae Haeckel,18?? emend. Popofsky
- Actinastrum* Haeckel,1885
- HELIozoa** Haeckel,1866
* Strongly polyphyletic. Now divided in eight distantly related groups: Actinophryida, Centrohelida, Cristidiscoidida (Rhizopoda, Filosea), Desmothoracidae (incertae sedis), Dimorphidae (incertae sedis), Gymnosphaerida, Pedinellidae & Taxopodida.
- Servetia* Poche,1913
= *Haeckelia* Mereschkowsky,1878, non Bessels,1875 (Foraminiferida)
= *Microlis* Dons,1918
* which order?
- S. borealis* (Mereschkowsky,1878)
= *Haeckelia borealis* Mereschkowsky,1878 (TM)
= *Podophrys gracilis* Calcins,1902
Norway, White Sea, E USA
- Gelatinophrys* ?Kufferath,1952
* which order?
- G. ostendensis* Kufferath,1952
* planktonic
Ostende
- ACTINOPHYRYIDA Kühn,1926
- Family?
- Camptonema* Schaudinn,1894
- C. nutans* Schaudinn,1894 (TM)
* Not seen since its discovery.
Norwegian coast at 10 m.
- Actinophryidae
- Actinophrys* Ehrenberg,1830 (TM *A. sol* Ehrenberg,1830)
- A. pontica* Valkanov,1940
Black Sea
- Actinosphaeridae
- Actinosphaerium* von Stein,1857
= *Echinosphaerium* ?,?!
- A. portuum* Kufferath,1952
* planktonic
Ostende
- DESMOTHORACIDA** Hertwig & Lesser,1874
* Taxon incertae sedis, according to Patterson & Zöllfel,1992
- Family?
- Hedriocystis* Hertwig & Lesser,1874
- Monostigocystis* Saedeleer,1930
- Clathrulinidae
- Clathrulina* Cienkowsky,1867
- Orbulinella* Entz senior,1877
- O. smaragdea* Entz senior,1877 (TM)
?= *O. salina* Labbé,1924
* in salty ponds
Hungary, Croise
- CENTROHELIDA** Hartmann,1913 = **CENTROHELIOZOA**
* Should the author be Kühn,1926
- Family?
- Chlamydaster* ?,???
- Raphidocystis* Pénaud,1904
- Actinolophus* Schulze,1874
- A. pedunculatus* Schulze,1874 (TM)
Warnemünde, S France
- Cienkowskya* Schaudinn,1896
= *Wagneria* Cienkowsky,1881, non Robineau-Desvoidy,1830, nec Gistl,1848, nec Alenitzin,1873
- C. mereschkowskii* (Cienkowsky,1881)
= *Wagneria mereschkowskii* Cienkowsky,1881 (TM)
White Sea, S France
- Wagnerella Mereschkowsky,1878
- W. borealis* Mereschkowsky,1878 (TM)
Norway, cosmopolitan
- Oxnerella* Dobell,1917
- O. marina* Dobell,1917 (TM)
Plymouth
- Actinocoryne Febvre-Chevalier,1980
- A. contractilis* Febvre-Chevalier,1980 (TM)
French Mediterranean
- Hedraiophryidae
- Hedraiophysys* Febvre-Chevalier,1973
- H. hovassei* Febvre-Chevalier,1973 (TM)
?French Mediterranean
- Gymnosphaeridae
- Gymnosphaera* Sasaki,1894
- G. albida* Sasaki,1894 (TM)
Mediterranean
- Acanthocystidae
- Acanthocystis* Carter,1863 (TM *A. turfacea* Carter,1863 - fresh water)
* This genus was divided into three genera (*Acanthocystis*, *Choanocystis* & *Pterocystis*) for ultrastructural reasons by Siemensma & Roijackers,1988. This was thought to be a bit premature by Patterson & al., 1993, until ecological studies separating the three groups in one ore another adaptive way is found.
- A. erinaceoides* Petersen & Hansen,1960
Tåvrminne, Danish & Dutch fresh water
- A. fortessa* Nicholls,1983
Tåvrminne, German & Dutch fresh water
- A. paliformis* Dürrschmidt,1987
Tåvrminne, Fresh waters of Chile & New Zealand
- A. pelagica* Ostenfeld,1904
* Autumn plankton form.
Baltic, Kattegatt, Skagerrak, W Jylland, W Limfjord
- A. perpusilla* Petersen & Hansen,1960
= *Choanocystis perpusilla* (Petersen & Hansen,1960) Siemensma & Roijackers,1988
Tåvrminne, Denmark, cosmopolitan, eurythermal in fresh water
- A. (Acanthocystis) turacea* Hertwig & Lesser,1874
Bothnian Sea, Denmark
- A. spinifera* Greeff,1869, non Penard,1904
* Freshwater species in Europe & North America, which also may be found pelagical in Seawater.
?
- A. wiesemskii* Ostroumoff,1917
* marine species
?
- Heterophryidae
- Heterophrys* Archer,1869
- H. myriopoda* Archer,1869 (TM - fresh water)
= *H. marina* Hertwig & Lesser,1874
Tåvrminne, German seawater aquarians, Fresh water in most of Europe
- Astrodisculus* Greeff,1869 sensu Penard,1904, non Stern,1924
- A. marinus* Kufferath,1952
* planktonic
Ostende
- Raphidiophryidae
- Raphidiophysys* Archer,1867
- R. marina* Ostenfeld,1904
* planktonic during October - January
Skagerrak, S & E North Sea, Engl. Channel, Shetland, Irish Sea
- R. marginata* Nicholls & Dürrschmidt,1985
Tåvrminne, Fresh water in Netherlands, Canada, Chile, New Zealand ...
- TAXOPODIIDA Fol,1883
- Sticholonchidae
- Sticholonche* Hertwig,1877
- S. zanklea* Hertwig,1877 (TM)
= *Spermatogeton antiqua* Fortmorel,1892
= *Sticholonche ventricosa* Meunier,1910
* pelagic at depths of 100-500 m. Often found with dinoflagellate parasite *Amoeboophrya sticholonchae* Koeppen
* Found in Scandinavian waters during winter (Nov.-Feb.) only.
Kattegatt, Skagerrak, North Sea, Barents Sea, Arctic Ocean, Mediterranean
- ROTOSPHAERIDA Rainer,1968
- Pinaciophora* Greeff,1875 (TM *P. fluviatilis* Greeff,1873 - fresh water)
= *Potamodiscus* Gerloff,1968?
- P. tridentata* Thomsen,1978
Tåvrminne, Denmark (Isfjord, Dybsø fjord)
- P. bifurcata* Thomsen,1978
Denmark (North Sea)
- Pinacocystis* Hertwig & Lesser,1874
?= *Pinaciocystis* Roskin,1929
- P. rubicunda* Hertwig & Lesser,1874
a Seawater aquarium
- Lithocolla* Schulze,1874

L. globosa Schulze,1874 (TM)

* Euryecous species, which may be found in fresh as well as brackish water.
Warinemünde

CHYTRIDIOMYCOTA

* in S Norway is *Thalassiosira nordenskioldii* Cleve parasited by a marine chytrid (Gaertner,1979)

Nowkowskella ??,1???

* Which order?

Sporomonas Chatton & Lwoff,1924

* Systematic placement very tentative. This flagellate is difficult to classify

S. infusoriorum Chatton & Lwoff,1924

@ internal parasite in *Folliculina* & *Vorticella*
France, Woods Hole

Topsisentella Duboscq & Tuzet,1936

* Systematic placement very tentative.

T. fallax Duboscq & Tuzet,1936

@ parenchyma of *Sycon raphanus*

? France

Nephrococcidioides Harant,1936

* Which order?

N. legeri (Harant,1931)

= *Pseudoklossia legeri* Harant,1931
@ pyloric glands of *Styela* & *Polycarpa* spp.
? France

CHYTRIDIALES

Achlyogotonaceae

Achlyogeton Schenk,18??

A. entophytum Schenk,18??

@ *Cladophora*

* marine?. Occurs primarily in fresh water *Cladophora* species.
?

A. satinum P.A. Dangeard,1932

@ *Cladophora* sp.

?

Rhizidiaceae

Catenochytridium Berdan,1939

* Correct family?

C. carolinianum Berdan, forma *marinum* Kobayasi & Ookubo,1953
@ *Cladophora japonica* Yamada
Japan

Chytridium Braun,1851

C. polysiphoniae Cohn,1865

= *Rhizophyllum polysiphoniae* (Cohn) Petersen,1905
= *Rhizophyllum olla* Petersen,1905
@ *Polysiphonia*, *Ceramium*, *Delesseria* & other red algae
Swedish coast, Denmark, Germany, Greenland, W USA, France

C. megastomum Sparrow,1934

@ *Striaria attenuata* & *Ceramium diaphanum*
Denmark, North Carolina

C. turbinatum Kobayasi & Ookubo,1954
@ *Bryopsis* sp.
Japan

C. codicola Zeller,1918
@ *Codium mucronatum*
North Carolina

C. lagenaria var. *japonense* Kobayasi & Ookubo,1953
@ *Cladophora japonica* Yamada
Japan

Entophyscylctis Fischer,1???

E. confervae-glosteratae (Cienkowski) Sparrow, forma *marina* Kobayasi & Ookubo,1954
* should be moved from this genus
Japan?

Phlyctidium (Braun) Rabenhorst,18??

P. brevipes
= *P. brevipes* var. *marinum* Kobayasi & Ookubo,1954
@ *Gracilaria*
* originally known from fresh water
Japan?

Physorhizopodium Scherffel,1???

P. pachydermum
@ *Amphora* & *Navicula*
?

Phlyctochytrium Schroeter,1892
= *Rhizidium* A. Fischer, in Rabenhorst,1892, non Braun,1856
@ filamentous algae

P. aestuarii Ulken,1972
Weser estuary

P. mangrovis Ulken,1972
Brazil, Hawaii

P. marinum Kobayasi & Ookubo,1953
@ *Cladophora* spp.
Japan

P. cladophorae Kobayasi & Ookubo,1954
@ *Cladophora* sp.

P. bryopsisidis Kobayasi & Ookubo,1954
@ *Bryopsis* sp. & *Cladophora* sp.
Japan

P. japonicum (Kobayasi & Ookubo) Sparrow,1960

= *Rhizophyllum japonicum* Kobayasi & Ookubo,1953

@ *Cladophora* sp. & *Gracilaria confervoides*

Japan

P. semiglobiferum Uebelmesser,1956

* not truly marine

European coast

Podochytrium Pfister,1869

P. chitinophilum Willoughby,1961

Tay estuary at very low salinity

Rhizidium Braun,1856

R. braunii Zopf,1888

@ diatoms & desmids

Finland, Germany (inland salt pool)

Rhizoclosmatium Petersen,1903

R. marinum Kobayasi & Ookubo,1954

@ *Codium fragile*

Japan

Rhizophlyctis Fischer, in Rabenhorst,1892

= *Karlingia Johanson*,1944

R. harderi Uebelmesser,1956

Tay estuary, Mediterranean

Rhizophydiuum Schenk,1858

R. planktonicum

@ *Asterionella*

?

R. littoreum Amon,1984

@ *Codium* sp. & *Cancer antonyi*

Hawaii

R. halophilum Uebelmesser,1956

@ Pine pollen

Baltic, North Sea, Mediterranean, USA

R. codicola Zeller,1918

@ *Codium mucronatum*

USA

R. subglobosum Kobayasi & Ookubo,1954

@ *Bryopsis* sp. & *Cladophora* sp.

Japan

R. cladophorae (Kobayasi & Ookubo) Sparrow,1960

= *Phlyctidium cladophorae* Kobayasi & Ookubo,1954

@ *Cladophora* sp.

Japan

R. distinctum Petersen,1905

@ e.g. *Acrosiphonia Polysiphonia*, *Ceramium* & *Spongomerpha*

* doubtful species

Sweden, Denmark, USA

R. gelatinosum Lind,1905

@ *Acrosiphonia pallida*

* Doubtful species, which may not be a fungus

Sweden

R. globosum (Braun) Rabenhorst,18???

@ e.g. ?*Acrochaetium* & *Bryopsis plumosa*

* Doubtful species, which may embrace a collection of taxa

Sweden, USA

? *R.* sp. : Drebels,1974

@ *Bacteriastrum hyalinum* & *Chaetoceros densus*

North Sea

Spizellomyces

S. acuminatus (Ulken,1972) (Barr,1984)

Weser estuary

Olpidiaceae

Olpidium (Braun) Rabenhorst,1868

O. laguncula Petersen,1905

@ *Dumontia incrassata* & *Dumontia filiformis*

* doubtful species

Swedish coast, Denmark, Isle of Man

O. maritimum Höhnk & Aleem,1953

?= *O. pendulum* Zopf,1???

* in brackish water

Sweden, ?Bohusl., Weser estuary

O. aggregatum P.A. Dangeard,1890-91

@ *Cladophora* sp.

* may possibly be a synonym of *Siroolidium bryopsisidis*

France, Japan, North Carolina

O. entosphaericum (Cohn) A. Fischer, in Rabenhorst,1892

= *Chytridium entosphaericum* Cohn,1865

@ *Bangia fuscopurpurea* & *Hormidium penicilliformis*

* should possibly be transposed to *Anisolpidium* or *Olpidiopsis*

Germany

O. lauderiae Gran,1900

@ *Lauderia borealis*

* should probably be transposed to *Ectrogella* or *Eurychasma*

Norway

O. spp. : Jirovec,1956

@ in a marine rotifer species

?

Rozella Cornu,1872

= *Rozia* Cornu,1872

= *Pleolpidium* A. Fischer, in Rabenhorst,1892

R. marina Sparrow,1938

@ *Chytridium polysiphoniae* Cohn

USA

Pleotrichelus Zopf,1884*P. olpidium* Petersen,1905@ algae, e.g. *Ectocarpus* spp. *Furcellaria*, *Polysiphonia* etc.
Swedish coast, Denmark, Faeroes, France*P. minutus* Petersen,1905@ *Chorda filum* & *Litosiphon pusillus*
Swedish coast, Denmark, France*P. inhabilis* Petersen,1905@ *Polysiphonia* spp.
* dubious species, seemingly consisting of different forms, belonging in *Petersenia* or *Olpidium*
Denmark*P. paradoxus* Petersen,1905@ in sporangia of *Rhizophydium distinctum* on *Spongomorpha vernalis*
NorwayCladophytriaeae*Coenomyces* Deckenbach,1902-1903
= *Deckenbachia* Jacewski,1931*C. consuens* Deckenbach,1902-1903 (TM)
@ marine blue green bacterians, e.g. *Calothrix*, *Rivularia* & *Tildenia*
Denmark, Albania, USA*Nephromyces* Giard,1888

* imperfectly known genus, very provisionally regarded as a member of this family

N. molgularum Giard,1888@ renal sac of *Molgula tubifera*
?France*N. sorokinii* Giard,1888@ renal sac of *Molgula complanata*
?France*N. rocovitanus* Giard,1888@ renal sac of *Molgula oculata*
?FranceHARPOCHYTRIALES

* No marine species?

BLASTOCLADIALESCatenariaceae*Catenaria* Sorokin,1876*C. anguillulae* Sorokin,1876

Mexico, India

MONOBLEPHARIDALES

* No marine species?

PHYCOMYCOTA**= MASTIGOMYCOTINA***Cycloptericola* ?Apstein,1910?

* Taxonomic position unclear

C. marina Apstein,1910@ oral cavity wall of *Cyclopterus lampus*
?OOMYCOTEAESAPROLEGNIIDA= SAPROLEGNIALESSaprolegniaceae*Leptolegnia* de Bary,1888*L. baltica* Höhnk & Vallin,1953
@ *Eurytemora hirundoides* (Nordquist)
N BalticEctrogellaceae*Eurychasma* Magnus,1905

@ Parasitic principally in Phaeophyta & Rhodophyta

E. dicksonii (Wright) Magnus,1905= *Rhizophydium dicksonii* Wright,1879= *Olpidium dicksonii* (Wright) Wille,1899= *Ectrogella dicksonii* (Wright) Scherffel,1925@ *Ectocarpus* spp., *Pyliella littoralis* & other algae

W Sweden, Denmark, Ireland, Britain, France, Greenland, Mediterranean, Kerguelen

E. sacculus Petersen,1905= *Palmaria palmata* & *Halosaccion ramentaceum*

* Doubtful species

North Sea, Greenland

Eurychasmidium Sparrow,1936

@ Parasitic principally in Phaeophyta & Rhodophyta

E. tumefaciens (Magnus) Sparrow,1936= *Chytridium tumefaciens* Magnus,1872= *Olpidium tumefaciens* (Magnus) Wright,1879= *Pleotrichelus tumefaciens* (Magnus) Petersen,1905@ nodal cells of *Ceramium* spp.

Denmark, Scotland, Ireland, France, Italy, USA

Ectrogella Zopf,18??, em. Scherffel,1925

@ Diatoms

E. perforans H.E. Petersen,1905@ *Licmophora* spp. & other pennate diatoms

Sweden, Denmark, Germany, France, USA

E. eurychasmoides Feldmann & Feldmann,1955@ *Licmophora lyngbyei*

France

E. licmophorae Scherffel,1925@ *Licmophora* sp.

Italy

E. marina (P.A. Dangeard) Feldmann & Feldmann,1955= *Olpidium marinum* P.A. Dangeard,1912

* doubtfully member of the genus

?

Leptolegniellaceae Dick,1971*Leptolegniella* Huneycutt,1???*L. marina* (Atkins,1954) Dick,1971?= *Leptolegnia marina* Atkins,1954@ eggs & gills of *Pinnotheres pisum* in *Barma candida* & *Cerastoderma echinata* & on eggs of *Callinectes sapidus*

Britain

Atkinsiella Vishniac,1958*A. dubia* (Atkins,1954) Vishniac,1958= *Plectospira dubia* Atkins,1954 (TM)@ Crustacean eggs, zoeas & pre-zoceans, e.g. *Macropodia*, *Liocarcinus* & *Crangon*

Britain

A. hamanaensis Bian & Egusa,1980@ ova of Mangrove crab, *Scylla serrata* (Forsskål)

Japan

LAGENIDIIDA= LAGENIDIALESFamilies?

?

? thalassium (Atkins,1955)= *Pythium thalassium* Atkins,1955@ *Pinnotheres pisum*

* Provisionally classified with Lagenidiida

Britain

? afertile (?1???)= *Pythium aferile* ?1???@ *Palaeomon serratus*

Britain

Gonimochaete*G. latitibus* Newell, Cefalu & Fell,1977

@ parasiting & killing littoral nematodes

?

Haptoglossa*H. heterospora* Drechsler,1???

@ parasiting & killing littoral nematodes

?

Lagenidiaceae*Lagenidium* Schenk,1859= *Resticularia* P.A. Dangeard,1890-91

@ eggs & juveniles of Crustacea

L. callinectes Couch,1942@ many crustaceans, including e.g. *Homarus gammarus*

North Carolina

L. ethamalophilum T.W. Johnson,1958

@ barnacle eggs

E USA

Myzocytium Schenk,18??*M. vermicola* (Zopf) Fischer, em. Newell, Cefalu & Fell,1977

@ parasiting & killing littoral nematodes

?

M. proliferum Schenk, forma *marinum* Kobayashi & Ookubo,1953

Japan?

Olpidiopsidaceae*Olpidiopsis* Cornu,1872*O. sphacelarum* (Kny,1???)

@ macro algae

W Sweden

O. andreei (Lagerheim) Karling,1942= *Pleotrichelus andreei* Lagerheim,1899= *Biciliatum andreei* (Lagerheim) Petersen,1905= *Pleotrichelus ectocarpii* Jökl,1916= *Petersenia andreei* (Lagerheim) Sparrow,1936@ macro algae, e.g. *Spongomorpha*, *Acrosiphonia*, *Ectocarpus* & *Ceramium*

W Sweden, Denmark, North Sea, Greenland, Arctic Seas, France, Adriatic, USA

O. feldmanni Aleem,1952@ macro algae, e.g. *Falkenbergia rufolanosa* & *Trailliella intricata*

W Sweden, France

O. antithamnionis Whittick & South,1972@ *Antithamnion floccosum* (O.F. Müller) Kleen

Newfoundland

O. globosa Anastasiou & Churchland,19??@ the deutericymete *Papulaspora halima* Anastasiou

?

O. paradoxus ??,1???

@ fungi

?

O. magnus Feldmann & Feldmann,1955@ rhizoids of *Ceramium flabelligerum*

France

Sirolpidiaceae*Lagenisma* Drebes,19??*L. coscinodisci* Drebes,1968

@ endoparasite in *Coscinodiscus* spp. & *Palmeria hardmaniana*
W Sweden, North Sea, Puget Sound, Jamaica

Petersenia Sparrow,1934
* Familial position uncertain

P. lobata (Petersen) Sparrow,1934
= *Pleotrichelus lobatus* Petersen,1905
@ red algae, often *Ceramium*
W Sweden, Denmark, Ireland, France, USA

P. pollagaster (Petersen) Sparrow,1934
= *Pleotrichelus pollagaster* Petersen,1905
@ macro algae, e.g. *Chondrus crispus*
W Sweden, Nova Scotia

P. palmariae van der Meer & Pueschel,1985
@ *Palmaria mollis*
? Canada

Pontisma Petersen,1905

P. lagenidoides Petersen,1905
= *Sirolipidium lagenidoides* (Petersen) Karling,1942
@ macro algae, especially *Ceramium* spp.
W Sweden, Denmark, North Sea, Germany, France, USA, Japan

Sirolipidium Petersen,1905
@ including Crustacean, oyster & clam larvae parasites

S. bryopsisidis (de Bruyne) Petersen,1905
= *Olpidium bryopsisidis* de Bruyne,1890
@ macro algae, e.g. *Bryopsis plumosa* & *Cladophora* spp.
W Sweden, Denmark, North Sea, France, Italy, USA, Japan

S. zoophithorum Vishniac,1955
@ larvae of *Crassostrea virginica* & *Venus* spp.
USA

Synchaetophagus Apstein,1911
* Family & order very provisional.

S. balticus Apstein,1911
@ the rotifer *Synchaeta monopus*
Baltic

Ostracoblabe
* Family & order very provisional. Should possibly be assigned to Deuteromycotina.

O. implexa Bornet & Flahault,19??
@ *Ostrea edulis*
?

Halipitoraceae

Haliphthoros Vishniac,1958

H. milfordensis Vishniac,1958
@ eggs & juveniles of Crustacea, e.g. *Homarus americanus* & eggs of *Urosalpinx*; also observed on
Enteromorpha sp.
USA

PERONOSPORIDA
= **PERONOSPORALES**

Phytiaeae

Phytophtora de Bary,1???

P. pythium Pringsheim,1858

P. marinum Sparrow,1934
@ *Porphyra perforata* & *Ceramium rubrum*
Denmark

P. maritimum Höhnk,1939
@ *Ceramium* sp.
Germany

P. imperfectum Höhnk,1953
Germany

P. aquatilis Höhnk,1953
Germany

P. porphyrae Takahashi, Ichinotani & Sasaki,19??
@ *Porphyra* spp.
* may be a synonym of *P. marinum* Sparrow
?

P. salinum Höhnk,1953
Germany

P. graminicolum var. *stagni* Höhnk,1953
* dubious taxon
?

P. undulatum var. *littorale* Höhnk,1953
@ *Fucus distichus*
British Columbia

LEPTOMITIDA

= **LEPTOMITALES**
* Exclusively freshwater, according to Moss,1991

Leptomitaceae

Leptomitus C.A. Agardh,18??
* Genus in need of revision

L. lanugo C.A. Agardh,18??
@ *Ceramium rubrum*
?

L. minutissimus (Dillwyn) C.A. Agardh,18??
@ filamentous marine green algae
?

HYPHOCHYTRIDIOMYCOTEA

HYPHOCHYTRIDA
= **HYPHOCHYTRIALES** = **ANISOCHYTRIALES**
* 6 genera known

Elinia ??,1???

* Which family?

E. marisalba

* not parasite of algae or animals
?

E. sinorifica

* not parasite of algae or animals
?

Anisopidiaceae

Anisopodium Karling,1943

A. sphacellarium (Kny) Karling,1943
= *Chytridium (Olpidium) sphacellarium* Kny,1871
= *Pleotrichelus sphacellarium* (Kny) Petersen,1905
@ *Sphacellaria*, *Cladostephus* & *Chaetopteris*
W Sweden, Helgoland, Britain, Denmark, Ireland, France, Italy, USA, Japan

A. ectocarpii Karling,1943

@ *Ectocarpus* spp.
North Carolina

A. rosenvingii (Petersen) Karling,1943

= *Pleotrichelus rosenvingii* Petersen,1905
@ *Pylaiella littoralis* (Linnaeus) Kjellman
Sweden, Denmark

Hypochytriaceae

Hypochytrium Zopf,18??

H. peniliae Artemchuk & Zelzinskaia,1969

@ *Penila avirostris* (Dana) & *Eurytemora*

H. oceanicum ??,1???

* not parasite of algae or animals
?

Rhizidiomyctaceae

Latrostium Zopf,18??

L. comprimens ??,1???

@ eggs of marine? *Vaucheria*

Rhizidiomyces Zopf,1884

R. bulbosus Karling,1968
@ ?
Helgoland, Oceania

LABYRINTHOMORPHA Page,1980

* Belongs in the informal group Straminopiles

Diplophryidae

* May belong in Labyrinthulea or Thraustochytrida

Diplophrys Barker,1868

D. archeri Barker,1868
= *D. marina* Dykstra & Porter,1984
Tåmminne, Arctic Canada, E & W USA, European fresh waters

Sorodiplophrys Dykstra & Olive,1975

* marine?

S. stercorea ?Dyksra,1976?

LABYRINTHULEA Levine & Corliss,1963

LABYRINTHULIDA Lankester,1877

Labryinthulidae Haeckel,1868

* monogenetic family

Labyrinthula Cienkowsky,1867 (8)

= *Chlamydonyxa* Archer,1875
= *Labryinthomyxa* Duboscq,1921 (TM) *L. sauvageau* Duboscq,1921 - in *Laminaria lejolisi* Sauvageau
= *Pseudoplasmodium* Molisch,1926
= *Labryinthodictyon* Valkanov,1972

L. vitellina Cienkowsky,1867

Sweden, Black Sea

L. macrocystis Cienkowsky,1867

@ *Zosteraceae* & marine algae

Sweden, Black Sea, Europe

L. roscoffensis Chadefaud,1956

@ marine algae

Roscoff

L. valkanovii Karling,1944

@ algae

Black Sea

L. coenocystis Schmoller,1960

?

L. magnifica (Valkanov,1972)

= *Labryinthodictyon magnificum* Valkanov,1972

@ diatoms

Black Sea

L. algeriensis Hollande & Enjumet,1955

@ *Spartina alterniflora*

?

L. aurantiacum (Molisch,1926)

= *Pseudoplasmodium aurantiacum* Molisch,1926

Japan

THRAUSTOCYTRIDEA

Thraustochytriidae Sparrow,1943, 1960

Thraustochytrium Sparrow,1936, emend. Johnson, in Johnson & Sparrow,1961 (16)

@ one species is parasitizing *Labyrinthula* sp.

- T. benthicola** Raghu-Kumar,197?
- T. multirudimentale** Goldstein,1963
off NW Africa or off Portugal
- T. aureum** Goldstein,1963
North Sea
- T. aggregatum** Ulken,1965
Weser estuary
- T. striatum** Schneider,1967
Kieler Bucht, Britain (Esk estuary), New Zealand
- T. kinnei** Gaertner,1967
Helgoland
- T. motivum** Goldstein,1963
E Scotland at 80 m
- T. pachydermum** Scholz,1958
North Sea, W Mediterranean, USA
- T. roseum** Goldstein,1963
US Atlantic, New Zealand
- T. proliferum** Sparrow,1936 (TM)
@ Bryopsis plumosa & Ceramium diaphanum
Germany, USA
- T. inglei** Quick,1972
@ thought to cause early summer mortalities in *Crassostrea virginica*
Florida
- T. globosum** Kobayasi & Ookubo,1953
@ Ulothrichales
Japan
- T. arudimentale** Artemchuk,19??
?
- T. rossii** Bahnweg & Sparrow,1974?
Antarctic
- T. antarcticum** Bahnweg & Sparrow,1974?
Antarctic
- Ulkenia** Gaertner,1977 (6)
- U. amoeboidae** (Bahnweg & Sparrow,1974)
* Is Gaertner the author?
@ Eledone cirrhosa
?
- U. minuta** Raghu-Kumar,1977
North Sea
- U. profunda** Gaertner,1977
off Portugal at 1883m
- U. radiata** Gaertner,1977
Helgoland rocky shore
- U. visurgensis** (Ulken,1965) Gaertner,1977 (TM)
= Thraustochytrium visurgense Ulken,1965
Weser
- U. sarkariana** Gaertner,1977
Weser estuary
- Schizochytrium** Goldstein & Belsky,1964 (2)
@ one species is parasitizing *Thalassionema nitzschiooides* in S Norway
- S. minutum** Gaertner,1981
North Sea, Barents Sea, River Tay, Portugal, W Africa, Madeira
- S. aggregatum** Goldstein & Belsky,1964 (TM)
Thames estuary, Florida
- Labyrinthuloides** Perkins,1973 (4)
* May be congeneric with *Aplanochytrium* Bahnweg & Sparrow,1972
- L. yorkensis** Perkins,1973 (TM)
North Sea, Virginia
- L. minuta** (Watson & Raper,1957)
= *Labyrinthula minuta* Watson & Raper,1957
@ marine algae
USA
- L. saliens** Quick,1974
@ *Halophila engelmanni*, but also isolated from *Crassostrea virginica*
USA
- L. schizochytrops** Quick,1974
- Japanochytrium** Kobayasi & Ookubo,1953 (1)
- J. marinum** Kobayasi & Ookubo,1953 (TM)
@ *Gracilaria confervoides* & *Cladophora japonica*
Japan
- Aplanochytrium** Bahnweg & Sparrow,1972 (1)
- A. kerguelensis** Bahnweg & Sparrow,1972 (TM)
?Sargasso Sea, Madeira, Kerguelen
- Althornia** Gareth-Jones & Alderman,1971 (1)
- A. crouchii** Gareth-Jones & Alderman,1971 (TM)
- Corallochytrium**
- APICOMPLEXA** Levine,1970
- PERKINSASIDA** Levine,1978
- PERKINSORIDA** Levine,1978
- Perkinsidae** Levine,1978
- Perkinsus** Levine,1978
- P. marinus** (Mackin, Owen & Collier,1950) Levine,1978 (TM - Gulf of Mexico)
= *Dinomyscystidium* Perez,1908 *marinum* Mackin, Owen & Collier,1950
= *Labyrinthomyxa marina* (Mackin, Owen & Collier,1950) Mackin & Ray,1966
= *Labyrinthula marina* : Auctt.
@ causes (mortal) tissue lysis in *Crassostrea virginica* (Gmelin)
Gulf of Mexico- Connecticut
- P.** sp.
@ *Macoma balthica*
Maryland
- SPOROZOASIDA** Leuckart,1879
- GREGARINASINA** Dufour,1828
- ARCHIGREGARINORIDA** Grassé,1953
- Selenidioididae** Levine,1971
@ 3 genera & 13 sp. in intestines of marine polychaetes, sipunculans, ascidians & enteropneusts
- Selenidioides** Levine,1971 (TM) *S. caulleryi* (Brasil,1907) Levine,1971
- S. hollandei** (Vivier & Schrével,1966) Levine,1971
@ posterior intestine of *Sabellaria alveolata*
- Meroselenidium** Mackinnon & Ray,1933 (TM) *M. keilini* Mackinnon & Ray,1933
- Merogregarina** Porter,1908 (TM) *M. amarouci* Porter,1908
- Exoschizontidae** Levine,1971
- Exoschizion** Hukui,1939
- E. siphonosomae** Hukui,1939 (TM)
@ *Siphonosoma kumanense*
* The only known species of the family.
- EUGREGARINORIDA** Léger,1900
- BLASTOGREGARINORINA** Chatton & Villeneuve,1936 emend.
- Siedleckiidae** Chatton & Villeneuve,1936
- Siedleckia** Caullery & Mesnil,1898
* 4 named species in marine pylychaetes
- S. nematoidea** Caullery & Mesnil,1898 (TM)
@ *Phylo norvegica*, *Scoloplos armiger* & *Aricia latreilli*
Trondheim, Roscoff, Anse St. Martin
- S. mesnili** Chatton & Dehorne,1929
@ *Aricia latreilli*
Roscoff
- S. dogieli** Chatton & Dehorne,1929
@ *Aricia foetida*
Bay of Naples
- S. caulleryi** Chatton & Villeneuve,1936
@ *Protoarcia oerstedi*
?
- ASEPTATORINA** Chakravarty,1960
- Selenidiidae** Brasil,1907
@ 4 genera & 45 sp. in marine invertebrates
- Selenidium** Giard,1884
= *Platyctysis* Léger,1892
* 41 named sp.
- S. pendula** Giard,1884 (TM)
@ intestine of *Nerine cirratulus* (*Scolelepis squamata*)
Engl. Channel
- S. cirratali** (Mingazini,1893)
@ *Audouinia tentaculata*
Engl. Channel
- S. branchiomatis** Ray,1930
@ *Branchiomma vesiculosum*
Plymouth
- S. allenii** Ray,1930
@ *Branchionoma vesiculosum*
Plymouth
- S. plicatum** Ray,1930
@ *Cirratulus cirratus*
Plymouth
- S. virgula** Caullery & Mesnil,1919
@ *Cirratulus cirratus*
Engl. Channel
- S. echinatum** Caullery & Mesnil,1899
@ *Dodecaceria concharium*
Engl. Channel
- S. mesnili** Brasil,1909
@ *Myxocola infundibulum*
Plymouth, Roscoff
- S. costatum** Siedlecki,1903
@ *Nicolea venustula* & *Eupolympnia nebulosa*
Engl. Channel, Cannes, Gulf of Naples
- S. brasili** Ray,1930
@ *Pomatoceros triqueter*
Plymouth, Wimereux
- S. caulleryi** Brasil,1907
@ *Protula tubularia*
Calvados, Plymouth
- S. serpulae** Lankester,1863
@ *Serpula vermicularis*
Plymouth, Gulf of Naples

| | |
|--|---|
| <i>S. terebellae</i> (Kölliker,1845) @ <i>Terebella lapidaria</i> Plymouth | <i>L.</i> sp. @ <i>Eupolynnia nebulosa</i> W France |
| <i>S. sabellae</i> (Lankester,1863) @ <i>Sabella pavonina</i> Plymouth | <i>L.</i> sp. : (de Beauchamp,1913) @ <i>Ophelia neglecta</i> Roscoff |
| <i>S. spinos</i> (Kölliker,1845) @ <i>Malacoceros fuliginosa</i> Helgoland, Engl. Channel | <i>L.</i> sp. @ <i>Petaloprocetus terricola</i> W France |
| <i>S. foliatum</i> Ray,1930 @ <i>Malacoceros fuliginosa</i> Helgoland, Engl. Channel | ? <i>L.</i> sp. @ <i>Temora</i> sp. Baltic, Kattegatt, North Sea |
| <i>S. intraepitheliale</i> Reichenow,1932 @ <i>Malacoceros fuliginosa</i> Helgoland, Engl. Channel | <i>Polyrhadina</i> Mingazzini,1891 @ 4 named sp. in polychaetes |
| <i>S. metchnikovi</i> Léger & Duboscq,1917 @ <i>Glossobalanus minutus</i> Biscay | <i>P. spinos</i> (von Kölliker,1848) Mingazzini,1891 (TM) @ <i>Malacoceros fuliginosa</i> Helgoland, Engl. Channel |
| <i>S. sipunculi</i> (Dogiel,1907) @ <i>Sipunculus mudis</i> Gulf of Naples | <i>P. polydorae</i> (Léger,1893) @ <i>Polydora</i> sp., ? <i>Polydora flava</i> Helgoland, Plymouth |
| <i>S. ceruzi</i> Faria, Cunha & Fonseca,19?? @ <i>Polydora</i> sp. Helgoland | <i>P. pygospionis</i> Caullery & Mesnil,1914 @ <i>Pygospio elegans</i> Engl. Channel |
| <i>S. sp.</i> @ <i>Scoloplos armiger</i> Engl. Channel | <i>P. bifurcata</i> Mackinnon & Ray,1931 @ <i>Malacoceros fuliginosa</i> Plymouth |
| <i>S. sp.</i> @ <i>Spio filicornis</i> Engl. Channel | <i>P. brasili</i> Caullery & Mesnil,1914 @ <i>Spio filicornis</i> Engl. Channel |
| <i>S. sp.</i> @ <i>Pygospio elegans</i> Engl. Channel | <i>Ulivina</i> Mingazzini,1891 @ 6 named sp. in polychaetes |
| <i>S. sp.</i> @ <i>Salmacina dysteri</i> Engl. Channel | <i>U. elliptica</i> Mingazzini,1891 (TM) @ <i>Audouinella filigera</i> , <i>A. tentaculata</i> , <i>Nicolea venustula</i> , <i>Petaloprocetus terricola</i> & <i>Eupolynnia nebulosa</i> Engl. Channel, W France, Cannes |
| <i>S. sp.</i> @ <i>Ophelia limacina</i> Murmansk | <i>Syctia</i> Léger,1892 (TM <i>S. spinipinata</i> Léger,1892) @ 4 named sp. in polychaetes |
| <i>S. sp.</i> @ <i>Polydora coeca</i> Engl. Channel | <i>Pontesia</i> Hasselmann,1926 (TM <i>P. kelschi</i> Hasselmann,1926) @ one named sp. in intestines of marine annelids |
| <i>S. sp.</i> @ <i>Ctenodrilus serratus</i> Engl. Channel | <i>Bhatiella</i> Setna,1931 (TM <i>B. morphysae</i> Setna,1931, emend. Levine,1977) = <i>Ferraria</i> Setna,1931 = <i>Coryloepimeritus</i> Hoshida (Tugawa),1944 @ 2 named sp. in polychaetes |
| <i>S. sp.</i> @ <i>Nephasoma minuta</i> Wimereux | <i>Viviera</i> Schrével,1963 (TM <i>V. morphysae</i> Schrével,1963) @ one named sp. in intestines of polychaetes |
| <i>S. sp.</i> @ <i>Cirratulus cirratulus</i> Engl. Channel | <i>Cochleomeritus</i> Hoshida (Tugawa),1944, emend. (TM <i>C. lysidici</i> Hoshida (Tugawa),1944) @ 2 named sp. in polychaetes |
| <i>S. sp.</i> : Brasil & Fantham,1907 @ <i>Phascolosoma elongatum</i> Engl. Channel | <i>A. sagittata</i> (Leuckart,1861) Labbé,1899 @ <i>Capitella capitata</i> Helgoland, Engl. Channel, Gulf of Naples |
| <i>S. sp.</i> : Brasil & Fantham,1907 @ <i>Phascolosoma vulgare</i> Engl. Channel | <i>A. prolifera</i> Clausen,1993 @ <i>Microphthalminus ephippiophorus</i> Clausen (Polychaeta, Hesionidae) Bergen, Trondheim |
| <i>Selenocystis</i> Dibb,1938 (TM <i>S. foliata</i> (Ray,1930) Dibb,1938) * one named sp. | <i>A. lutzi</i> Hasselmann,1918 * ?dubious species ? |
| <i>Ditrypanocystis</i> Burt, Denny & Thomasson,1963 (TM <i>D. cirratuli</i> Burt, Denny & Thomasson,1963) * 2 named sp. | <i>A. criodrilli</i> Sciacchitano,1931 @ <i>Criodrilus lacuum</i> ? |
| <i>Heterospora</i> de Saint Joseph,1907 * one named sp. | <i>A.</i> sp. : Caullery & Mesnil,1919 @ <i>Capitellides giardi</i> Engl. Channel |
| <i>H. eulaliae</i> de Saint Joseph,1907 (TM) @ coelom of <i>Eudalia</i> | <i>Hentschelia</i> Mackinnon & Ray,1931 @ one named sp. in intestines of echiuroids |
| <i>Lecudinidae</i> Kamm,1922 @ 26 genera & 138 sp. almost all in marine invertebrates | <i>H. thalassemae</i> Mackinnon & Ray,1931 (TM) @ <i>Thalassema neptuni</i> & <i>Phascolion strombi</i> Plymouth |
| <i>Lecudina</i> Mingazzini,1891 = <i>Dolycystis</i> Léger,1893 = <i>Koelikeria</i> Mingazzini,1891 = <i>Koelikerella</i> Labbé,1899 = <i>Cygnicollum</i> Bogolepova,1953 @ 42 sp. in polychaetes & other marine invertebrates | <i>Lecythion</i> Mackinnon & Ray,1931 @ one named sp. in intestines of echiuroids |
| <i>L. pellucida</i> (von Kölliker,1848) Mingazzini,1891 (TM) @ <i>Perinereis cultrifera</i> , <i>Platynereis dumerillii</i> & <i>Nereis</i> ? <i>pelagica</i> Helgoland, Mediterranean | <i>L. thalassemae</i> Mackinnon & Ray,1931 (TM) @ <i>Thalassema neptuni</i> & <i>Phascolion strombi</i> Plymouth |
| <i>L. tuzae</i> Schrével,1963 @ <i>Hediste diversicolor</i> ? | <i>Hyperidion</i> Mackinnon & Ray,1931 @ one named sp. in intestines of echiuroids |
| <i>L. elongata</i> (Mingazzini,1893) @ <i>Lumbrineris impatiens</i> , <i>Lumbrineris latreilli</i> Engl. Channel, Mediterranean | <i>H. thalassemae</i> Mackinnon & Ray,1931 (TM) @ <i>Thalassema neptuni</i> Plymouth |
| <i>L. legeri</i> (Brasil,1909) @ <i>Glycera convoluta</i> Engl. Channel | <i>Zygosoma</i> Labbé,1899 = <i>Conorhynchus</i> Greff,1880, <i>non</i> Motschoulsky,1860 (Coleoptera), <i>nec</i> Gill, <i>ex</i> Nozemann,1861 (Pisces), <i>nec</i> Bleeker,1863 (Pisces) @ 2 named sp. in intestines of echiuroids |
| <i>L. franciana</i> ?Arvy,1952? @ intestine of <i>Phascolion strombi</i> Dinard | <i>Z. gibbosum</i> (Greeff,1880) Labbé,1899 = <i>Conorhynchus gibbosus</i> Greeff,1880 (TM - Norway) @ <i>Echiurus pallasi</i> ?= <i>E. echinus</i> Norway |
| <i>L. lankesteri</i> (Desportes & Théodoridès,1986) = <i>Cygnicollum lankesteri</i> Desportes & Théodoridès,1986 @ <i>Laetmonice producta</i> & <i>L. hystrix</i> ? | <i>Filipodium</i> Hukui,1939 (TM <i>F. ozakai</i> Hukui,1939) @ 2 named sp. in intestines of sipunculids |
| | <i>Diplauxis</i> Vivier, Ormières & Tuzet,1964 (TM <i>D. hatti</i> Vivier, Ormières & Tuzet,1964) @ 2 named sp. in polychaetes |

- Chlamydocystis*** Dogel,1910 (TM *C. captiva* Dogel,1910)
@ one named sp. in polychaetes
- Contortiocorda*** Bhatia & Setna,1938 (TM *C. prashadi* Bhatia & Setna,1938)
* one named sp.
- Lankesteria*** Mingazzini,1891
= *Pleurozyga* Mingazzini,1891
@ 26 named sp. in ascidiants
- L. ascidiae*** (Lankester,1872) Mingazzini,1891
= *Gregaria (Monocystis) ascidiae* Lancester,1827 (TM)
@ *Ciona intestinalis*, *Dendrodoa grossularia*, *Polycarpa pomaria*, *Styela rustica* etc.
Roscoff
- L. amarouci*** (Giard,1873)
@ *Amaroucium nordmanni*, *A. punctatum*, *Aplidium zostericola*, *Leptoclinum listerianum*, *Morcheilium argus*,
Polysyncraton lacazei & *Sidnyum turbinatum*
Roscoff
- L. ascidiella*** Duboscq & Harant,1923
@ *Ascidella aspersa*
Roscoff
- L. acutissima*** ?,1???
- L. siedleckii*** Duboscq & Harant,1923
@ *Ascidia mentula* & *A. conchilega*
- L. parascidiae*** Duboscq & Harant,1923
@ *Parascidia elegans* = *Sydnium elegans*
Roscoff
- L. botrylli*** ?,1???
- L. maculata*** ?,1???
- L. striata*** ?,1???
- L. clavellinae*** ?,1???
- L. zonata*** ?,1???
- L. tethyi*** ?,1???
- L. diazona*** ?,1???
- L. monstrosa*** ?,1???
- L. distapliae*** ?,1???
- Monocystella*** Valkanov,1935 (TM *Monocystella arndti* (Valkanov,1931) Valkanov,1935)
= *Pseudolankesteria* Ormières,1965
@ 10 named sp. in turbellarians
- M. cyclopori*** (Poisson,1921)
= *Lankesteria cyclopori* Poisson,1921
@ *Cycloporus maculatus*
* Correct genus?
Engl. Channel
- Ophioidina*** Mingazzini,1891
@ one named sp. in echinoids
- O. bonelliae*** (Frenzel,1885) Labbé,1899
= *Gregaria bonelliae* Frenzel,1885 (TM - Mediterranean)
@ *Bonellia viridis*
Mediterranean
- Paraophioidina*** Levine,1977 (TM *Paraophioidina haekeli* (Mingazzini,1891) Levine,1977)
@ 11 named species in crustaceans
- Lateroprotomeritus*** Théodoridès & Desportes,1975 (TM *L. conicus* Théodoridès & Desportes,1975)
* one named sp.
- Extremocystis*** Setna,1931 (TM *E. dendrostomi* Setna,1931)
@ one named sp. in sipunculids
- Lobianchella*** Mingazzini,1891 (TM *L. belenoides* Mingazzini,1891)
@ one named sp. in polychaetes
- Urosporidae*** Léger,1892
@ 6 genera & 57 named species, mostly in the coelom of polychaetes, oligochaetes, sipunculids, nemertans, molluscs & echinoderms
* An undescribed species in *Antalis entale* from N Bretagne is thought to be related to *Gonospora testiculi* Tregouboff,1918 from *Cerithium vulgatum* as well as to another undescribed gregarine infesting *Turricula communis*
- Urospora*** Schneider,1875
= *Cystobia* Mingazzini,1891 (p.p.)
= *Pachysoma* Mingazzini,1891 = *Syncystis* Cuénnot,1891
@ 21 named species in annelids, sipunculids, nemertines & echinoderms
- U. nemertis*** (von Kölleker,1845) Schneider,1875 (TM)
* Is the spelling possibly *U. nemertes* ?
@ *Valencinia*
Engl. Channel
- U. echinocardii*** Pixell-Goodrich,1915
@ coelomic cavity of *Echinocardium cordatum* & *Spatangus purpureus*
Plymouth
- U. synaptae*** (Cuénnot,1891)
?= *Goniosporea mercieri* Cuénnot,1912
@ gut-associated hemal system & coelomic cavity of e.g. *Leptosynapta inhaerens*
Engl. Channel
- U. lagidis*** de St.-Joseph,1898
@ *Lagis koreni*
Calvados, Irish Sea
- U. chiridotae*** (Dogiel,1906)
= *Cystobia chiridotae* Dogiel,1906
@ gut-associated hemal system of *Chiridota laevis*
Murmansk, W Canada
- U. sipunculi*** (von Kölleker,1845) Léger,1892
@ *Sipunculus nudus*
W France, Mediterranean
- U. ovalis*** Dogiel,1910
@ *Travisia forbesi*
Bergen, Murmansk
- U. travisiæ*** Dogiel,1910
@ *Travisia forbesi*
Bergen, Murmansk
- U. neapolitana*** Pixell-Goodrich,1915
@ coelomic cavity of *Echinocardium cordatum*
Engl. Channel & Mediterranean
- Gonospora*** A. Schneider,1875
= *Cystobia* Mingazzini (p.p.)
= *Diplodina* Woodcock,1906 = *Kalpidorhynchus Cunningham*,1907
@ 17 named sp. in coelom or rarely intestines of polychaetes, testes of prosobranchs, coelom, respiratory tree & blood vessels of holothurians
- G. longissima*** Caullery & Mesnil,1898
@ *Dodecaceria concharum*
Engl. Channel
- G. arenicola*** (Cunningham,1907)
@ *Arenicola ecaudata*
Plymouth
- G. varia*** (Léger,1892) (TM)
@ *Audouinia tentaculata*
Engl. Channel
- G. irregularis*** (Minchin,1893)
@ *Holothuria forskaali*
Plymouth
- G. stichopi*** (Lützen,1968)
= *Cystobia stichopi* Lützen,1968
* Correct genus?
@ dorsal radial hemal system of *Parastichopus tremulus*
Gullmarsfjord, Oslofjord, Trondheimfjord
- G. gonadipertha*** (Djakonov,1923)
= *Diplodina gonadipertha* Djakonov,1923
@ on and in gonads of *Cucumaria frondosa*
Barents Sea
- G. minchinii*** Goodrich & Pixell Goodrich,1920
@ eggs & coelomic fluid of *Arenicola ecaudata*
Plymouth
- Lithocystis*** Giard,1876, em. Pixell-Goodrich,1916
@ 8 named sp. in sipunculids & echinoderms
- L. cucumariae*** Pixell-Goodrich,1929
@ respiratory trees of *Pawsonia saxicola*
Plymouth
- L. minchini*** (Woodcock,1904)
@ coelomic side of body wall of *Pawsonia saxicola*
Plymouth
- L. foliacea*** Pixell-Goodrich,1915
@ coelomic cavity of *Echinocardium cordatum*
Plymouth, Mediterranean
- L. schneideri*** Giard,1876 (TM)
@ coelomic cavity of *Echinocardium cordatum*, *Spatangus purpureus* & *Echinocardium* sp.
Engl. Channel, Mediterranean
- L. microspora*** Pixell-Goodrich,1915
@ *Spatangus purpureus*
Engl. Channel
- L. brachycercus*** Pixell-Goodrich,1925
@ *Chiridota laevis*
W Canada
- Ceratospora*** Léger,1892 (TM *C. mirabilis* Léger,1892)
@ one named sp. in polychaetes
- Pterospora*** Labbé & Racovitza,1897
@ 3 named sp. mostly in coelom of polychaetes
- P. maladenerum*** Labbé & Racovitza,1897 (TM)
@ *Leiochone liopygus* & *Petaloprocus terricola*
Roscoff
- P. ramificata*** Dogiel,1910
@ *Nicomedie lumbicalis*
Murmansk
- P. clymenellae*** (Porter,1897)
@ *Clymenella torquata*
?
- Paragonospora*** Lang,1954 (TM *P. typica* Lang,1954)
@ 1 named sp. in polychaetes
- Aikinetocystidae*** Bhatia,1930
@ 2 genera in coelom of oligochaetes
- Monocystidae*** Bütschli,1882
@ 27 genera & 188 named species, the great majority in oligochaetes
- Monocystinae*** Bhatia,1930

- @ 7 genera in seminal vesicles (or coelom) of oligochaetes
- Zygocystinae Bhatia,1930
@ 3 genera in oligochaetes
- Rhynchocystinae Bhatia,1930
@ 3 genera in oligochaetes
- Stomatophorinae Bhatia,1930
@ 10 genera in oligochaetes (or millipedes)
- Oligochaetocystinae Meier,1956 emend. Levine,1977
@ 4 named genera in oligochaetes, echiurids, chilopods & mites
- Oligochaetocystis Meier,1956 (TM *Oligochatocystis pachydrili* (Lankester,1863) Meier,1956, emend. Levine,1977)
* 3 named sp.
- Echiurocystis Levine,1977 (TM *Echiurocystis yumushii* (Iitsuka,1933) Levine,1977)
@ 3 named sp. in echiurids
- Diplocystidae Bhatia,1930
@ insects
- Allantocystidae Bathia,1930
@ ceratopogonid larvae
- Schaudinellidae Poche,1913
@ earthworms
- Gamynedidae Huxley,1910
@ one sp. known from *Anaspides tasmaniae*
- Enterocystidae Codreanu,1940
@ insects
- SEPTATORINA Lankester,1885
- Porosporoidea Chakravarty,1960
- Porosporidae Labb  ,1899
@ 3 genera & 31 sp. with vegetative & sexual development in decapod crustaceans and sporogony in connective tissue of Bivalvia
- Nematopsis Schneider,1892
= *Carcinoecetes* Ball,1938 (p.p.)
* 24 named sp.
- N. pectinis (L  ger & Duboscq,1925)
@ *Chlamys varia* ; decapod host unknown
Roscoff
- N. veneris (L  ger & Duboscq,1925)
@ gills of *Clausinella fasciata* ; decapod host unknown
Mediterranean
- N. legeri de Beauchamp,1910
= *N. mediterranea* L  ger,1905 (nomen oblitum)
= *Porospora galloprovincialis* L  ger & Duboscq,1925
@ oocysts in e.g. *Mytilus galloprovincialis*, *Lasaea rubra*, other bivalves & prosobranchs; definite host is the crab *Eriphia spinifrons*
French Atlantic, Mediterranean
- N. portunidarum (Frenzel,1885) Hatt,1931 (TM)
@ oocysts in gill vessels of *Cerastoderma edule* ; vegetative & sexual stages in *Liocarcinus depurator*, *Liocarcinus arcuatus* & ?*Carcinus maenas*
Europe
- N. schneideri ??,??
@ oocysts in *Mytilus edulis*, *Spirula solidula*, *Donax vittatus*, *Venerupis pullastra*, *Macoma balthica*, *Cerastoderma edule* & other bivalves; vegetative stage in *Carcinus maenas* & *Liocarcinus depurator*
Europe
- N. maraisi (L  ger & Duboscq,1911)
@ vegetative stage in *Liocarcinus depurator*
Mediterranean
- N. ostrearium ?,??
@ *Crassostrea virginica* & crabs
USA
- N. pryeri ?,??
@ *Crassostrea virginica* & crabs
USA
- Porospora Schneider,1875* 4 named sp.
- P. gigantea (van Beneden,1869) Schneider,1875 (TM)
@ various gastropods, e.g. *Trochocochea mutabilis* ; final host: *Homarus gammarus*
Engl. Channel, W France, Mediterranean
- P. nephropis L  ger & Duboscq,1911
@ alimentary channel of *Nephrops norvegicus*
W France
- Pachyoporospora Th  odorid  s,1961 (TM *P. laubieri* Th  odorid  s,1961)
* 3 named sp.
- Gregarinoidea Chakravarty,1960
- Cephaloidophoridae Kamm,1922
@ 3 genera & 61 sp.
- Cephaloidophora Mavrodiadi,1908
= *Carcinoecetes* Ball,1938 (p.p.)
@ 56 named sp. in intestines of amphipods, barnacles, crabs & shrimps
- C. duboscqi Poisson,1924
@ *Balanus nitescens*
Engl. Channel
- C. communis Mavrodiavi,1908
@ *Balanus improvisus*
Baltic
- C. ocellata (L  ger & Duboscq,1908)
@ *Pagurus prideauxii*
Banyuls-sur-mer
- C. maculata L  ger & Duboscq,1911
@ *Balanus duebeni*, *G. locusta* & *G. marinus*
Engl. Channel
- C. brasili Poisson,1920
@ *Orchestia littorea*
Engl. Channel
- C. talitri Mercier,1912
@ *Talitrus saltator*
Roscoff
- C. fossor (L  ger & Duboscq,1908)
@ *Pinnotheres pisum*
Engl. Channel
- C. conformis (Diesing,1851)
@ *Pachygrapsus marmoratus*
Mediterranean
- ?C., sp.
@ *Mysis mixta*
Baltic, Belt Sea
- Caridohabitans Ball,1959 (TM *C. setnai* Ball,1959)
@ 1 named sp. in digestive tract of Crustacea
- Rotundula Goodrich,1949 (TM *Rotundula maculata* (L  ger & Duboscq,1911) Goodrich,1949)
@ 4 named sp. in amphipods
- Cephalobidae Th  odorid  s & Desportes,1975
@ 2 genera & 4 named sp. in marine crustacea
- Cephalolobus Kruse,1959 (TM *C. penaeus* Kruse,1959)
@ 3 named sp. in guts of marine Crustacea
- Callynchrochlamys Frenzel,1885 (TM *C. phronimae* Frenzel,1885)
@ 1 named sp. in intestines of amphipods
- Uradiphoridae Grass  ,1953
@ 5 genera & 19 named sp. in intestine of Crustacea
- Uradiphora Mercier,1912 (TM *Uradiphora cuenoti* (Mercier,1911) Mercier,1912)
@ 5 named sp. in decapods & amphipods
- U. athanasi Poisson,1924
@ *Athanas nitescens*
Engl. Channel
- U. gammari Poisson,1924
@ *Gammarus locusta* & *Gammarus oceanicus*
Engl. Channel
- U. mercieri Poisson,1921
@ *Orchestia littorea*
Engl. Channel
- Heliospora Goodrich,1949 (TM *Heliospora longissima* (von Siebold, in von K  lliker,1848) Goodrich,1949)
@ 3 named sp. in amphipods
- Pyxinoides Tr  gouboff,1912
@ 10 named sp. in cirripedes
- P. balani (von K  lliker,1848) Tr  gouboff,1912 (TM)
@ *Balanus amphitrite* & *B. eburneus*
Mediterranean
- P. chthamali (L  ger & Duboscq,1909)
@ *Chthamalus stellatus*
Mediterranean
- Nematoidea Mingazzini,1891 (TM *N. fusiformis* Mingazzini,1891)
@ one named sp. in cirripedes
- Bifilida Tuzet & Ormi  res,1964 (TM *B. rara* Tuzet & Ormi  res,1964)
@ one named sp. in cirripedes
- Gregarinidae Labb  ,1899
@ typically in intestines of insects
- Degiustia Levine,1980 (TM *Degiustia hyalella* (Batten & DeGiusti,1949) Levine,1980)
@ one named sp. in amphipods
- Cirrigregarina Levine,1980 (TM *Cirrigregarina spissa* (Henry,1938) Levine,1980)
@ 2 named sp. in cirripeds
- Molluskocystis Levine,1980 (TM *Molluskocystis pterotracheae* (Stuart,1871) Levine,1980)
@ one named sp. in molluscs
- Metameridae Levine,1980
@ 4 genera & 5 named species in annelids
- Metamera Duke,1910 (TM *M. schubergi* Duke,1910)
@ 2 named sp. in leeches
- Gopalialla Ganpati, Kalavati & Sundaram,1974 (TM *G. morphsae* Ganpati, Kalavati & Sundaram,1974)
@ one named sp. in polychaetes
- Deutomera Bhatia & Setna,1938 (TM *D. cleava* Bhatia & Setna,1938)
@ one named sp. in intestines of polychaetes
- Cognettia Levine,1980 (TM *Cognettia legeri* (Cognetti de Martiis,1911) Levine,1980)
@ one named sp. in coelom of oligochaetes
- Didymophidae L  ger,1892
@ beetles
- Hirmocystidae Grass  ,1953
@ insects & arachnids
- Stenophoroidea Levine, in Lee, Hutner & Bovee,1985
= Solitaroidea Chakravarty,1960
- Stenophoridae L  ger & Duboscq,1904
@ millipedes
- Leidyaniidae Kudo,1954
@ intestines of insects
- Cnemidosporidae Grass  ,1953
@ intestines of millipedes
- Monodontidae Ray & Chakravarty,1933
@ millipedes & insects
- Sphaerocystidae Chakravarty,1960
@ insects

- Trichorhynchidae Ormières, Marquès & Puisségur,1977
@ centipedes
- Dactylophoridae Léger,1892
@ mostly in centipedes
- Stylocephalidae Ellis,1912
@ insects
- Actinocephalidae Léger,1892
@ intestines of arthropods, chaetognaths & chordates
- Actinocephalinidae Léger,1899
@ 34 genera in insects, millipedes, opilionids, chaetognaths & chordates
- Tricystis Hamon,1951 (TM *T. planctonis*) Hamon,1951
@ 2 named sp. in chaetognaths
- Thalicola Ormières,1965 (TM *Thalicola salpae*) (Frenzel,1885) Ormières,1965
@ 4 named sp. in salps
- Acanthosporinae Léger,1892
@ 15 genera in insects, opilionids & chilopods
- Menosporinae Léger,1892
@ 2 genera in insects
- Acutidae Stejskal,1965
@ intestines of honey bees
- Monoicidae Geus,1969
@ intestine of honey bee
- Fusionoidea Stejskal,1965
- Fusionidae Stejskal,1965
@ in a S American cockroach
- NEOGREGARINORIDA Grassé,1953
@ all in insects
- Gigaductidae Filippini,1948
- Ophryocystidae Léger & Duboscq,1908
- Schizocystidae Léger & Duboscq,1908
- Caullervellidae Keilin,1914
- Syncystidae Schneider,1886
- Lipotrophidae Grassé,1953
- Incertae sedis
- Sawayella*** Marcus,1939 (TM *S. polyzoorum*) Marcus,1939
@ one named sp. in midgut of marine bryozoa
* probably not a neogregarine (Grassé,1953)
- COCCIDIASINA*** Leuckart,1879
@ chiefly vertebrate parasites
* an undescribed species in the epithelial cells of the digestive gland of *Antalis entale* is reported from N Bretagne by Arvy,1949
* an unidentified species, probably belonging in Aggregatidae or Caryotrophidae, was reported from Swedish waters in *Boreohydra simplex* by Nyholm,1963
* Lühling,1942 described in detail, without giving it a name, a species from the gut of *Priapulus caudatus* & *Halicryptus spinulosus*
- Hyalosphaera*** Dogiel,1906
* To which family does this belong?
- H. gregarinicola*** Dogiel,1906
@ *Urospora chiridotea*
? Murmansk
- Piridium*** Patten,1936
* To which family does this belong?
- P. sociabile*** Patten,1936
@ in subepithelial connective tissue of ventral part of foot in *Buccinum undatum*
Oresund, England
- AGAMOCOCCIDIORIDA*** Levine,1979
- Rhytidocystidae*** Levine,1979
* only one genus known
- R. henneguyi*** Henneguy,1907
= *Dehornia* Porchet-Henneré,1972
@ 3 named sp. in marine annelids
- R. opheliae*** Henneguy,1907 (TM)
@ *Ophelia bicornis*
?
- R. henneguyi*** de Beauchamp,1913
@ gut epithel of *Ophelia neglecta*
Roscoff
- PROTOCOCCIDIORIDA*** Kheisin,1956
- Eleutheroschizidae*** Chatton & Villeneuve,1936
- Eleutheroschizidae*** Brasil,1906
@ 2 named sp. in intestinal lumen of marine polychaetes
- E. muranicum*** Averinzew,1908
@ *Ophelia limacina*
Murmansk
- E. duboscqi*** Brasil,1906 (TM)
@ intestines of *Scoloplos armiger* & *Protoaricia oerstedi*
Engl. Channel
- Grellidae*** Levine,1973
@ 6 genera & 12 sp. in annelids
- Grellia*** Levine,1973
@ 2 named sp. in coelom of archiannelids & polychaetes
- G. dinophili*** (Grell,1953) Levine,1973 (TM)
@ coelom of *Dinophilus gyroculatus*
- Coelestropha*** Henneré,1963 (TM *C. vivieri*) Henneré,1963
@ 2 named sp. in coelom of polychaetes
- Myriospora*** Lermantoff,1913
@ 4 named sp. in polychaetes
- M. trophoniae*** Lermantoff,1913 (TM)
@ *Stylaroides plumosus*
Gulf of Naples
- Myriospoides*** Henneré,1966 (TM *M. amphiglenae*) Henneré,1966
@ one named sp. in polychaetes
- Defretinella*** Henneré,1966 (TM *D. eulaliae*) Henneré,1966
@ one named sp. in polychaetes
- Mackinnonia*** Janiszewska,1963 (TM *M. tubificis*) Janiszewska,1963
@ 2 named sp. in oligochaetes
- EUCOCCIDIORIDA*** Léger & Duboscq,1910
- ADELEORINA*** Léger,1911
- Adeleidae*** Mesnil,1903
@ 6 genera & 38 named sp. in invertebrates
- Adelea*** Schneider,1875 (TM *A. ovata*) Schneider,1875
@ 4 named sp. in chilopods & molluscs
- Adelina*** Hesse,1911 (TM *A. octospora*) Hesse,1911
@ 17 named sp. in annelids, chilopods, priapuloids & insects
- Klossia*** Schneider,1875 (TM *K. helicina*) Schneider,1875
@ 9 named sp. in molluscs & perhaps birds & mammals
- K. tellinae*** Buchanan,1979
@ in epithelial cells of the kidney of *Tellina tenuis*
Scotland
- Orcheobius*** Schuberg & Kunze,1906 (TM *O. herpobellae*) Schuberg & Kunze,1906
* 3 named sp.
- Legerellidae*** Minchin,1903
@ 1 genus & 6 named sp. in insects, diplopods or nematodes
- Legerella*** Mesnil,1900 (TM *Legerella nova*) (Schneider,1881) Mesnil,1900
- Haemogregarinidae*** Léger,1911 = ***Adelaidae*** Léger,1911
* 3 genera & 378 named sp.
- Cytilia*** Lainson,1981 (TM *C. gomesi*) (Neiva & Pinto,1926) from *Synbranchus marmoratus*)
- C. uncinata*** (Kahn,1978)
@ erythrocytes of *Lycodes vahlii*
?
- Haemogregarina*** Danilewsky,1885 (TM *H. stepanowi*) Danilewsky,1885
@ 301 named sp. in blood cells of fish, reptiles, amphibians, leeches or arthropods
- H. delagei*** Lavaren & Mesnil,1902
@ erythrocytes of *Rajja* spp.
Engl. Channel, Millport
- H. cataphracti*** Henry,1913 (nomen nudum)
@ Blood of *Agonus cataphractus*
Plymouth
- H. anarrhichadis*** Henry,1912
@ blood corpuscles of *Anarrhichas lupus*
Shetlands
- H. cotti*** Brumpt & Lebailly,1904
@ erythrocytes of *Taurulus bubalis* & *Myoxocephalus scorpius*
Engl. Channel
- H. aegelefini*** Henry,1913
@ blood of *Melanogrammus aegelefinus*
Shetlands
- H. pollachi*** Henry,1913 (nomen nudum)
@ blood of *Pollachius pollachius*
Isle of Man
- H. callionymi*** Brumpt & Lebailly,1904
@ blood corpuscles of *Callionymus lyra*
Engl. Channel
- H. bothi*** Lebailly,1905
@ blood corpuscles of *Rhombus maximus* (= *Bothus rhombus*)
Engl. Channel
- H. quadrigemina*** Brumpt & Lebailly,1904
@ blood of *Callionymus lyra*
Engl. Channel
- H. laternae*** Lebailly,1904
@ erythrocytes of *Arnoglossus laterna*
Engl. Channel
- H. minutula*** Neumann,1909
@ blood of *Pomatoschistus minutus*
Mediterranean
- H. blanchardi*** Brumpt & Lebailly,1904
@ blood corpuscles of *Gobius niger*
Engl. Channel
- H. gobii*** Brumpt & Lebailly,1904
@ blood of *Gobius niger*
Engl. Channel
- H. zeugopteri*** Henry,1913 (nomen nudum)
@ blood of *Zeugopterus punctatus*
Isle of Man
- H. flesi*** Lebailly,1904
@ blood corpuscles of *Platichthys flesus*
Engl. Channel, Millport
- H. platessae*** Lebailly,1904
@ blood corpuscles of *Pleuronectes platessa* & *Microcephalus kitt*
Shetlands, Engl. Channel

- H. labri** Henry,1913 (nomen nudum)
@ blood of *Labrus bergylta*
Isle of Man
- H. simondi** Laveran & Mesnil,1901, em. Kirmse,1979
@ blood of *Solea solea*
Engl. Channel
- H. clavata** Neumann,1909
@ blood of *Solea lutea*
Plymouth, Gulf of Naples
- H. bigemina** Laveran & Mesnil,1901
@ blood of *Bleennius* spp., *Zoarces viviparus* & many other species
Engl. Channel, N America, S Africa, S Pacific
- H. polyparita** Neumann,1909
@ blood of *Gobius paganellus*
Plymouth
- H. sachai** Kirmse,1978
@ blood of *Scophthalmus maximus*
Scotland
- H. sp.**
@ *Raja batis*
Engl. Channel
- H. sp.**
@ *Scyllium catulus*
Engl. Channel
- H. sp.** : Coles,1914
@ blood of *Trachinus viperus*
Engl. Channel
- ?**H. sp.** : Henry,1913
@ blood of *Scomber scombrus*
Engl. Channel
- Klossiellidae** Smith & Johnson,1902
@ mammal kidneys
- EIMERIORINA** Léger,1911
- Cardiosporidium** van Gaver & Stephan,1907
* Which family, if belonging in this suborder?
- C. cionae** van Gaver & Stephan,1907
@ *Ciona intestinalis*
?
- Spirocystidae** Léger & Duboscq,1915
@ the only sp. in *Lumbriculus variegatus*
- Selenococcidiidae** Poche,1913
- Selenococcidium** Léger & Duboscq,1910
one named sp.
- S. intermedium** Léger & Duboscq,1910 (TM)
@ intestinal lumen & epithelium of *Homarus gammarus*
?Norway, Mediterranean
- Dobelliidae** Ikeda,1914
- Dobellia** Ikeda,1914
* one named sp.
- D. binucleata** Ikeda,1914 (TM)
@ intestinal mid gut epithelium of the sipunculan *Nephasoma minuta*
?Plymouth
- Aggregatidae** Labbé,1899
@ 7 genera & 29 named sp. in marine invertebrates
* A coccidian similar to *Pseudoklossia pectinis* parasitizes in the renal epithelium of *Ostrea edulis* from Auray, France, sensu Tigé & al.,1977
- A. frenzeli** Frenzel,1885
= *Eucoccidium* Lühe,1902
@ 17 named sp. in decapod crustaceans (merogony) & cephalopoda (gametogony)
- A. eberthi** (Labbé,1895)
= *A. arcuata* ?,1?? = *A. frenzeli* ?,1?? = *A. mamillana* ?,1??
= *A. mingazzinii* ?,1?? = *A. minima* ?,1?? = *A. portundarum* ?,1??
@ sporogony & merogony in intestine of e.g. *Lioacrinus depurator* & *L. arcuatus*, *L. holsatus* & *Necora puber*, Roscoff, Mediterranean, German Bight
- A. coelomica** (Léger,1901)
@ *Pinnotheres pisum* & *P. pinnotheres*; cephalopod host unknown
Engl. Channel, Spain
- A. octopiana** (Schneider,1875) Frenzel,1885 (TM)
= *A. duboscqi* Moroff,1908
= *A. reticulosa* ?,1??
@ *Octopus vulgaris*; crustacean host : ?
Roscoff, Mediterranean
- A. vagans** Léger & Duboscq,1903
@ *Pagurus prideauxi*, *P. cuanensis* & *P. sculptimanus*; cephalopod host unknown
Engl. Channel, Mediterranean
- A. spinosa** Moroff,1906
= *A. jacquemeti* ?,1?? = *A. labbei* ?,1?? = *A. legeri* ?,1??
= *A. ovata* ?,1?? = *A. reticulosa* ?,1?? = *A. schneideri* ?,1??
= *A. stedlecki* ?,1?? = *A. stellata* ?,1??
@ *Octopus vulgaris*; crustacean host unknown
Europe
- A. sepiæ** (Lancaster,1863)
= *Monocystis sepiæ* Lancaster,1863
@ **Sepia*
?Europe
- A. inachi** Smith,1905
@ e.g. *Inachus dorsettensis*; cephalopod host unknown
Mediterranean
- A. sp. : Theodorides & Desportes,1975
@ e.g. *Sergestes arcticus*, *Pasiphæa sivado* & *P. multidentata*
Mediterranean
- A. sp.**
@ *Homarus gammarus*
Roscoff
- A. sp.**
@ *Carcinus maenas*
Engl. Channel
- A. sp.**
@ *Inachus phalangium*
Roscoff
- A. sp.**
@ *Pachygrapsus marmoratus*
Mediterranean
- Angeiocystis** Brasil,1904
@ one named sp. in hearts of polychaetes
- A. audouiniae** Brasil,1904 (TM)
@ *Audouinia tentaculata*
Engl. Channel
- ?**A. sp.**
@ *Hediste diversicolor*
Roscoff
- Merocystis** Dakin,1911
* one named sp. (apparently at least 2)
- M. kathae** Dakin,1911 (TM)
@ *Buccinum undatum*
Öresund, Kattegatt, North Sea, Isle of Man
- M. tellinovum** Buchanan,1979
@ germ cells of female *Tellina tenuis*
Millport
- Pseudoklossia** Léger & Duboscq,1915
@ 5 named sp. with known stages in molluscs
- P. chitonis** Debaisieux,1920
@ digestive gland cells of *Acanthochiton fascicularis*
Engl. Channel
- P. patellae** Debaisieux,1920
@ *Patella vulgata*
Engl. Channel
- P. pelseneeri** (Léger,1897)
@ nephridia of *Macoma balthica* & *Donax vittatus*
Engl. Channel
- P. pectinis** Léger & Duboscq,1917
@ kidney of *Pecten maximus*
Roscoff
- P. glomerata** Léger & Duboscq,1915 (TM - Mediterranean)
@ *Tapes floridus* & *T. virginicus*
Mediterranean
- Grassella** Tuzet & Ormières,1960 (TM) *G. microcosmi* Tuzet & Ormières,1960
@ one named sp. in ascidians
- Ovivora** Mackinnon & Ray,1937
@ one named sp. in echinoids
- O. thalassemae** (Lankester,1885) Mackinnon & Ray,1937 (TM)
@ eggs of *Thalassema neptuni*
Plymouth
- Selysina** Duboscq,1917
@ 3 named sp. in ascidians
- S. perforans** Duboscq,1917 (TM)
@ *Stolonica socialis*
Roscoff
- S. incerta** Duboscq & Harant,1923
@ *Sydinium elegans*
Roscoff
- S. dubosequi** ?,1??
@ *Polycarpa pomaria*
?
- Caryotrophidae** Lühe,1906
@ 2 genera & 2 sp. in polychaetes
- Caryotropha** Siedlecki,1902
- C. mesnili** Siedlecki,1902 (TM)
@ *Eupolymlnia nebula*
Wimereux, Mediterranean
- Dorisella** Ray,1930
- D. scolelepidis** Ray,1930 (TM)
@ gut epithel of *Malacoboceros fuliginosus*
Plymouth
- Cryptosporidiidae** Léger,1911
@ mammals & birds
- Pfeifferinellidae** Grasse,1953
@ 1 genus & 2 sp. in Mollusca
- Pfeifferinella** von Wasielewski,1904 (TM) *P. ellipsoidea* von Wasielewski,1904
@ 2 named sp. in molluscs
- Lankesterellidae** Nöller,1920
@ Vertebrata
- Eimeriidae** Minchin,1903
@ 16 genera, chiefly in vertebrates, but some in invertebrates
- Caryospora** Léger,1904 (TM) *C. simplex* Léger,1904
- C. cheloniae** Leibovitz, Rebell & Boucher,1976
@ *Chelonia mydas*
Grand Cayman

- Crystallospora*** Labbé,1896
- C. crystalloides*** (Thélohan,1893) Labbé,1896
= *Eimeria crystalloides* : Auctt.
@ intestine of *Gadropsaurus mediterraneus* & *G. vulgaris*
Roscoff, W France, Mediterranean
- Eimeria*** A. Schneider,1875 (TM *Eimeria falciformis* (Eimer,1870) Schneider,1875)
= *Coccidium* Leuckart,1879
@ >1050 sp. in vertebrates & invertebrates
- E. ?beauchampi*** Léger & Duboscq,1917
@ liver blind sac epithel of *Glossobalanus minutus*
Biscay
- E. clupearum*** (Thélohan,1892)
@ *Clupea* & *Sprattus*
Denmark
- E. ?epidermica*** Léger & Duboscq,1917
@ proboscis epidermis of *Glossobalanus minutus*
Gulf of Naples
- E. gadi*** Fiebiger,1913
@ swim bladder of *Gadus*, *Melanogrammus* & *Pollachius* (appearing as a creamy yellow mass of spores contained in 26-28 µm Ø oocysts)
Denmark, Nova Scotia
- E. gigantea*** (Labbé,1896)
@ spiral gut of *Lamna nasus*
Roscoff
- E. bigemina*** (Labbé,1896)
@ gut of *Ammodytes tobianus*
Roscoff
- E. anguillae*** (Léger & Hollande,1922)
@ gut of *Anguilla anguilla*
Corsica
- E. variabilis*** (Thélohan,1893) Reichenow,1921
@ gut epithel cells of *Anguilla anguilla*, *Gobius paganellus*, *Crenilabrus melops*, *Taurulus bubalis* & *Lepadogaster gouani*
Roscoff
- E. sardinae*** (Thélohan,1892)
= *E. oxyphora* Dobell,1919
@ male sex glands of *Sprattus sprattus*, *Clupea harengus* & *Sardina pilchardus* & other clupeoids
Denmark, White Sea, W Europe, W France
- E. gasterostei*** (Thélohan,1890)
@ liver of *Gasterosteus aculeatus*
Vilaine
- E. motellae*** (Labbé,1893)
@ gut & pylorus blind sacs of *Gadropsaurus mediterraneus*
Roscoff
- E. thelohani*** (Labbé,1896)
@ liver of *Labrus* sp.
?
- E. raiarum*** ??,19??
@ gut of *Raja batis*
?
- E. ivanae*** Lom & Dyková,1981
@ pyloric caeca of *Serranus cabrilla*
Mediterranean
- E. dingleyi*** ??,19??
@ gut of *Blennius pholis*
?
- E. phoca*** Hsu, Melby & Altman,1974
@ *Phoca vitulina concolor*
Maine
- E. lari*** ??,1???
@ intestinal epithelium of *Larus argentatus*
?
- E. somateriae*** Christiansen,1952
@ kidney epithelium of *Somateria mollissima*
Denmark
- E. bucephala*** Christiansen & Madsen,1948
@ causing summer mortalities among *Bucephala clangula*
Denmark
- E. cruciata*** Thélohan,1892
@ liver of *Trachurus trachurus*
?
- E. sp. : MacKenzie,1978**
@ liver of *Micromesistius poutassou*
off NE Scotland
- E. sp. : Debaisieux,1919**
@ salivary gland cells of *Acanthochiton fascicularis*
Plymoth
- Epieimeria*** Dyková & Lom,1981
- E. isabellae*** Dyková & Lom,1981
@ *Conger conger*
Mediterranean
- Goussia*** Labbé,1896
- G. lucida*** (Labbé,1893) Labbé,1896
= *Eimeria lucida* : Auctt.
@ posterior part of intestine of *Scyliorhinus canicula*, *Squalus acanthias* & *Mustelus vulgaris*
Roscoff, Mediterranean
- G. luciae*** Lom & Dyková,1982
@ *Mullus barbatus*
France
- G. clupearum*** (Thélohan,1894) Labbé,1896
= *Eimeria clupearum* : Auctt.
@ liver of *Clupea harengus*, *Sprattus sprattus*, *Sardina pilchardus*, *Scomber scombrus* & *Engraulis encrasicholus*
Roscoff, W France, London fish-market
- G. cruciata*** (Thélohan,1892) Labbé,1896
= *Eimeria cruciata* : Auctt.
@ liver of *Trachurus trachurus*
Engl. Channel, W France
- G. gadi*** (Fiebiger,1913) Grassé,1953
= *Eimeria gadi* Fiebiger,1913
@ swimbladder of *Gadus morhua*, *Melanogrammus aeglefinus* & *Pollachius virens*
?Baltic, North Sea, Faeroes, Iceland, Nova Scotia
- G. caseosa*** Lom & Dyková,1981
@ *Macrourus berglax*
?
- G. spraguei*** Morrison & Poynton,1989
@ kidney tubules of *Gadus morhua*
Canada
- Iospora*** Schneider,1881 (TM *I. rara* Schneider,1881)
= *Diplospora* Labbé,1893
* 207 sp., usually in vertebrates
- Sarcocystidae** Poche,1913
@ mammals & birds
- Sarcocystinae** Poche,1913
- Sarcocystis*** Lankester,1882
- S.* sp. : Cowan 1966 / : Owen & Kakulas,1967**
@ skeletal muscles of *Globicephala melaena* & *Physeter macrocephalus*
?
- Toxoplasmatinae** Biocca,1957
- Toxoplasma*** Nicolle & Manceaux,1909
- T. gondii*** ??,19??
@ *Phoca vitulina richardi* & California Sea Lion
Alaska
- HAEMOSPORORINA** Danilewsky,1885
- Plasmodiidae** Mesnil,1908
@ chiefly mammals, birds, reptiles & amphibians
- Mesnilium*** Misra, Haldar & Chakravarty,1972 (TM *M. malariae* Misra, Haldar & Chakravarty,1972)
@ one named sp. in fish with leeches as probable vectors
- PIROPLASMASINA** Levine,1961
* Marine fish harbour 3 genera & - as far as known - 5 species, sensu Lom,1984
- PIROPLASMORIDA** Wenyon,1926
- Anthemosomatidae** Levine,1980
@ erythrocytes of Ethiopian rodent
- Babesiidae** Poche,1913
@ mammals
- Theileriidae** du Toit,1918
@ mammals
- Dactylosomatidae** Jakowska & Nigrelli,1955 emend. Levine,1971
@ 3 genera & 21 sp. in amphibians, reptiles & fish
- Dactylosoma*** Labbé,1894 (TM *Dactylosoma ranarum* (Lankester,1882) Wenyon,1926)
@ 6 named sp. in amphibians, reptiles & fish
- Haematractidium** Henry,1910
* Correct family?
- H. scombi*** Henry,1913
@ erythrocytes of *Scomber scombrus*
Irish Sea, E USA
- Haemohormidium*** Henry,1910 (TM *H. cotti* Henry,1910)
@ 14 named sp. in fish, reptiles & amphibians
- H. beckeri*** So,1972
@ 4 species of perciform fishes; vectors : *Johanssonia arctica* & *Platybdella obriti*
Labrador
- Incertae sedis**
- Cristalloidophora*** Dehorne,1934 (TM *C. dodecaceriae* Dehorne,1934)
* one named sp.
- Toxocystis*** Léger & Duboscq,1910
@ one named sp. in marine decapods
- T. homari*** Léger & Duboscq,1910 (TM)
@ rear gut blind sac of *Homarus gammarus* & *Pagurus prideauxii*
Engl. Channel, W France, Mediterranean
- Trophosphera*** Le Calvez,1939 (TM *T. planorbulinae* Le Calvez,1939)
@ one named sp. in Foraminifera
- Globidiellum*** Brumpt,1913 (TM *Globidiellum multifidum* (Neumann,1909) Brumpt,1913)
@ one named sp. in fish
- Joyeuxella*** Brasil,1902
@ one named sp. in annelids
- J. toxoides*** Brasil,1902 (TM)
@ *Lagis koreni*
Calvados
- Rhabdospora*** Laguesse,1895 (TM *R. thelohani* Laguesse,1895)
@ one named sp. in fish epithelial cells
- Echinococcidium*** Porchet,1978 (TM *E. notomasti* Porchet,1978)
@ one named sp. in polychaetes
- MICROSPORA** Sprague,1977
- RUDIMICROSPOREA** Sprague,1977
- METCHNIKOVELLIDA** Vivier,1975
- Metchnikovellidae** Caulley & Mesnil,1914

- @ 3 genera, many sp., most parasiting Eugregarina in annelids
- Metchnikovella** Caullery & Mesnil,1897
= Caulleyetta Dogiel,1922 = *Microsporidiosis* Schereschewsky,1925
- M. hovassei** Vivier,1965
@ *Lecudina pellucida* from *Perinereis cultrifera*
France
- M. spinos** Caullery & Mesnil,
@ *Polyrhabdina brasili* from *Spiro marinensis*. *Polyrhabdina spinos* from *Malacoceros fuliginosus* & *P. bifurcata* from *Malacoceros fuliginosus*
Plymouth
- M. brasili** Caullery & Mesnil,
@ *Polyrhabdina brasili* from *Spiro marinensis*?
- M. incurvata** Caullery & Mesnil,
@ *Polyrhabdina pygospionis* from *Pygospio elegans*?
- M. oviformis** Caullery & Mesnil,
@ *Polyrhabdina pygospionis* from *Pygospio elegans*?
- M. nereidis** Caullery & Mesnil,
@ *Lecudina pellucida* from *Platynereis* sp.
?
- M. polydorae** Reichenow,1932
@ *Selenidium ?cruzi* from *Polydora* sp.
Helgoland
- M. selenidii** Awerinzew,1908
@ *Selenidium* sp. from *Ophelia limacina*
Murmansk
- M. minima** Caullery & Mesnil,
@ *Selenidium cirratum* from *Audoinia* sp.
Belle-Isle-en-mer
- M. berliozi** Arvy,1952
@ *Lecudina franciana* from *Phascolion strombi*
Dinard
- M. caulleryi** ??,19??
@ *Polyrhabdina polydorae* from *Polydora flava*?
- Amphiambylys** Caullery & Mesnil,1914
- A. capitellidis** Caullery & Mesnil,1914?
@ *Ancora* sp. from *Capitellides giardi*?
- A. capitellae** Caullery & Mesnil,1914?
@ ?*Lecudina* sp. from *Capitella capitata*
- A. ancorae** Reichenow,1932
@ *Ancora sagittata* from *Capitella capitata*
Helgoland
- A. caulleryi** (Mackinnon & Ray,19??)
@ *Polyrhabdina polydorae* from *Polydora flava*?
- Amphiacantha** Caullery & Mesnil,1914, non Popofsky,1904, nec Matsumoto,1917 (Echinodermata)
* should need a new name
- A. longa** Caullery & Mesnil,1914
@ *Lecudina elongata* from *Lumbrineris tingens*?
- A. ovalis** Stubblefield,1955
@ *Lecudina* sp. from *Lumbrineris* spp.
California
- A. attenuata** Stubblefield,1955
@ *Lecudina* sp. from *Lumbrineris* spp.
California
- MICROSPORA** Delphy,1963
- MINISPORIDA** Sprague,1972
- Chytridiopsidae** Sprague, Ormieres & Manier,1972
?(@ the only genus in gut wall of beetles, myriapods & snails)?
- Steinhauida** Sprague, Ormieres & Manier,1972
- S. ovicola** (Léger & Hollande,1917)
= *Chytridiopsis ovicola* Léger & Hollande,1917
@ ova in females of *Ostrea edulis*
Marennes (France)
- S. mytilolum** (Field,1923)
= *Haplosporidium mytilolum* Field,1923
@ ova in females of *Mytilus edulis* & *M. galloprovincialis*
Atlantic USA, Mediterranean
- Chytridiopsis** Schneider,1884
= *Chytridioides* Tregouboff,1913
- Hesseidae** Ormieres & Sprague,1973
@ the only genus infects epithelial cells of Diptera larvae
- Burkeidae** Sprague,1977
- Buxthuidae** Larsson,1980
- MICROSPORIDA** Balbiani,1882
- PANSPOROBLASTINA** Tuzet, Maurand, Fize, Michel & Fenwick,1971
- Glugeidae** Thélohan,1892
- Glugea** Thélohan,1891
- G. anomala** (Moniez,1887) Gurley,1893 (TM)
?= *G. gasterostei* Voronin,1974
?= *G. weisenbergi* Sprague & Vernick,1968
- G. subcutan** tissue & different organs of *Gasterosteus aculeatus*, *Pungitius pungitius* & *Pomatoschistus minutus*
? *Apeltes quadratus*
Baltic, Kattegatt, Bohusl. & fresh water, & ?N America
- G. laverani** Caullery & Mesnil,1899
@ *Malacoboceros fuliginosus* & *Scoloplos armiger*
Engl. Channel
- G. destruens** Thélohan,1892
@ muscles of *Callionymus lyra*, *Melanogrammus aeglefinus*
?Plymouth, Roscoff, Concarneau
- G. cordis** Thélohan,1895
@ heart of *Sardina pilchardus*
Marseille
- G. punctifera** Thélohan,1895
@ Eye muscle of *Pollachius pollachius*. This is probably also the Microsporean species encountered frequently in European *Micromesistius poutassou*
W France
- G. acuta** Thélohan,1895
@ fin muscles of *Entelurus aequeorus* & *Syngnathus acus*
Roscoff, Concarneau
- G. hertwigi** Weissenberg,1911
@ subcutan tissue & different organs of *Osmerus eperlanus* & *O. mordax*
Rügen, Canada
- G. stephani** (Hagenmüller,1899) Woodcock,1904
@ gut wall of "0 group" of *Limanda limanda*, *Pleuronectes platessa* & 9 other flatfish species
Baltic, W Sweden, Helgoland, White Sea, Mediterranean, Black Sea, NW Atlantic
- G. capverdensis** Lom, Gaevskaya & Dyková,1980
@ *Myctophum punctatum*
Cape Verde
- C. berglax** Lom & Laird,1976
@ *Macrourus berglax*
?
- G. atherinæ** Berrebi,1978
@ *Atherina boyeri*
French Mediterranean
- Pleistophora** Gurley,1893
= *Plistophora* Gurley,1893
@ e.g. muscles of fish, insects
- P. ehrenbaumi** Reichenow,1929
@ muscles of *Anarrhichthys lupus* & *A. minor*
Baltic, North Sea
- P. typicalis** Gurley,1893, em. Canning & Nicholas,1980 (TM)
@ muscles & liver of *Blennius pholis*, *Taurulus bubalis*, *Myoxocephalus scorpius*, *Pungitius pungitius*
Roscoff, Concarneau
- P. hippoglossoides** Bosanquet,1910
@ forming whitish nodules (10 mm Ø) in fin muscles of *Hippoglossoides platessoides* & other flatfish species
North Sea
- P. shiplei** (Drew,1910)
@ muscles & gut of *Trisopterus luscus*
Plymouth
- P. gadi** Polyansky,1955
?= *P. sp.*; Drew,1909
@ muscles of *Gadus morhua*
?Iceland
- P. destruens** Delphy,1916
@ muscles of *Liza aurata*
Engl. Channel
- P. littoralis** Canning & Nicholas,1980
@ muscles of *Blennius pholis*, other blennies & cottids
Europe
- P. giganta** ??,19??
@ muscles of *Crenilabrus melops*
?
- P. duodecimae** Lom, Gayevskaya & Dyková,1981
@ muscle fibres of *Coryphaenoides nasutus*
Atlantic Ocean
- Mitoplustophora** Codreanu,1966 (TM *M. angularis* Codreanu,1966)
- Vavraia** Weiser,1977 (TM *V. cuculus* (Weiser,1947) Weiser,1977)
- Baculea** Loubès & Akbarieh,1978
- B. daphniae** Loubès & Akbarieh,1978 (TM)
@ intestinal epithel of *Daphnia pulex*
?
- Loma** Morrison & Sprague,1981
- L. morhua** Morrison & Sprague,1981 (TM)
@ gills of *Gadus morhua*
?
- L. brachialis** (Nemeczek,1911) Morrison & Sprague,1981
= *Nosema brachiale* Nemeczek,1911
@ gills of *Melanogrammus aeglefinus*, *Gadus morhua* & other Gadidae
W Sweden, Vienna fish market
- Pseudopleistophoridae** Sprague,1977
* 2 genera
- Octosporea** Flu,1911 (TM *O. muscaedomesticae* Flu,1911)
- O. effeminans** Bulnheim & Vávra,1968
@ *Gammarus duebeni*
Elbe estuary, W Baltic
- Pseudopleistophora** Sprague,1977 (TM *P. szollissi* Sprague,1977)
- Duboscuidae** Sprague,1977
* 2 genera
- Duboscqia** Pérez,1908

| | |
|--|---|
| <i>Trichoduboscqia</i> Léger,1926 | Engl. Channel |
| <i>Thelohanidae</i> Hazard & Oldacre,1975 * 13 genera | <i>Spraguidae</i> Weissenberg,1976 * 1 genus |
| <i>Thelohania</i> Henneguy,1892 = <i>Orthothelohania</i> Codreanu & Codreanu-Balcescu,1974 | <i>Spraguea</i> Weissenberg,1976 |
| <i>T. giardi</i> Henneguy,1892 @ muscles of <i>Crangon crangon</i> | <i>S. lophii</i> (Doflein,1898) Sprague,1977 = <i>Nosema lophii</i> (Doflein,1898) @ nerve system gangliae of <i>Lophius piscatorius</i> , <i>L. americanus</i> & <i>L. gastrophysus</i> North Sea, Iceland, Liverpool, W France, Mediterranean, E USA |
| Engl. Channel | |
| <i>T. herediteria</i> Buinheim,1971 @ ovary of <i>Gammarus duebeni</i> Bornholm, Elbe estuary | <i>Pereziidae</i> Loubès, Maurand, Comps & Campillo,1977 * 2 genera |
| <i>T. maenadis</i> Pérez,1904 @ muscles of <i>Carcinus maenas</i> Arcachon | <i>Perezia</i> Léger & Duboscq,1909 |
| <i>T. paguri</i> ?Pérez,1927? @ <i>Pagurus bernhardus</i> France | <i>P. lankesteriae</i> Léger & Duboscq,1909 (TM) = <i>Glugea lankesteriae</i> : Auctt. @ infects the gregarine <i>Lankesteria ascidiae</i> from <i>Ciona intestinalis</i> Cette (Mediterranean) |
| <i>T. baueri</i> Voronin,1974 @ 3- & 9-spined sticklebacks | <i>Ameson</i> Sprague,1977 @ Includes former <i>Nosema</i> species infecting arthropods. See Sprague,1977 |
| ? <i>T. sp.</i> : Mercier,1906 @ body muscles & heart muscle of <i>Talitrus</i> sp. Roscoff | <i>A. pulvis</i> (Pérez,1905) = <i>Nosema pulvis</i> Pérez,1905 @ muscles of <i>Carcinus maenas</i> Arcachon |
| <i>Indosporus</i> Overstreet & Weidner,1974 | <i>A. sp.</i> = <i>Nosema</i> sp. : Mercier & Poisson,1926 @ muscles of <i>Mysis spiritus</i> Engl. Channel |
| <i>I. octospora</i> (Henneguy,1892) = <i>Telohania octospora</i> Henneguy,1892 | <i>A. sp.</i> = <i>Neomyasis vulgaris</i> Engl. Channel |
| = <i>Indosporus spraguei</i> Overstreet & Weidner,1974 (TM) @ <i>Palaeomon serratus</i> , <i>P. elegans</i> & <i>Palaeomon rectirostris</i> | <i>Cougourdeillidae</i> Poisson,1953 * 1 genus |
| Roscoff, Plymouth, W France | <i>Caudosporidae</i> Weiser,1958 * 4 genera |
| <i>Ormieresia</i> Viviers, Bouix & Manier,1977 | <i>Nosematidae</i> Labbé,1899 * 3 genera |
| <i>O. careni</i> Viviers, Bouix & Manier,1977 (TM) @ <i>Carcinus mediterraneus</i> Czerniavsky,1884 | <i>Ichthyosporidium</i> Caullery & Mesnil,1905 |
| Mediterranean | <i>I. giganteum</i> (Thélohan,1895) Swarczewsky,1914 = <i>Glugea gigantea</i> Thélohan,1895 = <i>Plistophora laborum</i> Le Danois,1910 = <i>Ichthyosporidium phymogenes</i> ?,??? |
| <i>Burenellidae</i> Jouvenaz & Hazard,1978 * 2 genera | @ mesenteria of <i>Crenilabrus melops</i> Roscoff |
| <i>Amblysporididae</i> Weiser,1977 * 3 genera in Diptera larvae | <i>Nosema</i> Nägeli,1857 @ Lepidoptera. (parasites of other hosts have been transposed to other genera) |
| <i>Culicosporidae</i> Weiser,1977 * 2 genera, probably lacking marine species. | <i>N. frenzeliae</i> Léger & Duboscq,1909 @ <i>Cephaloidophora conformis</i> (a gregarine) from <i>Pachygrapsus marmoratus</i> Banyuls-sur-mer |
| <i>Gurleyidae</i> Sprague,1977 | * Should be moved to another genus. Which? ? |
| <i>Gurleya</i> Doflein,1898 = <i>Marssonella</i> Lemmermann,1900 | <i>N. vivieri</i> Vinckier, Devauchelle & Prensier,1970 @ monocystid gregarine living in the coelom of an unspecified nemertean * Should be moved to another genus. Which? Wimereux |
| <i>G. nova</i> Hovasse,1950 @ hyperparasite in <i>Spirobutschilliella chattoni</i> from <i>Pomatoceros triquetus</i> | <i>N. marionis</i> ?Kudo,1924 @ <i>Leptothecca coris</i> from <i>Coris julis</i> * Should be moved to another genus. Which? Mediterranean |
| Banyuls-sur-mer | |
| <i>Pyrotheca</i> Hesse,1935 | <i>N. cotti</i> ?,?? @ forming 7mm Ø nodes in testes of <i>Taurulus bubalis</i> * Should be moved to another genus. Which? ? |
| <i>Stempellia</i> Léger & Hesse,1910 | <i>Hirsutosporos</i> Batson,1983 |
| <i>S. muelleri</i> ?,?? @ <i>Gammarus</i> spp. Elbe estuary | <i>Mřížekidae</i> Léger & Hesse,1922 * 2 genera in oenocytes of Oligochaeta |
| <i>Norlevinea</i> Vávra,1985 | <i>Jirovecia</i> Weiser,1977 |
| <i>Episeptum</i> Larsson,1986 | <i>Mrazekia</i> Léger & Hesse,1916 |
| <i>Telomyxidae</i> Léger & Hesse,1910 * 4 genera | <i>M. piscicola</i> Cépède,1924 @ pylorus appendix of <i>Merlangius merlangus</i> Bergen |
| <i>Berwaldia</i> Larsson,1981 | Unclassified Microspora (Collective group name) |
| <i>B. singulais</i> Larsson,1981 (TM) @ <i>Daphnia pulex</i> Skåne | <i>Microsporidium</i> Balbiani,1884 |
| <i>Tuzetiidae</i> Sprague, Tuzet & Maurand,1977 * 4 genera | <i>M. ovoideum</i> (Thélohan,1895) Sprague,1977 = <i>Nosema ovoideum</i> (Thélohan,1895) @ liver of <i>Gadropsaurus mediterraneus</i> & several other fish species Scotland, Roscoff |
| <i>Tuzetia</i> Maurand & al.,1971 @ infects aquatic hosts | <i>M. cotti</i> (Chatton & Courrier,1923) Lom,1984 = <i>Nosema cotti</i> Chatton & Courrier,1923 @ testis of <i>Taurulus bubalis</i> Roscoff |
| <i>T. infirma</i> (Kudo,1921) = <i>Nosema infirmum</i> Kudo,1921 (TM) @ Cyclopoid Copepoda | ASCETOSPORA |
| <i>APANSPOROBLASTINA</i> Tuzet, Maurand, Fize, Michel & Fenwick,1971 | STELLATOSPOREA |
| <i>Unikaryonidae</i> Sprague,1977 * 7 genera | OCCLUSOSPORIDA |
| <i>Tetramicra</i> Matthews & Matthews,1980 | <i>Marteilia</i> Grizel, Comps, Bonami, Cousserans, Duthoit & Le Pennec,1974 |
| <i>T. brevifilum</i> Matthews & Matthews,1980 (TM) @ skeletal muscles of <i>Scophthalmus maximus</i> Britain | <i>M. refringens</i> Grizel, Comps, Bonami, Cousserans, Duthoit & Le Pennec,1974 (TM - Aber Wrach, W Bretagne) @ causing "Aber disease" in <i>Ostrea edulis</i> ; also sporadically found in <i>Mytilus edulis</i> & <i>Cerastoderma edule</i> , albeit the coccle inhabitant may however be another species |
| <i>Unikaryon</i> Canning & al.,1974 (TM <i>U. piriformis</i>) Canning, Lai & Lie,1974 = <i>Oligosporidium</i> Codreanu-Balcescu, Codreanu & Traciuc,1981 | |
| <i>U. mytilicola</i> ?Vallmitjana & Durfort,1976? @ hyperparasite in <i>Mytilicola intestinalis</i> Steuer from <i>Mytilus</i> Spanish Mediterranean | |
| <i>U. legeri</i> (Dollfus,1912) = <i>Nosema legeri</i> Dollfus,1912 @ hyperparasite in <i>Meiogymnophallus minutus</i> from <i>Cerastoderma edule</i> & <i>Gymnophallus strigatus</i> from razor clams Exe estuary, Wimereux | |
| <i>U. spelotremae</i> (Guyénot, Naville & Ponse,1925) = <i>Nosema spelotremae</i> Guyénot, Naville & Ponse,1925 @ <i>Spelotrema carcinii</i> (from <i>Carcinus maenas</i>) * Correct genus? | |

N & W Bretagne

M. maurini Comps, Pichot & Papagianni, 1982
@ *Mytilus galloprovincialis*
France (in mussels imported from the lagoon of Venice)

Paramarteilia Ginsburger-Vogel & Desportes, 1979
P. orchestiae Ginsburger-Vogel & Desportes, 1979 (TM)
@ *Orchestia gammarellus* (Pallas)
France

BALANOSPORIDA

Haplosporidiidae

Haplosporidium Lühe, 1900
= *Aplosporidium* Caullery & Mesnil, 1899
* 23 species

H. ascidiarum Duboscq & Harant, 1923
@ *Syndium elegans* & *Amaroucium proliferum*
Roscoff

H. tapetis Vilela, 1951
@ *Tapes decussatus*
Portugal

H. nemertis Debaissieux, 1920
@ *Lineus bilineatus*
Plymouth

H. malacobdellae Jennings & Gibson, 1968
@ *Malacobdella grossa*
Yorkshire

H. heterocirri Caullery & Mesnil, 1899
@ *Heterocirrus viridis* ?= *Dodecaceria* sp.
Engl. Channel

H. caulleryi Mercier & Poisson, 1922
@ *Neanthes fucata*
Engl. Channel

H. potamillae Caullery & Mesnil, 1905
@ *Potamilla vellei*
Engl. Channel

H. marchouxii Caullery & Mesnil, 1905
@ *Salmacina dysteri*
Engl. Channel

H. scolopli Caullery & Mesnil, 1899
@ *Scoloplos armiger*
Engl. Channel

H. mytilorum Field, 1922
@ *Mytilus edulis*
USA

H. lusitanicum Azevedo, 1984
@ *Patina pellucida*
Portugal

H. sp. : Caullery & Mesnil, 1905
@ *Eupolymlnia nesidenensis*
Engl. Channel

H. sp. : Duboscq & Harant, 1923
@ *Cerastoderma edule*
?

H. sp., non Minchinia armoricana : Vivarès & al., 1981
@ connective tissues of gills and digestive gland of *Ostrea edulis*
French Atlantic coast

Minchinia Labbé, 1896
* 4 species. May be an invalid genus; if so, *Haplosporidium* is so far the only alternative genus.

M. chitonis (Lankester, 1885)
= *Klossia chitonis* Lankester, 1885 (TM - Roscoff)
@ *Lepidochiton cineraceus* (chiefly in liver)
E Scotland, SE, S & SW Britain, Dinard

M. armoricana van Banning, 1977
@ connective tissues of *Ostrea edulis*
Netherlands (import oysters from Bretagne), Bretagne

M. dentali (Arvy, 1949)
= *Haplosporidium dentali* Arvy, 1949
@ *Antalis entale*
N Bretagne

M. ascidiarum (Ormières, 1965)
= *Haplosporidium ascidiarum* Ormières, 1965
@ *Aplidium elegans* & *A. nordmanni*
?

Bonamia Pichot, Comps, Tigé, Grizel & Rabouin, 1980
* Family placement tentative

B. ostreae Pichot, Comps, Tigé, Grizel & Rabouin, 1980 (TM)
@ in cytoplasm of haemocytes of *Ostrea edulis*
?France

Anurosporidiidae

Urosporidium Caullery & Mesnil, 1905
@ hyperparasites of larval trematodes & a nematode parasiting marine bivalves
* 6 species

U. fuliginosum Caullery & Mesnil, 1905 (TM)
@ *Syllis gracilis* & *Sphaerosyllis ?erinaceus*
Engl. Channel

Anurosporidium Caullery & Chappellier, 1906

A. pelseneeri Caullery & Chappellier, 1906 (TM)
@ cercariae (*Cercaria lutea* ?= *C. pectinata* & *Bucephalus haimeanus*) from *Donax vittatus* & ?*Barnea candida*
Boulogne, ?W France

PARAMYXEA

* with one sp. only

PARAMYXIDA

Paramyxidae

Paramyxxa Chatton, 1911

P. paradoxa Chatton, 1911 (TM - Banyuls-sur-Mer)
@ intestinal cells of Polychaete larvae
S France

STEPHANOPOGONOMORPHA

* Taxon incertae sedis, sensu Patterson & Zölfel, in Patterson & Larsen, 1992

PRIMOCILIATIDA

Corliss, 1974

Stephanopogonidae

Corliss, 1961

Stephanopogon

Entz Sr., 1884

S. colpoda Entz Sr., 1884

Kieler Bucht, Helgoland, Plymouth, Naples

S. mesnili Lwoff, 1923

Plymouth, Roscoff, Banyuls-sur-mer

S. apogon Borror, 1965

New Hampshire

CILIOPHORA

Doflein, 1901

POSTCILIODESMATOPHORA

Gerassimova & Seravin, 1976

KARYORELICTEA

Corliss, 1974

* all marine or brackish water living

PROTOSTOMATIDA

Small & Lynn, in Lee, Hutner & Bovee, 1985

Trachelocercidae

Kent, 1881

Gruvelina Delphy, 1939

* incertae sedis

G. longissima Delphy, 1939 (TM)

Dinard

Trachelocerca Ehrenberg, 1833

T. entzi Kahl, 1927 (TM)

Kiel, Helgoland, Plymouth, Roscoff

T. tenuicollis Quennerstedt, 1867

Baltic, Halland, Roscoff

T. multinucleata Dragesco, 1960

Roscoff

T. binucleata Dragesco, 1960

Finistère

T. minuta Dragesco, 1960

Roscoff

T. dogieli Raikov, 1957

?

Tracheloraphis Dragesco, 1960

T. discolor Raikov, 1962

Sylt, Barents Sea, White Sea

T. phoenicopterus (Cohn, 1866) (TM)

= *Trachelocerca phoenicopterus* Cohn, 1866

Sylt, Barents Sea, Murmansk, White Sea, Plymouth, Roscoff, W Africa, Cape Cod

T. kahli Raikov, 1962

= *Trachelocerca phoenicopterus* : Kahl, 1930 ; Dragesco, 1960, non Cohn, 1866

W Baltic, Kieler Bucht, Hiddensee, Königshafen, Sylt, White Sea, French Atlantic, Mediterranean, E USA

T. remanei Dragesco, 1960

Roscoff

T. drachi Dragesco, 1960

Roscoff

T. incaudatus (Kahl, 1933)

= *Trachelocerca incaudata* Kahl, 1933

* belonging to the "incaudatus" group

Kieler Bucht, Drøbak, White Sea, Yorkshire, Roscoff, Mediterranean, Black Sea, Bermuda

T. incaudatus f. **quadrimicronucleata** Raikov & Kovaleva, 1968

* belonging to the "incaudatus" group

Japan Sea

T. fasciolatus (Sauerbrey, 1928)

= *Trachelocerca fasciolata* Sauerbrey, 1928

S Baltic, Helgoland, Roscoff

T. teissieri Dragesco, 1960

Roscoff

T. prenanti Dragesco, 1960

Roscoff

T. prenanti f. **multicineta** Raikov & Kovaleva, 1968

Sylt, ?White Sea, French Atlantic, Mediterranean, Black Sea

T. prenanti f. **oligocineta** Raikov & Kovaleva, 1968

White Sea, Yorkshire, Japan Sea

T. hyalinum Dragesco, 1960

Roscoff

T. similis Raikov & Kovaleva, 1968

Japan Sea

T. sarmaticus Agamaliev & Kovaleva, 1966

?

- T. swedmarki** Dragesco,1960
* belonging to the "mcaudatus" group
Roscoff
- T. gracilis** Dragesco,1960
* belonging to the "mcaudatus" group
Roscoff
- T. enigmaticus** Dragesco,1960
Roscoff
- T. griseus** (Kahl,1933)
= *Trachelocerca grisea* Kahl,1933
Kieler Bucht, Roscoff
- T. haloetes** Borror,1973
Bahamas hypersaline waste water
- T. arenicolus** (Kahl,19???)
S Baltic
- T. margaritatus** (Kahl,1933)
S Baltic
- T. coluber** (Kahl,19???)
S Baltic, Helgoland
- T. subviridis** Sauerbrey,19??
S Baltic
- T. monocaryon** Dragesco,1965
* belonging to the "mcaudatus" group
?
- T. stephani** Dragesco,1965
* belonging to the "mcaudatus" group
?
- Kentrophoridae** Jankowski, in Lee, Hurner & Bovee,1985
- Ciliofaurea** Dragesco,1960
* Incertae sedis. Correct family?
- C. mirabilis** (Dragesco,1954)
Öresund, Sylt, Yorkshire, French Atlantic, Mediterranean
- Trachelonema** Dragesco,1960
- T. oligostriata** Raikov,1962
Sylt, White Sea, French Atlantic
- T. longicollis** Dragesco,1960
Roscoff
- T. minima** Dragesco,1960
Roscoff
- T. grassei** Dragesco,1960
Roscoff
- Kentrophoros** Sauerbrey,1928
= *Centrophorella* = *Centrophorus* (homonym)
W Baltic, Kieler Bucht, Öresund, Sylt, Helgoland, Norway, White Sea, French Atlantic, W Africa, Mediterranean, Black Sea, Cape Cod
- K. lanceolatum** (Fauré-Fremiet,1951)
= *Centrophorella lanceolata* Fauré-Fremiet,1951
S Baltic, Cape Cod
- K. grandis** Dragesco,19???
?
- K. latum** Raikov,1962?
Öresund, Black Sea
- K. fistulosa** Fauré-Fremiet,1950?
Bretagne
- K. longissima**
?
- K. trichocystus**
?
- LOXODIDAE** Jankowski, in Lee, Hurner & Bovee,1985
- Loxodidae** Bütschli,1889
- Loxodes** Ehrenberg,1833
- Remanella** Kahl,1933
- R. multinucleata** Kahl,1933
Kieler Bucht, Sylt, White Sea, French Atlantic, Mediterranean, Black Sea
- R. faurei** Dragesco,1954
Sylt, French Atlantic, Mediterranean, Black Sea
- R. rugosa** Kahl,1933
W Baltic, Öresund, Sylt, Helgoland, N Russia, Plymouth, French Atlantic, Mediterranean, Black Sea, E USA
- R. rugosa v. unicorpulculata** Kahl,19??
S Baltic
- R. unirugosa** Hartwig,1973
Sylt
- R. margaritifera** Kahl,1933
S Baltic, Öresund, Sylt, Helgoland, Yorkshire, Plymouth, W France, Black Sea, E USA, Brazil
- R. brunnea** Kahl,19??, non : Dragesco,1965
S Baltic, Öresund
- R. gigas** Dragesco,19??
Öresund
- R. minuta** Dragesco,19??
Öresund
- R. trichocystis** Dragesco,19???
- R. swedmarki** Dragesco,19???
?
- R. caudata** Dragesco,19???
?
- R. granulosa** Kahl,19??
S Baltic
- R. obtusa** Fauré-Fremiet,1951
Cape Cod
- Cryptopharyngidae** Jankowski,1967
- Cryptopharynx** Kahl,1928
- C. setigerus** Kahl,19???
Öresund
- PROTOHETEROTRICHIDA** Nouzarède,1977
- Geleidae** Kahl,1933
- Avelia** Nouzarède,1977
- Corlissia** ?Dragesco,1960?
= *Corlissa* ?Dragesco,19??
* Incertae sedis. Family placement tentative.
- C. picta** (Dragesco,19??) (TM)
?
- Geleia** Kahl,1933
- G. nigriceps** Kahl,1933
Kieler Bucht, Öresund, Drøbak, N Russia, Sylt, Atlantic France, Black Sea, Bermuda
- G. fossata** Kahl,19???
S Baltic, Öresund, Plymouth, Cape Cod
- G. orbis** Fauré-Fremiet,1950
Öresund, Drøbak, Barents Sea, White Sea, Bretagne, Mediterranean, Black Sea, Cape Cod, Bermuda, Brazil
- G. decolor** Kahl,19???
S Baltic, Öresund, Helgoland, ?Bretagne, ?Cape Cod
- G. gigas** Dragesco,19???
?
- G. luci** Dragesco,19???
?
- G. swedmarki** Dragesco,19???
?
- G. simplex** Fauré-Fremiet,1951
Cape Cod
- PROTCRUZIIDAE** Jankowski, in Lee, Hurner & Bovee,1985
- Protocruziidae** Jankowski, in Lee, Hurner & Bovee,1985
- Protocruzia** de Faria, da Cunha & Pinto,1922
- P. pigerimma** (Cohn,18??) da Cunha,19???
Plymouth
- P. granulosa** Kahl,19???
Helgoland
- P. depressa** Ammermann,1968
New Hampshire
- P. tuzeti** Villeneuve-Brachon,1940?
?
- SPIROTRICHEA** Bütschli,1889
- HETEROTRICHIA** von Stein,1859
- HETEROTRICHIDA** von Stein,1859
- HETEROTRICHINA** von Stein,1859
- Blepharismidae** Jankowski, in Lee, Hurner & Bovee,1985
- Anigsteinia** Isquith,1968
- Blepharisma** Perty,1849
- B. clarissimum** Anigstein,1???
Askö, Öresund, Ifsejord, Cape Cod
- B. clarissimum f. arenicola** Kahl,1932
Kieler Bucht, Sylt, N Russia, Yorkshire, French Atlantic, Mediterranean, Black Sea, E USA
- B. clarissimum v. longissimum** Kahl,19???
S Baltic
- B. salinarum** (Florentin,1???)
Askö, S Baltic, Öresund
- B. grayi** Hartwig & Parker,1977
Yorkshire
- B. tardum** Kahl,19???
?
- B. dileptus** Kahl,19???
?
- B. vestitum** Kahl,19???
?
- B. velatum** Vacelet,1961
S France
- B. sp. ; Fenichel,1968**
Öresund
- Parablepharisma** Kahl,1932

- P. pellitum* Kahl,19??
Kieler Bucht, Öresund, Cape Cod
- P. chlamydophorum* Kahl,19??
Öresund
- P. bacteriophaga* Villeneuve-Brachon,1940
Sylt, E USA
- P. collare* Kahl,19??
Öresund
- Pseudoblepharisma* Kahl,1926
- Climacostomidae Repak,1972
- Climacostomum* von Stein,1859
* Fresh water genus with one species occurring in brackish water
- C. vires* (Ehrenberg,18??)
Askö
- Copemetopus* Villeneuve-Brachon,1940
- C. subsalsus* Villeneuve-Brachon,1940?
?
- Fabrea* Henneguy,1890
- F. salina* Henneguy,1890?
?France
- Condylostomatidae Kahl, in Doflein & Reichenow,1929
- Condylostoma* Bory de St. Vincent,1826
- C. patulum* Claparède & Lachmann,1858
S Baltic, Öresund
- C. arenarium* Spiegel,1926
Askö, Hiddensee, Kieler Bucht, Öresund, Sylt, Helgoland, N Russia, Yorkshire, French Atlantic, Mediterranean, Black Sea, Mauretania, E USA, Bermuda, Brazil
- C. arenarium* f. *proturostyla* Kahl,19??
S Baltic
- C. remanei* Spiegel,1928
Askö, Kieler Bucht, Öresund, Sylt, Norway, N Russia, ?Yorkshire, French Atlantic, Mediterranean, Black Sea, E USA, Bermuda, Brazil
- C. remanei* v. *oxyura* Dragesco,1960
?
- C. feldii* Hartwig,1973
= *C. remanei* : Fjeld,1955, non Spiegel,1928
Sylt, Drøbak, Yorkshire
- C. patens* (O.F. Müller,1786) Dujardin,18??
= *Trichoda patens* O.F. Müller,1786
Kieler Bucht, Halland, Plymouth, Whitstable, Cape Cod
- C. acuta* Dragesco,1954
?
- C. ancestralis* Villeneuve-Brachon,1940?
?
- C. tenuis* Fauré-Fremiet,1958
Yorkshire, French Atlantic, Brazil
- C. psammophilum* Bock,1952
S Baltic
- C. enigmatica* Dragesco,1954
?
- C. caudatum* Lauterborn,1908
- Spirostomidae von Stein,1867
- Gruberia* von Stein,1867
- G. uninucleata* Kahl,19??
Plymouth
- G. lanceolata* (Gruber,1???)
?
- G.* sp. : Fenchel,1968
Öresund
- Spirostomum* Ehrenberg,1833
- S. teres* Claparède & Lachmann,18??
* this is the only species of the genus, which may occur in brackish water, the other species being strictly fresh water forms
Askö
- Stentoridae Carus,1863
- Stentor* Oken,1815
- S. acrobaticus* Silén,1947
* on *Fucus serratus*
Gullmann
- Chattonidiidae Villeneuve-Brachon,1940
* not included in Snall & Lynn,1985
- Chattonidium* Villeneuve,1937
- C. setense* Villeneuve-Brachon,19?? (TM)
* in polluted brackish water
?
- COLIPHORINA** Jankowski,1967
- Folliculinidae Dons,1914
- Ampullofolliculina* Hadzi,1951
- A. lageniformis* Hadzi,1951 (TM)
- = *Folliculina (Vorticella) ampulla* : Andrews,1949, non (O.F. Müller,1786)
@ oyster shells
Delaware Bay
- Ascobius* Henneguy,1884
= *Semifolliculina* Dons,1914 (p.p.)
- A. claparedi* Hadzi,1951
= *Freia ampulla* Claparède & Lachmann,1858, non *Vorticella ampulla* O.F. Müller,1786
- Bergen
- A. faurefremieti* Hadzi,1951
= *Folliculina ampulla* : Fauré-Fremiet,1932, non (O.F. Müller,1786)
Aquarium in Concasseau
- A. simplex* (Dons,1918)
= *Semifolliculina simplex* Dons,1918 = *Platycola nigra* Walles,1???
@ *Delesseria*
?Kieler Bucht, Trondheim, W France
- A. sileni* Hadzi,1951
= *Folliculina simplex* : Silén,1947, non Dons,1917
@ *Spirorbis spirorbis*
Gullmarfjord
- Atriofolliculina* Hadzi,1951
- A. andrewsi* Hadzi,1951
= *Parafolliculina amphora* : Andrews,1946, non Andrews,1941
@ *Elodea*
Chesapeake Bay
- A. hirundo* (Kent,1881-82) Hadzi,1951
?
- A. faureana* Hadzi,1951
= *Parafolliculina hirundo* Fauré-Fremiet,1936, non Kent,1881-82
?
- Aulofolliculina* Hadzi,1951
- A. labyrinthica* Hadzi,1951
@ *Sertella beaniiana*
Adriatic
- Claustrofolliculina* Hadzi,1951
- C. clausa* Hadzi,1951 (TM)
Adriatic
- Diafolliculina* Hadzi,1951
- D. rotunda* Hadzi,1951
Helgoland, Adriatic
- D. similis* Hadzi,1951
@ hydroids
Adriatic
- Donsia* Hadzi,1951, non Jankowski,1967
= *Semifolliculina* Dons,1914 (p.p.)
- Echinofolliculina* Dons,1934 (TM) *E. mortensenii* Dons,1934 - New Zealand
- Epifolliculina* Hadzi,1951
- E. diaphana* Hadzi,1951 (TM)
Adriatic
- Eufolliculina* Hadzi,1951
- E. ampulacea* Hadzi,1951
= *Folliculina ampulla* : Lackmann,1910, non (O.F. Müller,1786)
Adriatic
- E. moebiusi* (Kahl,1932) Hadzi,1951
?
- E. latemarginata* (Hadzi,1938) Hadzi,1951
= *Folliculina latemarginata* Hadzi,1938
Adriatic
- Folliculina* Lamarck,1816
- F. ampulla* (O.F. Müller,1786)
= *Vorticella ampulla* O.F. Müller,1786 (TM)
= *Ascobius lentus* Henneguy,1884
Baltic, ?Kieler Bucht, ?Helgoland
- Folliculinopsis* Fauré-Fremiet,1936
= *Alexandrina* Hadzi,1938
- F. annulata* Fauré-Fremiet,1936
= *Alexandrina karagjorgjevici* Hadzi,1937-38
@ crusts of *Lithothamnion* & tubes of *Spirorbis tridentatus*
Gullmarfjord, Adriatic
- F. moebiusi* Hadzi,1951
= *Folliculina ampulla* Möbius,1887, non (O.F. Müller,1786)
?Kieler Bucht
- F. lignicola* Fauré-Fremiet,1936
?
- Halofolliculina* Hadzi,1951
= *Freia* Claparède & Lachmann,1858, non Koch,1850 (Arachnida) = *Freya* : Auctt.
- H. elegans* (Claparède & Lachmann,1859) Hadzi,1951
= *Freia elegans* Claparède & Lachmann,1859
Norway
- H. annulata* (Andrews,1944) Hadzi,1951
= *Parafolliculina annulata* Andrews,1944
Hawaii
- Lagotia* T.S. Wright,1858
= *Angustofolliculina* Silén,1947 (TM) *Folliculina spirorbis* Dons,1912 - Norway
= *Semifolliculina* Dons,1914 (p.p.)

- L. viridis* T.S. Wright,1858 (TM)
@ *Carcinus Liocarinus*, serpulids, etc.
?Sweden, Norway, Britain, W France, ?Adriatic
- L. lutea* (Fauré-Fremiet,1936) Hadzi,1951
?
- L. flava* (Silén,1947) Hadzi,1951
Gullmarfjord
- L. aculeata* (Claparède & Lachmann,1858) Hadzi,1951
?
- L. faurefremieti* Hadzi,1951
?
- L. expansa* (Levinsen,1893) Hadzi,1951
= *Fidellum* (?) *expansum* Levinsen,1893
@ hydroids, mollusc shells, serpulids, bryozoans, etc.
Gullmarfjord, Greenland, Adriatic
- L. gigantea* (Dons,1914) Dons,1934
= *Semifolliculina gigantea* Dons,1914
* according to Hadzi,1951 is this possibly only a form of *L. expansa* (Levinsen)
Norway, Spitsbergen
- L. abyssorum* (Giard,1883) Hadzi,1951
?
- L. stylifer* (T.S. Wright,1862) Hadzi,1951
Britain
- L. coerulea* Hadzi,1951
= *Folliculina elegans* : Fauré-Fremiet,1936, non Claparède & Lachmann,18
?
- L. obstetrica* (T.S. Wright,1862) Hadzi,1951
= *Freya obstetrica* T.S. Wright,1862
Britain
- L. donsi* Hadzi,1951
@ bryozoans, hydroids etc.
Adriatic
- L. minima* Hadzi,1951
@ bryozoans, hydroids etc.
Adriatic
- L. spirorbis* (Dons,1912) Dons,19??
= *Folliculina spirorbis* Dons,1912 = *Angustofolliculina spirorbis* : Silén,1947
@ *Spirorbis spirorbis* bryozoans, etc.
Gullmarfjord, Norway, Spitsbergen, W France, Adriatic, E North America
- Latifolliculina* Hadzi,1951
- L. incolorata* Hadzi,1951 (TM)
= *Parafolliculina amphora* : Fauré-Fremiet,1936, non Dons,1914
Woods Hole aquarium tanks
- Magnifolliculina* Uhlig,1964
* See also *Perifolliculina* Hadzi,1951
- M. alata* Uhlig,1964
German Bight
- M. binatala* Uhlig,1964
German Bight
- Metafolliculina* Dons,1925
- M. andrewsi* Hadzi,1938
Helgoland, Adriatic, E North America
- M. producta* (T.S. Wright,1859) Dons,1934
= *Lagotia producta* T.S. Wright,1859
Britain
- M. longicollis* Hadzi,1938
= *M. producta* : Silén,1947, non (T.S. Wright,1859)
W Sweden, Adriatic
- M. elongata* (S.M. Das,1949) Hadzi,1951
Britain
- M. nordgardi* Dons,1925 (TM)
W Sweden-Trondheimsfjord
- Mirofolliculina* Dons,1927
= *Microfolliculina* : Auctt.
- M. limnoriae* (Giard,1883) Dons,1927
= *Freya limnoriae* Giard,1883 (TM)
@ telson of *Limnoria*
Gullmarfjord, Denmark, Trondheim, W France
- Pachyfolliculina* Hadzi,1951
- P. gunneri* (Dons,1927) Hadzi,1951
= *Folliculina gunneri* Dons,1927 (TM)
@ oostegites of *Limnoria lignorum* & their tunnels & crabs
Gullmarfjord, Trondheim, W France
- Parafolliculina* Dons,1914
- P. amphora* Dons,1914
Gullmarfjord, N Norway, Iceland, W France, Adriatic
- P. violacea* (Giard,1888) Dons,1914
= *Folliculina violacea* Giard,1888 = *Folliculina telesto* Laackmann,1910
Gullmarfjord, S Norway, W France, Adriatic, Woods Hole
- P. labiata* Dons,1942
?Norway
- P. glutinata* Hadzi,1951
Adriatic
- P. americana* Hadzi,1951
= *P. amphora* Andrews,1941, non Dons,1914
@ *Elodaea*
Chesapeake Bay
- Pebrilla* Giard,1888
- P. paguri* Giard,1888 (TM)
?
- Pedifolliculina* Hadzi,1951
- P. arctica* (Dons,1915) Hadzi,1951
= *Pseudofolliculina arctica* Dons,1915 (TM)
Tromsø
- Perifolliculina* Hadzi,1951
- P. roestensis* (Dons,1931) Hadzi,1951
= *Parafolliculina roestensis* Dons,1931 (TM)
* Sensus Uhlig,1964 should this species go to *Magnifolliculina* Uhlig
Lofoten, German Bight
- Planifolliculina* Hadzi,1951
- P. cumbens* Hadzi,1951 (TM)
@ Rhodophyceae, Phaeophyceae, *Posidonia* & Bryozoa
* more common than, and easily confused with *Parafolliculina amphora*
Adriatic
- Platyfolliculina* Hadzi,1938
- P. sahrhageana* Hadzi,1938 (TM)
Kieler Bucht
- P. paguri* Andrews & Reinhard,1943
@ abdomen of *Pagurus pubescens*
?
- Pseudofolliculina* Dons,1914 (TM) *Folliculina melitta* Laackmann,1910 - Antarctic
- Splitofolliculina* Hadzi,1951
- S. adherens* Hadzi,1951
Adriatic
- S. longicollis* Hadzi,1951
Adriatic
- Stentofolliculina* Hadzi,1938
- S. tubicola* Hadzi,1938 (TM)
Adriatic
- Tapetofolliculina* Andrews & Nelson, in Hadzi,1951
- T. portitor* Andrews & Nelson, in Hadzi,1951
@ on turbellarian egg capsules on gills of *Limulus*
E North America
- Valletofolliculina* Andrews,1953
- V. bicornis* Andrews & Reinhard,1953 (TM)
German Bight, California
- CLEVELANDELLIDA* de Puytorac & Grain,1976
- Clevelandellidae* Kidder,1938
- Clevelandella* Kidder,1938
- Nyctotheridae* Amaro,1972
- Nyctotherus* Leidy,1849
- Nyctotheroides* Grassé,1928
- Metanyctotherus* Albaret,1970
- Pronyctotherus* Albaret & Njiné,1976
- Sicuophoridae* Amaro,1972
- Metasicuophora* Albaret,1973
- Parasicuophora* Albaret,1968
- Prosicuophora* de Puytorac & Oktem,1967
- Sicuophora* de Puytorac & Grain,1969
- PLAGIOTOMIDA* Albaret,1974
- Plagiotomidae* Bütschli,1887
- Plagiotoma* Dujardin,1841
- ARMOPHORIDA* Jankowski,1964
- Metopidae* Kahl,1927
- Bothrostoma* Stokes,1887
- Brachonella* Jankowski,1964
- Eometopus* Small & Lynn, in Lee, Hutner & Bovee,1985
- Metopus* Claparède & Lachmann,1858
- M. contortus* (Quennerstedt,1867)
= *Metopides contorta* Quennerstedt,1867
Askö, S Baltic, Öresund, Iscfjord, Halland, W Kattegatt, Limfjorden
- M. contortus* v. *pellitus* Kahl,19??
Kieler Bucht
- M. vestitus* Kahl,19??
Öresund
- M. verrucosus* (Cunha,1???)
Öresund
- M. sigonoides* Claparède & Lachmann,18??
Öresund
- M. brachoni* Fenchel,1968
= *M. vestitus* : Villeneuve-Brachon,1940, non Kahl,19??
?

- M. halophilus** Kahl,19??
Øresund
- M. sigmoides** ?Stein,18??
* fresh water species
Baltic (Wismar)
- Palmarella** Jankowski,1975
- Parametopus** Grolière, de Puytorac & Grain,1980
- Tesnospira** Jankowski,1964
- Caenomorphidae** Poche,1913
- Caenomorpha** Perty,1852
- C. levanderi** Kahl,19??
Askö, Øresund, Isefjord, Kattegatt
- Cirranter** Jankowski,1964
- Ludio** Penard,1922
- PHACODINIIDA** Small & Lynn, in Lee, Hutner & Bovee,1985
- Phacodiniidae** Corliss,1979
- Phacodinium** Prowazek,1900
- Transitella** Gellér,1950
- LICNOPHORIDA** Corliss,1957
- Licnophoridae** Bütschli,1897
- Licnophora** Claparède,1867
- L. auerbachii**
@ ectoparasite on eyes of *Aequipecten opercularis* (& may rarely be found also on *Pecten maximus* & *Chlamys varia*)
Ireland
- L.**, sp. : Fenichel,1965
@ *Thysinia sarsi* (Philippi) (& *Abra nitida* - maybe another species)
Gullmarsfjord
- ODONTOSTOMATIDA** Sawaya,1940
- Epalkellidae** Corliss,1960
- Atopodinium** Kahl,1932
- Epalkella** Corliss,1960
- Saprodnium** Lauterborn,1908
- S. halophilum** Kahl,19??
Askö, Øresund, Kattegatt
- Discomorphellidae** Corliss,1960
- Discomorphella** Corliss,1960
- Mylestomatidae** Kahl, in Doflein & Reichenow,1929
- Mylestoma** Kahl,1928
- M. bipartitum** Gourret & Roeser,19??
Øresund
- PERITROMIDA** Jankowski, in Lee, Hutner & Bovee,1985
* Incertae sedis.
- Peritromidae** von Stein,1867
- Peritromus** von Stein,1862
- P. faurei** Kahl,1932
Kieler Bucht, Øresund, Sylt, Norway, White Sea, Yorkshire, French Atlantic, Mediterranean, Black Sea, E USA, Bermuda
- P. californicus** Kirby,1???
Plymouth
- P. emmae** von Stein,18??
Plymouth
- P. kahli** Villeneuve-Brachon,19??
?
- P. montanus** Kahl,19??
?
- CHOREOTRICHIA** Small & Lynn, in Lee, Hutner & Bovee,1985
- CHOREOTRICHIDA** Small & Lynn, in Lee, Hutner & Bovee,1985
- TINTINNINA** Kofoid & Campbell,1929
- Ascampbelliellidae** Corliss,1964
- Acanthostomella** Jørgensen,1927
- A. elongata** Kofoid & Campbell,1929
Arctic Seas
- A. gracilis** (Brandt,1896)
Greenland
- A. norvegica** (von Daday,1887) (TM)
= *Cyrtocylis norvegica* : Auctt.
Kattegatt, Skagerrak, W Norway, North Sea, Barents Sea, Greenland, English Channel
- Ascampbelliella** Corliss,1960
= *Craterella* Kofoid & Campbell,1929, non Schrammen,1901 (Spongiaria)
- A. obscura** (Brandt,1906)
N North Atlantic
- A. urceolata** (Ostenfeld,1899)
Greenland, N North Atlantic
- Codonellidae** Kent,1882
- Poroecus** Cleve,1902
- P. apiculatus** (Cleve,1900)
N North Atlantic
- P. curtus** Kofoid & Campbell,1929
N North Atlantic
- Tintinnopsis** von Stein,1867
- T. acuminata** von Daday,1887
Baltic, Kattegatt, Skagerrak, Norway, North Sea, Britain, Biscay
- T. amphora** Kofoid & Campbell,1929
W Sweden, S Europe
- T. angusta** Meunier,1910
Arctic Seas
- T. annulata** (Claparède & Lachmann,1858)
= *Tintinnus annulatus* Claparède & Lachmann,1858
Norw. Sea
- T. baltica** Brandt,1896
= *T. vasculum* Meunier,1919
= *T. striigosa* Meunier,1919
* May be a synonym of *Tintinnopsis lobiancoi* von Daday
Baltic, Kattegatt, Bohusl., S North Sea, Southampton, Barents Sea, S Europe
- T. beroidea** von Stein,1867
Kattegatt, Skagerrak, S North Sea, Zuiderzee, Barents Sea, Plymouth, Mediterranean, all N Atlantic
- T. brandtii** (Nordqvist,1890)
= *Codonella brandtii* Nordqvist,1890
Baltic, Kattegatt, Skagerrak, Norway
- T. campanula** (Ehrenberg,1840)
Baltic, Kattegatt, Bohusl., Oslofjord, Skagerrak, SE North Sea, Zuiderzee, Southampton, Barents Sea, Mediterranean
- T. campanula** (Ehrenberg,1840), var. *cincta* (Claparède & Lachmann,1858)
Baltic, Kattegatt, Skagerrak, W Norway, Netherlands, Mediterranean
- T. campanula** (Ehrenberg,1840), var. *buetschlii* von Daday,1887
W Norway, ?Skagerrak, Zuiderzee, Mediterranean
- T. campanula** (Ehrenberg,1840), var. *cyathus* von Daday,1887
Baltic, Norway, North Sea, Engl. Channel, S Europe
- T. cochleata** (Brandt,1906)
Baltic, Kattegatt, ?Skagerrak
- T. compressa** von Daday,1887
Irminger Sea, S Europe
- T. cylindrica** von Daday,1887
Baltic, Norway, Southampton, Plymouth
- T. ecaudata** Kofoid & Campbell,1929
Norway
- T. everta** Kofoid & Campbell,1929
Baltic
- T. fennica** Kofoid & Campbell,1929
Baltic, Kattegatt, ?Skagerrak
- T. fimbriata** Meunier,1919
= *Tintinnopsis ventricosa* Levander,1???
Kieler Bucht, SW North Sea, Zuiderzee, Barents Sea
- T. incurvata** Meunier,1910
Arctic Seas, S Europe
- T. karjacensis** Brandt,1896
= *T. beroidea* von Stein,1867, var. *rotundata* Jørgensen,1899
* May be a synonym of *Tintinnopsis lobiancoi* von Daday
Baltic, Kattegatt, Skagerrak, Bohusl., Norway, Southampton, Barents Sea
- T. lata** Meunier,1910
* May be a synonym of *Tintinnopsis lobiancoi* von Daday
Bohusl., Arctic Seas, Plymouth, S Europe
- T. lobiancoi** von Daday,1887
S North Sea, Southampton
- T. magna** Merkle,1909
?Kattegatt, Skagerrak, Norway
- T. major** Meunier,1910
Baltic, Norway, Arctic Seas
- T. meunieri** Kofoid & Campbell,1929
* Possibly a synonym of *Tintinnopsis fimbriata* Meunier
Baltic, Kattegatt, Skagerrak, North Sea, Arctic Seas
- T. minuta** Wailes,1925
* Possibly a synonym of *Tintinnopsis nana* Lohmann
Limfjorden, S Europe
- T. nana** Lohmann,1908
= *Tintinnopsis fistularis* Meunier,19??
Kieler Bucht, Zuiderzee, S Europe
- T. nitida** Brandt,1896
Baltic, Norway, Barents Sea, Greenland
- T. nucula** (Fol,1884)
Baltic, Kattegatt, Skagerrak, North Sea, Norway
- T. parvula** Jørgensen,1912
Baltic, Kattegatt, Bohusl., SW North Sea, Barents Sea, Greenland, S Europe
- T. patula** Meunier,1910
Arctic Seas
- T. pistillum** Kofoid & Campbell,1929

- * May be a synonym of *Tintinnopsis lobiancoi* von Daday
Baltic, Kattegatt, Skagerrak
- T. rapa* Meunier,1910
Baltic, Norway, Arctic Seas
- T. rotundata* Jørgensen,1899
Baltic, S Europe
- T. sacculus* Brandt,1896
Baltic, Bohusl., Barents Sea, N Greenland
- T. sinuata* Brandt,1896
Norway, Greenland, Arctic Seas
- T. spiralis* Kofoid & Campbell,1929
Arctic Seas
- T. strigosa* Meunier,1919
Baltic, North Sea, W France
- T. subacuta* Jørgensen,1899
* May be a synonym of *Tintinnopsis lobiancoi* von Daday
* a variety of *T. tubulosa* (Levander)?
Skagerrak, W Norway
- T. tubulosa* Levander,1900
* May be a synonym of *Tintinnopsis lobiancoi* von Daday
Baltic, Kattegatt, Skagerrak, SW North Sea, Zuiderzee, W Norway, Barents Sea
- T. tubulosoides* Meunier,1910
* May be a synonym of *Tintinnopsis tubulosa* Levander
Baltic, Arctic Seas, Engl. Channel
- T. turbo* Meunier,1919
* May be a synonym of *Tintinnopsis lobiancoi* von Daday
* Possibly a synonym of *Tintinnopsis lohmanni* Laackmann
Southampton, Barents Sea
- T. undella* Meunier,1910
Barents Sea, S Europe
- T. urniger* (Entz, Sr.,18??) Daday
Plymouth
- T. urnula* Meunier,1910
Arctic Seas
- T. vasculum* Meunier,1919
North Sea, W & S Europe
- T. ventricosoides* Meunier,1910
Arctic Seas
- T. lohmanni* Laackmann,1906
Bohusl., Zuiderzee
- T. elongata* von Daday,1887
Bohusl.
- T. turbinata* Balech,1948
Bohusl.
- T. gracilis* Kofoid & Campbell,1929
Barents Sea
- T. japonica* ?,1???
- Codonella* Haeckel,1873
- C. amphorella* Bielermann,1893
North Sea, Irminger Sea
- C. cratera* (Leidy,1877)
Baltic, Bohusl., Plymouth
- C. elongata* Kofoid & Campbell,1929
W & S Europe
- C. galea* Haeckel,1873
Mediterranean
- C. lagena* (Claparède & Lachmann,1858)
= *Tintinnus lagena* Claparède & Lachmann,1858
Norway, W Europe
- C. nationalis* Brandt,1906
Mediterranean
- C. relicta* Minkiewitsch,1905
* a brackish & fresh water form
Baltic, Kattegatt, Skagerrak
- Codonaria* Kofoid & Campbell,1939
- C. cistellula* (Fol,1884)
Mediterranean
- C. oceanica* (Brandt,1906)
W & S Europe
- Codonellopsidae* Kofoid & Campbell,1929
- Luminella* Kofoid & Campbell,1939
* Incertae sedis
- L. punctata* (Wales,1925)
Mediterranean
- Codonellopsina* Kofoid & Campbell,1929
- Codonellopsis* Jørgensen,1925
- C. bulbulus* (Meunier,1919)
W Europe
- C. contracta* Kofoid & Campbell,1929
Norway, W Europe
- C. caudata* (Brandt,1906)
North Sea, Engl. Channel
- C. inornata* (Brandt,1906)
Greenland
- C. orthoceros* (Haeckel,1873)
Baltic, Mediterranean
- C. pusilla* (Cleve,1900)
= *Codonella pusilla* Cleve,1900
Bohusl., Barents Sea
- C. ovata* Jørgensen,1925
?= *Codonella lagena* (Claparède & Lachmann,1858), var. *ovata* Jørgensen,1899
Skagerrak, W Norway, Barents Sea
- C. pusilla* (Cleve,1900)
North Sea, Arctic Seas, W Europe
- C. lagena* (Claparède & Lachmann,1858)
= *Tintinnus lagena* Claparède & Lachmann,1858
N North Sea
- Stenosemellinae* Campbell & Moore,1954
- Stenosemella* Jørgensen,1925
- S. avellana* (Meunier,1919)
North Sea, W Europe
- S. nivalis* (Meunier,1910)
* Possibly a synonym of *S. nucula* (Fol)
Bohusl., Norway, North Sea, Arctic Seas, Plymouth, W Europe, Mediterranean
- S. oliva* (Meunier,1910)
* Possibly a synonym of *S. nucula* (Fol)
Barents Sea, Plymouth, W Europe
- S. producta* (Meunier,1919)
Bohusl., North Sea, W Europe
- S. steinii* Jørgensen,1912
Baltic, Kattegatt, Skagerrak, Zuiderzee, Barents Sea, Mediterranean
- S. ventricosa* (Claparède & Lachmann,1858) Jørgensen,19??
= *Tintinnus ventricosa* Claparède & Lachmann,1858
Baltic, Skagerrak, SW North Sea, W Norway, Plymouth, Hebrides, Barents Sea, Mediterranean
- S. nucula* (Fol,1884) Laackmann,19??
= *Tintinnopsis nucula* Fol,1884
W Baltic, Zuiderzee
- Cyrtarociliidae* Kofoid & Campbell,1929
- Petalotricha* Kent,1881
- P. ampulla* (Fol,1881) Kent,18??
Engl. Channel, W Mediterranean
- P. major* Jørgensen,1925
Engl. Channel, S Europe
- Cyrtarocylis* Fol,1881
- C. cassis* (Haeckel,1873) Fol,18??
Mediterranean
- C. longa* Kofoid & Campbell,1929
Mediterranean, N North Atlantic
- Dictyocystidae* Haeckel,1873
- Dictyocysta* Ehrenberg,1854
- D. dilatata* Brandt,1906
W & S Europe
- D. duplex* Brandt,1906
W & S Europe
- D. elegans* Ehrenberg,1854
Baltic, Kattegatt, Skagerrak, W Norway, N North Sea, Engl. Channel, W Europe, Mediterranean
- D. grandis* Brandt,1906
Engl. Channel
- D. lepida* Ehrenberg,1854
= *D. templum* Haeckel,18??
N North Sea, W Norway, Engl. Channel, Mediterranean
- D. minor* Jørgensen,1925
North Sea, Mediterranean
- D. mitra* Haeckel,1873
* Possibly a synonym of *Dictyocysta elegans* Ehrenberg
Engl. Channel, ?Mediterranean
- D. nidulus* Kofoid & Campbell,1929
W & S Europe
- D. reticulata* Kofoid & Campbell,1929
W & S Europe
- D. speciosa* Kofoid & Campbell,1929
?Kattegatt, Skagerrak, North Sea, W & S Europe
- Epiplocylidae* Kofoid & Campbell,1939
- Epiplocylis* Jørgensen,1925
- E. acuminata* (von Daday,1887)
North Sea, Engl. Channel, S Europe
- Epiplocyloides* Hada,1938
- E. reticulata* (Ostenfeld & Schmidt,1901)
Engl. Channel, Mediterranean
- Epicancella* Kofoid & Campbell,1929
- Metacylididae* Kofoid & Campbell,1929
- Metacylidinae* Kofoid & Campbell,1929
- Helicostomella* Jørgensen,1925

- H. edentata* (Ehrenberg,1833)
Baltic, North Sea, W & S Europe
- H. fusiformis* (Meunier,1919)
North Sea, W Europe, Arctic Seas
- H. kiliensis* (Laackmann,1906)
Baltic, Norway, North Sea, Mediterranean
- H. subulata* (Ehrenberg,1833)
Baltic, Kattegatt, Limfjorden, Bohusl., Skagerrak, Netherlands, W Norway, Barents Sea, Engl. Channel, W & S Europe
- Metacylis** Jørgensen,1925
- M. annulata* (Meunier,1910)
Arctic Seas
- M. jørgensenii* (Cleve,1902)
?Kattegatt, Skagerrak, North Sea, Norway, W & S Europe
- M. vitreoides* Kofoid & Campbell,1929
Barents Sea, Greenland
- M. mediterranea* (Mereschowsky,18???)
= *Ptychocylis amphorella* Meunier,19??
N Kattegatt, Skagerrak, SW North Sea, Mediterranean
- Coxliellina* Kofoid & Campbell,1929
- Climacocylis* Jørgensen,1925
- C. scalaroides* Kofoid & Campbell,1929
W Europe
- Coxliella* Brandt,1908
* Nomen inquirendum. The species in this genus may be only life cycle stages of species in other genera.
- C. ampla* (Jørgensen,1899) Laackmann,19??
= *Amphorella ampla* Jørgensen,1899
N Kattegatt, Skagerrak, W Norway, North Sea, Arctic Seas, W & S Europe
- C. annulata* (von Daday,1886)
Bohusl., S Europe
- C. calyptra* (Cleve,1899)
Arctic Seas, Greenland
- C. helia* (Claparède & Lachmann,1858)
= *Tintinnus helix* Claparède & Lachmann,1858
= *Tintinnopsis fistularis* Möbius,18??
S & W Baltic, Kattegatt, Bohusl., Skagerrak, W Norway, North Sea, Barents Sea
- C. intermedia* (Laackmann,1907)
Arctic Seas
- C. meunieri* Kofoid & Campbell,1929
Barents Sea
- C. pseudannulata* (Jørgensen,1899)
N Kattegatt, Skagerrak, W Norway, North Sea, Arctic Seas, Greenland, S Europe
- C. tubularis* (Meunier,1910)
Arctic Seas
- Ptychocylididae** Kofoid & Campbell,1929
- Ptychocyclinae** Kofoid & Campbell,1929
- Ptychocylis** Brandt,1896
- P. acuta* Brandt,1896
* May be a synonym of *P. obtusa* Brandt
?Kattegatt, Skagerrak, Norway, North Sea, Arctic Seas, Greenland
- P. arctica* Brandt,1896
* Possibly a synonym of *P. obtusa* Brandt
?Kattegatt, Skagerrak, Norway, North Sea, Greenland, Barents Sea
- P. basiscurvata* Meunier,1910
?Kattegatt, Skagerrak, Norway, North Sea, Arctic Seas
- P. cylindrica* Meunier,1910
* May be a synonym of *P. obtusa* Brandt
Arctic Seas
- P. drygalskii* Brandt,1896
* May be a synonym of *P. obtusa* Brandt
?Kattegatt, Skagerrak, Norway, North Sea, Arctic Seas, Greenland
- P. glacialis* Meunier,1910
* May be a synonym of *P. obtusa* Brandt
Arctic Seas
- P. minor* Jørgensen,1899
W Baltic, Kattegatt, Skagerrak, Bohusl., W & N Norway, NE Scotland, Greenland
- P. obtusa* Brandt,1896
= *P. elongata* Meunier,18??
?Kattegatt, Skagerrak, Norway, North Sea, Greenland, Barents Sea
- P. ostenfeldi* Kofoid & Campbell,1929
N North Atlantic
- P. urnula* (Claparède & Lachmann,1858)
= *Tintinnus urnula* Claparède & Lachmann,1858
Baltic, Bohusl., Skagerrak, N Norway, North Sea, Barents Sea, Engl Channel
- Favellinae** Kofoid & Campbell,1929
- Cymatocylis* Laackmann,1910
- C. keruelensis* Laackmann,1910
Shetland - Faeroe - Iceland area
- Favella* Jørgensen,1925
- F. attingata* Kofoid & Campbell,1929
Norway, S Europe
- F. brevis* Kofoid & Campbell,1929
?Kattegatt, Skagerrak, Mediterranean
- F. ehrenbergii* (Claparède & Lachmann,1858) Jørgensen,1???
= *Tintinnus ehrenbergii* Claparède & Lachmann,1858
?Bothnian Sea, Kattegatt, Bohusl., Skagerrak, Norway, S North Sea, Zuiderzee, Engl. Channel, Mediterranean
- F. ehrenbergii* (Claparède & Lachmann,1858) var.*subannulata* Jørgensen,1899
Skagerrak, W Norway
- F. franciscana* Kofoid & Campbell,19??
Plymouth
- F. helgolandica* (Brandt,1906) Jørgensen,1???
* *Duboscquella tintinnicola* may be found parasitizing this species.
Helgoland, Norway, Engl. Channel, W Europe, Mediterranean
- F. infundibulum* Kofoid & Campbell,1929
Norway
- F. markusovszkyi* (von Daday,1887)
North Sea, S Europe
- F. meunieri* Kofoid & Campbell,1929
North Sea, W Europe
- F. serrata* (Möbius,1887) Jørgensen,1???
W Baltic, Kattegatt, Skagerrak, S & central North Sea, W Norway, Arctic Seas, Engl. Channel, W & S Europe
- Rhabdonellidae** Kofoid & Campbell,1929
- Protorhabdonella* Jørgensen,1925
- Rhabdonella* Brandt,1907
- R. amor* (Cleve,1900)
Greenland, W & S Europe
- R. brandti* Kofoid & Campbell,1929
Greenland
- R. chavesi* Brandt,1907
N North Atlantic, S Europe
- R. conica* Kofoid & Campbell,1929
N North Atlantic, S Europe
- R. hebe* (Cleve,1900)
N North Atlantic, S Europe
- Rhabdonellopsis** Kofoid & Campbell,1929
- R. apophysata* (Cleve,1900)
N North Atlantic
- Tintinnidae** Claparède & Lachmann,1858
- Tintinninae** Claparède & Lachmann,1858
- Tintinnus* Schrank,1803
- T. inquinilum* (O.F. Müller,1776)
@ symbiotic with *Chaetoceros*
Baltic, Kattegatt, Skagerrak, ?German Baltic, Denmark, Norway, Barents Sea, Engl. Channel, W Europe, Mediterranean
- T. obliqua* Claparède & Lachmann,1858
Norway, North Sea, W Europe
- T. vitreus* Brandt,1896
Greenland
- Stenstrupiella* Kofoid & Campbell,1929
- S. steenstrupii* (Claparède & Lachmann,1858)
= *Tintinnus steenstrupii* Claparède & Lachmann,1858
?Kattegatt, Skagerrak, North Sea, W Norway, Engl. Channel, W & S Europe
- Amphoridae** Strand,1928
- = *Amphorella* von Daday,1887, *non* Lowe,1852 (Mollusca), *nec* Léger,1892 (Gregarina)
- A. amphora* (Claparède & Lachmann,1858)
= *Tintinnus amphora* Claparède & Lachmann,1858
North Sea, Norway
- A. infundibulum* (Kofoid & Campbell,1929)
Norway
- A. quadrilineata* (Claparède & Lachmann,1858)
= *Tintinnus quadrilineatus* Claparède & Lachmann,1858
Kieler Bucht, Kattegatt, Skagerrak, N North Sea, W Norway, Barents Sea, W Europe, Mediterranean
- A. borealis* (Hensen,1890)
= *Tintinnus borealis* Hensen,1890
= *Cothurnia borealis* (Hensen,1890) Ostenfeld,1916
= *Cothurnia innata* : Hamburger & von Buddenbrock,1911, *non* *Trichoda innata* O.F. Müller,1786
= *Cothurnia maritima* : Auct., *non* Ehrenberg,18??
@ *Chaetoceros danicum* Cleve
Baltic, Belt Sea, Öresund, Kattegatt, W Norway
- Amphorellopsis* Kofoid & Campbell,1929
- Albatrossiella* Kofoid & Campbell,1929
- A. minutissima* (Meunier,1910)
Greenland, Arctic Seas
- Dadayiella* Kofoid & Campbell,1929
- D. bulbosa* (Brandt,1906)
N North Atlantic
- D. ganymedes* (Entz Sr.,1884)
Mediterranean, N North Atlantic
- D. joergensei* Kofoid & Campbell,1929
Engl. Channel
- Steliidiellina* Kofoid & Campbell,1929
- Ormosella* Kofoid & Campbell,1929
- Brandtiella* Kofoid & Campbell,1929
- Steliidiella* Kofoid & Campbell,1929

- Salpingellinae** Kofoid & Campbell,1939
- Eutintinnus** Kofoid & Campbell,1939
- E. apertus** (Kofoid & Campbell,1929)
W & S Europe
- E. elongatus** (Jørgensen,1925)
N North Atlantic, S Europe
- E. fraknoi** (von Daday,1887)
= *Tintinnus fraknoi* von Daday,1887
Norway, North Sea, W Europe, Mediterranean, N North Atlantic
- E. latus** (Jørgensen,1925)
N North Atlantic, Mediterranean
- E. latusundae** (Entz,1885)
= *Tintinnus latus undae* Entz,1885
Skagerrak, N North Sea, W Norway, Barents Sea, Biscay, N North Atlantic, Mediterranean
- E. macilentus** (Jørgensen,1925)
North Sea, S Europe
- E. pinguis** (Kofoid & Campbell,1929)
N North Atlantic, W Europe
- E. tubulosus** (Ostenfeld,1899)
?Kattegatt, Skagerrak, Norway, N North Atlantic, Britain, W & S Europe
- Daturella** Kofoid & Campbell,1929
- D. gaussi** Kofoid & Campbell,1929
N North Atlantic
- Salpingella** Jørgensen,1925
- S. acuminata** (Claparède & Lachmann,1858) Jørgensen
= *Tintinnus acuminatus* Claparède & Lachmann,1858
W Baltic, Skagerrak, North Sea, W Norway, Barents Sea, Greenland, Engl. Channel, W Europe, Mediterranean
- S. curta** Kofoid & Campbell,1929
Baltic, W Europe
- S. decurtata** Jørgensen,1925
N North Atlantic
- S. gracilis** Kofoid & Campbell,1929
N North Atlantic, W & S Europe
- S. minutissima** Kofoid & Campbell,1929
N North Atlantic, W & S Europe
- S. secata** (Brandt,1896)
Kattegatt - Skagerrak area, Arctic Seas, Greenland, N North Atlantic
- Salpingelloides** Campbell,1942
- S. atiplicata** (Merkle,1909)
Norway
- Salpingacantha** Kofoid & Campbell,1929
- S. undata** (Jørgensen,1899)
Norway, Arctic Seas, Shetlands - >Faeroes - Iceland area
- Tintinnidiidae** Kofoid & Campbell,1929
- Tintinnidium** Kent,1882
- T. mucicola** (Claparède & Lachmann,1858) von Daday
= *Tintinnus mucicola* Claparède & Lachmann,1858
Kieler Bucht, Kattegatt, Skagerrak, W Denmark, Norway, Netherlands
- T. incertum** Brandt,1906
= *T. primitivum* Bush,1925
Zuiderzee
- Lepotintinnus** Jørgensen,1900
- L. pellucidus** (Clevé,1886) Jørgensen,1???
= *Tintinnus pellucidus* Clevé,1886
* Year may be wrong, perhaps 1899?
Belt Sea, Bohusl., W Norway, Barents Sea
- L. bottnicus** (Nordqvist,1890) Jørgensen,1899 (TM)
= *Tintinnus bottnicus* Nordqvist,1890
= *Codonella bottnica* : Levander,1895
all Baltic, W Norway, Barents Sea, Zuiderzee
- Undellidae** Kofoid & Campbell,1929
- Undellinae** Kofoid & Campbell,1929
- Undella** von Daday,1887
- U. hyalina** von Daday,1887
Norway, S Europe
- Ampectella** Kofoid & Campbell,1929
- Undellosinae** Campbell,1942
- Undellosis** Kofoid & Campbell,1929
- Propectellinae** Campbell,1942
- Propectella** Kofoid & Campbell,1929
- P. acuta** (Jørgensen,1925)
N North Atlantic, S Europe
- P. claparedei** (Entz Sr.,1885)
N North Atlantic, W Europe, Mediterranean
- P. ellipsoida** Kofoid & Campbell,1929
N North Atlantic, S Europe
- P. fastigata** (Jørgensen,1925)
= *Undella claparedei* forma *fastigata* Jørgensen,19??
* Spelling? (*P. fastigata* or *fastigiata*?)
N North Atlantic, Mediterranean
- P. subcaudata** (Jørgensen,1925)
N North Atlantic, S Europe
- Xystonellidae** Kofoid & Campbell,1929
- Xystonella** Brandt,1907
- X. acus** (Brandt,1906)
Engl. Channel
- X. lohmanni** (Brandt,1906)
N North Atlantic
- X. treforti** (von Daday,1887)
N North Atlantic, Mediterranean
- Xystonellopsis** Jørgensen,1925
- X. paradoxa** (Clevé,1900)
N North Atlantic
- Parafavella** Kofoid & Campbell,1929
- P. acuminata** (Ehrenberg,1854)
= *P. greenlandica* (Kofoid & Campbell,1929)
Greenland, S Europe
- P. acuta** (Jørgensen,1901)
= *Cyrtarocylis denticulata* var. *typica* Jørgensen,1901 f. *acuta* Jørgensen,1901
Norway, North Sea, Greenland, Barents Sea
- P. calycina** (Jørgensen,1901)
= *Cyrtarocylis edentata* var. *paramundata* Jørgensen,1901
N Norway, North Sea, Arctic Seas
- P. curvata** Kofoid & Campbell,1929
Norway, Arctic Seas
- P. cylindrica** (Jørgensen,1899) Kofoid & Campbell,1929
= *Cyrtarocylis denticulata* Ehrenberg,1899
Baltic, Kattegatt, Skagerrak, North Sea, W & N Norway, Greenland, Arctic Seas
- P. denticulata** (Ehrenberg,1840)
= *Cyrtarocylis denticulata* Ehrenberg,1840
S & W Baltic, Kattegatt, Bohusl., Skagerrak, N & central North Sea, W & N Norway, Barents Sea, Greenland, Engl. Channel, W & S Europe
- P. digitalis** Kofoid & Campbell,1929
Norway, North Sea
- P. dilatata** (Jørgensen,1899)
N Norway, North Sea, Arctic Seas
- P. edentata** (Brandt,1896)
= *Cyrtarocylis edentata* Brandt,1896
North Sea, N Norway, Greenland, Barents Sea
- P. elegans** (Ostenfeld,1899)
North Sea, Arctic Seas
- P. gigantea** (Brandt,1896)
= *Cyrtarocylis gigantea* Brandt,1896
Bohusl., N Norway, North Sea, Arctic Seas, Engl. Channel, W Europe
- P. hemifusus** (Meunier,1910)
Arctic Seas
- P. inflata** Kofoid & Campbell,1929
Arctic Seas
- P. media** (Brandt,1896)
Baltic, Norway, Arctic Seas, Greenland
- P. obtusa** (Aurivillius,1899)
North Sea, Barents Sea
- P. obtusangula** (Ostenfeld,1899)
= *Cyrtarocylis obtusangula* Ostenfeld,1899
Bohusl., N Norway, North Sea, Arctic Seas
- P. paramundata** (Brandt,1906)
= *Cyrtarocylis edentata* var. *paramundata* Brandt,1906
N Norway, Spitsbergen, Greenland, Arctic Seas
- P. robusta** (Jørgensen,1901)
= *Cyrtarocylis denticulata* var. *robusta* Jørgensen,1901
N Norway, North Sea, Greenland, Arctic Seas
- P. rotunda** (Jørgensen,1899)
?Kattegatt, Skagerrak, North Sea, W Norway, Arctic Seas
- P. subedentata** (Jørgensen,1905)
Norway
- P. subtrotunda** (Jørgensen,1899)
?Kattegatt, Skagerrak, North Sea, W Norway, Arctic Seas
- P. subula** Kofoid & Campbell,1929
North Sea, Arctic Seas, W Europe
- P. ventricosa** (Jørgensen,1899)
= *Cyrtarocylis denticulata* var. *cylindrica* f. *ventricosa* Jørgensen,1899
Norway, Arctic Seas
- P. tenuis** (Schulz & Wulff,1929)
W Barents Sea
- Parundella** Jørgensen,1925
- P. aculeata** Jørgensen,1925
Shetlands - Faeroes - Iceland area, W & S Europe
- P. attenuata** Kofoid & Campbell,1929
W & S Europe, N North Atlantic
- P. caudata** (Ostenfeld,1899)
North Sea, Norway, W & S Europe, Shetlands - Faeroes - Iceland area
- P. difficilis** Kofoid & Campbell,1929
W & S Europe
- P. grandis** Kofoid & Campbell,1929

- N North Atlantic ?
P. lachmanni (von Dayad,1887)
 Baltic, Norway
P. lata Ringdal Gaarder,1946
 W Europe
P. lohmanni Jørgensen,1925
 Shetlands - Faeroes - Iceland area, W & S Europe
P. longa Jørgensen,1925
 North Sea, S Europe
P. pellucida (Jørgensen,1899)
 = *Ündella pellucida* Jørgensen,1899
 North Sea, Norway, Barents Sea
STROMBIDINOPSISNA Small & Lynn, in Lee, Hutner & Bovee,1985
Strombidinopsidae Small & Lynn, in Lee, Hutner & Bovee,1985
Strombidinopsis Kent,1881
S. spiniferum (Leegaard,1915) Lynn, Montagnes,Dale, Gilron,Strom,1991
 = *Strombidiitum spiniferum* (Leegaard,1915) Kahl,1932
 = *Strombidiitum spiniferum* Leegaard,1915
 North Sea, Barents Sea (off N Norway; pelagic in neritic surface waters)
STROBILIDIINA Small & Lynn, in Lee, Hutner & Bovee,1985
Strobilidiidae Kahl, in Doflein & Reichenow,1929
S. strobiliidium Schewiakoff,1893
 * Mainly pelagic
S. conicum Kahl,1932
 Øresund
S. caudatum Kahl,1932
 * interstitial
 Kieler Bucht
S. minimum (Gruber,1???)
 * interstitial
 Kieler Bucht
S. epacrum Lynn & Montagnes,1988
 Gulf of Maine
S. multinucleatum Lynn & Montagnes,1988
 Gulf of Maine
S. sphaericum Lynn & Montagnes,1988
 Gulf of Maine
S. spiralis (Leegaard,1915) Lynn & Montagnes,1988
 = *Lohmanniella spiralis* Leegaard,1915
 Tjärnö, Kattegatt, Southampton, Gulf of Maine
S. gyrans (Stokes,1887) Schewiakoff,1892
?
Lohmanniella Leegaard,1915
L. oviformis Leegaard,1915, em. Lynn & Montagnes,1988
 Kattegatt, Limfjorden, Gulf of Maine
L. spiralis ??,1???
 Kattegatt
Sphaerotrichium Wulff,1919
S. elegans (Wulff,1919) Kahl,1932
?
Leegaardellidae Lynn & Montagnes,1988
Leegaardellia Lynn & Montagnes,1988
L. sol Lynn & Montagnes,1988 (TM)
 = *Lohmanniella oviformis* Leegaard,1915 (p.p. - larger form)
 Gulf of Maine
L. ovalis Lynn & Montagnes,1988
 = *Lohmanniella oviformis* Leegaard,1915 (p.p. - medium-sized form)
 Gulf of Maine
OLIGOTRICHIDA Bütschli,1887
Halteriidae Claparède & Lachmann,1858
Halteria Dujardin,1841
H. grandinella (Müller,1???)
?
Meseres Schewiakoff,1893
M. corlissi Petz & Foissner,1992
?
Metastrombidium Fauré-Fremiet,1924
M. sonnifer Jankowski,1980
?
Strombiidiidae Fauré-Fremiet,1970
Cyrtostrombidium Lynn & Gilron,1993
C. longisomum Lynn & Gilron,1993 (TM)
 Jamaica
C. walesi Lynn & Gilron,1993
 Jamaica
Pelagostrombidium Krainer,1991
P. mirabile (Penard,1916) Krainer,1991
?
P. fallax (Zacharias,1896) Krainer,1991
Strombidium Claparède & Lachmann,1859
S. latum Kahl,1932
 S Baltic, Øresund
S. acutum Leegaard,1915
 Plymouth, Gulf of Maine
S. cinctum Kahl,1932
 Kieler Bucht, Sylt, Yorkshire, Plymouth
S. datum Lynn, Montagnes & Small,1988
 Plymouth, Gulf of Maine, Jamaica
S. sauerbreyae (Sauerbrey,1928)Kahl,1932
 W Baltic, Øresund, Sylt, N Russia, French Atlantic, Mediterranean, Black Sea, Mauretania, E USA
S. sulcatum Claparède & Lachmann,1858
 Limfjorden, Plymouth, Gulf of Maine
S. costatum Tucoleesco,1962
?
* possibly a synonym of *S. sulcatum* C. & L.
?
S. lagunula Fauré-Fremiet,19??
 Plymouth
S. wulffi (Wulff,1919) Kahl,1932
 Plymouth, Gulf of Maine, Jamaica
S. kahli Bock,1952
 S Baltic, New Hampshire
S. constrictum (Meunier,1910) Wulff,1919
 Gulf of Maine, Jamaica
S. viride von Stein,1859
?
S. macronucleatum Dragesco,1960
?
S. styliferum (Levander,1???)
?
S. elegans Florentin,1901
?
S. faurei Dragesco,1960
?
S. arenicola Dragesco,1960
?
S. calkinsi Kahl,19??
?
S. typicum (Lancester,1874) Bütschli,1889
?
S. epidemum Lynn, Montagnes & Small,1988
 Gulf of Maine, Jamaica
S. purpureum Kahl,19??
 New Hampshire
S. rhynchum Lynn, Montagnes & Small,1988
 Gulf of Maine
S. tressum Lynn, Montagnes & Small,1988
 Gulf of Maine
S. alveolare Bullington,1940
?
S. opisthostomum Tucoleesco,1962
?
S. conicum (Lohmann,1908) Wulff,1919
=? *Laboea conica* Lohmann,1908
?Kieler Bucht, Southampton
S. oculatum Gruber,1884
?
S. ovale (Leegaard,1915) Kahl,1932
=? *S. oblongum* Leegaard,1915
?
S. tintinnoides Entz,1884
?
S. reticulatum (Leegaard,1915) Busch,1921?
?
S. vestitum (Leegaard,1915) Kahl,1932
 North Sea
S. delicatissimum (Leegaard,1915) Busch,1921
 North Sea, N Atlantic
S. emergens (Leegaard,1915) Kahl,1932
=? *S. sulcatum* : Lynn & al.,1988, non Claparède & Lachmann,1858
?
S. inclinatum Montagnes, Taylor & Lynn,1990
 Jamaica
S. oculatum Gruber,1884
?
S. hadai (Hada,1970) Maeda & Carey,1985
?
S. cornucopiae (Wailes,1929) Kahl,1932
?
S. clavellina Buddenbrook,1922
?
S. grande Levander,1894

- ?
- S. obliquum* Kahl,1932
- ?
- S. compressum* (Leegaard,1915) Kahl,1932
Gulf of Maine
- ?
- S. conicoides* (Leegaard,1915) Kahl,1932
- ?
- S. caudatum* (Fromentel,19???)
- ?
- S. rehwaldi* Petz & Foissner,1992
- ?
- S. bilobum* Lynn & Gilron,1993
Jamaica
- S. maedai* Lynn & Gilron,1993
Jamaica
- S. pollostomum* Lynn & Gilron,1993
Jamaica
- S. sphaericum* Lynn & Gilron,1993
Jamaica
- S. eurystomum* Lynn & Gilron,1993
Jamaica
- S. ioanum* Lynn & Gilron,1993
Jamaica
- Laboea* Lohmann,1908
= *Conocylis* Meunier,1910
- L. strobila* Lohmann,1908
Kieler Bucht, Skagerrak, Plymouth, Jamaica
- Tontonia* Fauré-Fremiet,1914
- T. gracillima* Fauré-Fremiet,1924
Gulf of Maine
- ?
- T. caudata* (Lohmann,1908) Kahl,1932
- ?
- T. appendiculariformis* Fauré-Fremiet,1914 (TM)
- ?
- T. ovalis* (Leegaard,1915) Laval-Peuto & Rassoulzadegan,1987
?
- T. poopsia* Montagnes & Lynn,1988
Maine, USA
- T. simplicidens* Lynn & Gilron,1993
Divided by the authors in two "forms", *octocarya* & *quadricarya*
Jamaica
- STICHOTRICHIA** Small & Lynn, in Lee, Huttner & Bovee,1985
- STICHOTRICHIDA** Fauré-Fremiet,1961
- STICHOTRICHINA** Fauré-Fremiet,1961
- Amphisiellidae* Small & Lynn, in Lee, Huttner & Bovee,1985
- Amphisiella* Gourret & Roeser,1888
- A. milnei* Kahl,19???
- ?
- A. annulata* Kahl,19???
- ?
- A. marioni* Gourret & Roeser,1888 (TM)
off Yucatan peninsula
- A. arenicola* Fernandez-Leborans & Novillo,1992
Mediterranean Spain
- A. ovalis* Fernandez-Leborans & Novillo,1992
Mediterranean Spain
- Balladyna* Kowalewski,1882
- B. euplotes* Dragesco,1950?
- ?
- Eschaneustyla* Stokes,1886
- Kahliella* Corliss,1960
- Onychodromopsis* Stokes,1887
- Onychodromus* von Stein,1859
- Paraurostyla* Borror,1972
- Chaetospiridae* Jankowski, in Lee, Huttner & Bovee,1985
- Chaetospira* Lachmann,1856
- Cladotrichidae* Small & Lynn, in Lee, Huttner & Bovee,1985
- Cladotricha* Gajewskaja,1926
- Engelmanniella* Foissner,1982
- Lamtostyla* Buitkamp,1977
- Perisincirra* Jankowski,1978
- Uroleptooides* Wenzel,1953
- U. kihni* Wenzel,1953
E USA
- Gonostomatidae* Small & Lynn, in Lee, Huttner & Bovee,1985
- Gonostomum* Sterki,1878
- Trachelochaeta* Srámkov-Husek,1954
- Wallackia* Foissner,1977
- Keronidae* Dujardin,1840
- Epiclantes* von Stein,1862
- * Incertae sedis
- E. ambiguus* (O.F. Müller,1786) Bütschli,1889
E & W Baltic, Sylt, White Sea, Yorkshire, Plymouth, French Atlantic, Mediterranean, Black Sea, Mauretania, E USA, Bermuda, Brazil
- Keronopsis* Penard,1922
- * using symbiotic bacteriae
- K. rubra* (Ehrenberg,1838)
Elsinore laboratory salt water system, Yucatan peninsula
- K. monilata* (Kahl,1928)
Sylt, France
- K. flavigans* Kahl,19???
- S Baltic, Plymouth
- K. multistilata* Kahl,19???
- Plymouth
- K. gracilis* Kahl,19???
- Helgoland
- K. pulchra* Kahl,19???
- S Baltic
- K. globulifera* Kahl,19???
- ?
- K. coronata* (Gourret & Roeser,1???)
- ?
- K. ovalis* Kahl,19???
- S Baltic
- K. arenivorus* Dragesco,19???
- ?
- Psilotrichidae** Bütschli,1889
- Caryotricha* Kahl,1932
- Kiitricha* Nozawa,1941
- Psilotricha* von Stein,1859
- Spirofilidae* von Gelei,1929
- Atractos* Vörösváry,1950
- Hypotrichidium* Ilowaisky,1921
- Spiratella* Borror,1972
- Stichotricha* Perty,1849
- S. ?gracilis* Möbius,18???
- Kieler Bucht
- S. secunda* Perty,1849?
- Halland, E USA
- Urostrongylum* Kahl,1932
- U. caudatum* Kahl,1932
W Baltic, Öresund, Isefjord, Sylt, White Sea, Plymouth, Roscoff, Mediterranean, E USA, Bermuda
- Strongyliididae* Fauré-Fremiet,1961
- Strongylidium* Sterki,1878
- S. areniculus* Dragesco,19???
- ?
- UROSTYLINA** Jankowski,1979
- Pseudokeronopsidae* Borror & Wicklow,1983
- Pseudokeronopsis* Borror & Wicklow,1983
- Thigmokeronopsis* Wicklow,1981
- Pseudourostylidae* Jankowski,1979
- Pseudourostyla* Borror,1972
- Irostylidae* Bütschli,1889
- Bakuella* Agamaliev & Alekperov,1976
- B. marina* Agamaliev & Alekperov,1976 (TM)
- ?
- Banyulsella* Dragesco,1953
- B. viridis* Dragesco,1953? (TM)
- ?
- Holosticha* Wrzesniewski,1877
= *Trichotaxis* Stokes,1891 (p.p.)
- H. diademata* (Rees,1884)
W Baltic, Sylt, Helgoland, White Sea, E USA
- H. discocephalus* Kahl,1932
Kieler Bucht, Sylt, White Sea, Plymouth, Roscoff, Mediterranean
- H. arenicola* Kahl,19???
- Kieler Bucht, Öresund

- H. kessleri* Wrzesniowski,1877 (TM)
S Baltic, Öresund, New Hampshire
- H. alveolata* Kahl,19??
Kieler Bucht
- H. extensa* Kahl,19??
Kieler Bucht
- H. multistilata* Kahl,1928
Kieler Bucht
- H. velox* Kahl,19??
= *Trichotaxis velox* Kahl,19??
* Correct genus?
S Baltic
- H. manca* Kahl,19??
?
- H. violacea* Kahl,19??
?
- H. scutellum* (Cohn,1866)
?
- Paraholosticha* Kahl,1932
- P. polychaeta* Borror,1966
New Hampshire
- Paraurostyla* Borror,1972
- P. dispar* (Kahl,19??) Borror,1972
Kieler Bucht
- Urostyla* Ehrenberg,1830 (TM *U. grandis* Ehrenberg,1830 - fresh water)
- U. marina* (Kahl,1932) Borror,19??
Kieler Bucht, New Hampshire
- SPORADOTRICHINA** Fauré-Fremiet,1961
- Oxytrichidae* Ehrenberg,1838
- Ancystropodium* Fauré-Fremiet,1907
- Gastrostyla* Engelmann,1862
- G. pulchra* (Perejaslawzewska,1885)
Sylt, White Sea, Mediterranean, Florida
- Histiculus* Corliss,1960
- H. similis* (Quennerstedt,1867) Corliss,1960?
= *Stylyonchia similis* Quennerstedt,1867
S Baltic, Halland
- Laurentiella* Dragesco & Njiné,1971
- Oxytricha* Bory de St. Vincent,1825
- O. pellionella* (O.F. Müller,17??) Ehrenberg,18??
Plymouth
- O. stenocephala* Borror,1963?
?
- O. discifera* Kahl,19??
Kieler Bucht
- O. marina* Kahl,19??
S Baltic
- O. gibba* (O.F. Müller,1786) von Stein,18??
= *Trichoda gibba* O.F. Müller,1786
Halland
- O. rubra* Ehrenberg,18??
Halland
- O. velox* Quennerstedt,1869
Visby
- O. crassa* Claparède & Lachmann,18??
Norway
- Parastylonychia** Dragesco,1963
- Pleurotricha** von Stein,1859
- P. planensis* Fernandez-Leborans,1984
?Spain
- Stylyonchia* Ehrenberg,1830
- Tachysoma* Stokes,1887
- T. saltans* (Cohn,18??) Borror,19??
= *Actinotricha saltans* Cohn,18??
S Baltic, Halland
- Trachelostylidae* Small & Lynn, in Lee, Hurner & Bovee,1985
- Psammomitra* Borror,1972
- P. brevicauda* (Kahl,19??) Borror,1972?
S Baltic, Helgoland
- P. retractilis* (Claparède & Lachmann,18??) Borror,1972?
Kieler Bucht, Helgoland
- P. radiosa* (Quennerstedt,1867)
= *Mitra radiosa* Quennerstedt,1867
Halland
- Trachelostyla* Kahl,1932
- T. caudata* Kahl,1932
Kieler Bucht, Sylt, Norway, N Russia, French Atlantic, Mediterranean, Black Sea
- T. pediculiformis* (Cohn,1866) Kahl,1932
- E & W Baltic, Öresund, Norway, Sylt, Yorkshire, Plymouth, Mediterranean, Black Sea, New Hampshire, Florida, Brazil
- Urosoma* Kowalewski,1882
- Urosomoida* Hemberger,1981
- RHABDOPHORA** Small,1976
- PROSTOMATEA** Schewiakoff,1896
- PROSTOMATIDA** Schewiakoff,1896
- Holophryidae* Perty,1852
- Bursellopsis* Corliss,1960
- Holophrya* Ehrenberg,1833
- H. marina* Mansfeld,1923
Berliner Aquarium
- H. biconica* Sauerbrey,1928
Kieler Bucht
- H. tarda* Quennerstedt,1869
Visby, Whitstable
- H. vorax* Dragesco,19??
?
- Metacystidae* Kahl,1926
- Metacystis* Cohn,1866
- M. truncata* Cohn,18??
Plymouth
- Pelatractus* Kahl,1930
- Vasicola* Tatem,1869
- PRORODONTIDA** Corliss,1974
- Balanionidae* Small & Lynn, in Lee, Hurner & Bovee,1985
- Balanion* Wulff,1922
- B. somatum* Wulff,1922
Kieler Bucht, North Sea, Barents Sea
- Colepidae* Ehrenberg,1838
- Coleps* Nitzsch,1827
- C. pulcher* Spiegel,1926
Kieler Bucht, Sylt, Helgoland, Plymouth, Roscoff, Mediterranean, N Russia
- C. simili* Kahl,1933
Kieler Bucht, Sylt, Mediterranean, White Sea
- C. spiralis* Noland,1937
Sylt, White Sea, E USA
- C. tessellatus* Kahl,1930
Askö, W Baltic, Kieler Bucht, Öresund, Hiddensee, Sylt, Roscoff, Mediterranean, N Russia, New Hampshire, Cape Cod, Bermuda
- C. remanei* Kahl,19??
S Baltic, Plymouth
- C. crenarius* Bock,1952?
S Baltic
- Nolandia* Small & Lynn, in Lee, Hurner & Bovee,1985
- Plagiopogon* von Stein,1859
- P. loricatus* Kahl,1933
W Baltic, Öresund, Kieler Bucht, Hiddensee, Sylt, White Sea
- Tiarina* Bergh,1881
- T. fusus* (Claparède & Lachmann,18??) Bergh,1881
= *Coleps fusus* (Claparède & Lachmann,18??) (TM - North Sea)
* found chiefly during late summer-late autumn. Contains living zooxanthellae according to Ostenfeld,1916
S Baltic, Belt Sea, Kattegatt, Skagerrak, North Sea, Plymouth
- T. levigata* Kofoid-Campbell
Plymouth
- Placidae* Small & Lynn, in Lee, Hurner & Bovee,1985
- Placus* Cohn,1866
- P. striatus* Cohn,1866
Kieler Bucht, Hiddensee, White Sea, Sylt, Plymouth, Roscoff, Woods Hole
- P. luciae* (Kahl,19??)
Plymouth
- P. socialis* Fabre-Domergue,1889
Baltic, North Sea, Woods Hole
- P. salinus* Dietz,1964
French Mediterranean
- P. buddenbrocki* Sauerbrey,1928
Kieler Bucht, Cape Cod
- Spathidiopsis* Fabre-Domergue,1889
- Plagiocampidae* Kahl,1926
- Plagiocampa* Schewiakoff,1892
- P. margaritata* Kahl,1932
Sylt
- P. posticeconica* Kahl,1932
Kieler Bucht, Sylt

| | |
|---|--|
| <i>P. rouxi</i> Kahl,1932 Kieler Bucht & fresh water | <i>Chilophrya</i> Kahl,1930 |
| <i>P. multiseta</i> Kahl,1932 Helgoland | <i>Enchelys</i> O.F. Müller,1773 |
| <i>P. marina</i> Kahl,1932 Helgoland | <i>E. marina</i> Meunier,1907 * Pelagic Barents Sea |
| <i>P. perlata</i> Kahl,1933 Helgoland | <i>E. tarda</i> Quennerstedt,1869 Baltic (Visby) |
| <i>P. acuminata</i> Kahl,1933 Neuwerk brackish water | <i>E. gracilis</i> Mansfeld,1923 North Sea Aquarium |
| <i>P. incisa</i> Kahl,1933 Kieler Bucht | <i>E. pectinata</i> Kahl,1930 Elbe mouth brackish water |
| <u>Prorodontidae</u> Kent,1881 | <i>E. pterotrachea</i> Collin,1913 @ on gills of <i>Pterotrachea coronata</i> ? |
| <i>Mimeticus</i> Small & Lynn, in Lee, Hutner & Bovee,1985 | <i>Enchelyodon</i> Claparède & Lachmann,1859 |
| <i>Paraspadidium</i> Noland,1937 * Incertae sedis. This probably belongs to another family | <i>E. farctus</i> Claparède & Lachmann,18?? Plymouth |
| <i>P. fuscum</i> (Kahl,1928) Askö, W Baltic, Öresund, Hiddensee, Sylt, Norway, White Sea, Yorkshire, France, Black Sea, Mauretania | <i>E. laevis</i> (Quennerstedt,1867) = <i>Lagynus laevis</i> Quennerstedt,1867 Halland |
| <i>P. obliquum</i> Dragesco,19?? ? | <i>E. sulcatus</i> Kahl,1930 Oldesloe |
| <i>Prorodon</i> Ehrenberg,1833 | <i>E. vacuolatus</i> Dragesco,19?? ? |
| <i>P. discolor</i> (Ehrenberg,18??) Askö, W Baltic, Öresund, White Sea, Sylt, Mediterranean, Kamerun | <i>E. fascinucleatus</i> Kahl,1933 Sylt |
| <i>P. marinus</i> Claparède & Lachmann,1958? Sylt, White Sea, France, Black Sea | <i>E. trepida</i> (Kahl,1928) = <i>Trachelocerca trepida</i> Kahl,1928 New Hampshire |
| <i>P. teres</i> Ehrenberg,1838 Schlei, Hiddensee, Sylt, White Sea, Roscoff, E USA | <i>Homalozoon</i> Stokes,1890 |
| <i>P.</i> sp. : Hartwig,1973 Jordsand | <i>H. caudatum</i> Kahl,19?? Öresund |
| <i>P. morganii</i> Kahl,1930 Kieler Bucht, Öresund, Elbe mouth, Oldesloe, Plymouth | <i>Nannophrya</i> Kahl,1933 |
| <i>P. ovum</i> Ehrenberg,1838 Kieler Bucht, Oldesloe, Sylt | <i>N. truncata</i> Kahl,1933 Helgoland |
| <i>P. arenarius</i> Dragesco,19?? ? | <u>Lacrymariidae</u> de Fromental,1876 |
| <i>P. luteus</i> Kahl,1930 Oldesloe | <i>Lacrymaria</i> Bory de St. Vincent,1826 |
| <i>P. elegans</i> Kahl,1928 Oldesloe | <i>L. caudata</i> Kahl,1933 Kieler Bucht, Norway, White Sea, Sylt, French Atlantic, Mediterranean, Black Sea |
| <i>P. karianus</i> Meunier,1907 * pelagic Barents Sea | <i>L. marina</i> Kahl,1933 = <i>L. olor</i> (O.F. Müller,17??) var <i>marina</i> Kahl,1933 * Fenchel, 1968 gives Dragesco, 1963 as author Kieler Bucht, Öresund, Barents Sea, White Sea, Sylt, Roscoff, Mediterranean |
| <i>P. dubius</i> Kahl,1930 Helgoland | <i>L. acuta</i> Kahl,1933 Kieler Bucht |
| <i>P. binucleatus</i> (von Buddenbrock,1920) Berliner Aquarium | <i>L. pupula</i> (O.F. Müller,17??) Plymouth |
| <i>P. opalescens</i> Kahl,1928 Sylt, Oldesloe | <i>L. versatilis</i> (O.F. Müller,17??) Quennerstedt,1867 = <i>Trichoda versatilis</i> O.F. Müller,17?? Halland, Cuxhaven |
| <i>P. moebiusi</i> Kahl,1930 Kieler Bucht, Sylt | <i>L. coronata</i> Claparède & Lachmann,1858 S Baltic, White Sea, Yorkshire, W France, Mediterranean, Black Sea, W Africa |
| <i>P. mimeticus</i> Kahl,1930 Oldesloe, Kaiser-Wilhelm-Kanal, Holtenau | <i>L. delamarei</i> Dragesco,19?? Öresund |
| <i>Pseudoprorodon</i> Blochmann,1886 * Sensu Curds, 1982 is the generic name erected by Kahl, 1930, although Kahl attributed it to Blochmann, 1886 | <i>L. trichocystis</i> Dragesco,19?? ? |
| <i>P. arenicola</i> Kahl,1933 Kieler Bucht, Öresund, Sylt, Norway, Barents Sea, Murmansk, White Sea, Yorkshire, French Atlantic, Mediterranean | <i>L. lagunula</i> Claparède & Lachmann,18?? Halland |
| <i>P. halophilus</i> Kahl,1930 Oldesloe | <i>L. affinis</i> Bock,1952? S Baltic |
| <i>P. incisus</i> Bock,1952? S Baltic | <i>L. cohnii</i> Kent,1881 S Baltic, White Sea, Yorkshire, Whitstable, Caspian Sea |
| <i>P. mononucleatus</i> Bock,1952? S Baltic | <i>L. salinarum</i> Kahl,19?? S Baltic |
| <u>Urotrichidae</u> Small & Lynn, in Lee, Hutner & Bovee,1985 | <i>L. olor</i> (O.F. Müller,17???) Whitstable |
| <i>Patschia</i> Small & Lynn, in Lee, Hutner & Bovee,1985 | <u>Mesodiniidae</u> Jankowski, in Lee, Hutner & Bovee,1985 |
| <i>Rhagadostoma</i> Kahl,1926 | <i>Askenasia</i> Blochmann,1895 |
| <i>Urotricha</i> Claparède & Lachmann,1859 | <i>A. stellaris</i> ??,1??? |
| LITOSTOMATEA Small & Lynn,1981 | <i>Mesodinium</i> von Stein,1863 |
| HAPTORIA Corliss,1974 | <i>M. pupula</i> Kahl,1933 W Baltic, Öresund, Hiddensee, Sylt, White Sea, Yorkshire, French Atlantic, Mediterranean, Ivory Coast, E USA |
| <u>HAPTORIDA</u> Corliss,1974 | <i>M. pulex</i> (Claparède & Lachmann,1858) von Stein,1863 Kieler Bucht, Hiddensee, Sylt, White Sea, Plymouth, Roscoff, Black Sea, Florida |
| <u>Didiniidae</u> Poche,1913 | <i>M. cinctum</i> Calkins,1??? |
| <i>Didinium</i> von Stein,1859 | <i>Myrionecta</i> Jankowski,1976 |
| <i>D. nasutum</i> (O.F. Müller,17???) von Stein,1859 Kieler Bucht, Öresund | <i>M. rubrum</i> (Lohmann,1908) Jankowski,1976 = <i>Halteria rubra</i> Lohmann,1908 |
| <i>D. balbianii</i> Fabre-Domergue,18?? Kieler Bucht | = <i>Mesodinium rubrum</i> (Lohmann,1908) Hamburger & von Buddenbrock,1911 = <i>Cyclotrichium meunieri</i> Powers,1932 |
| <i>Monodinium</i> Fabre-Domergue,1888 | |
| <u>Enchelydidae</u> Ehrenberg,1838 | |

- Baltic, Öresund, Isefjord, Kattegatt, Plymouth, Iceland, Black Sea, Canada, Maine, Florida
Rhabdoaskenasia Krainer & Foissner, 1990
- R.** sp.: Pilling & al., 1992
 Plymouth
- Spathidiidae** Kahl, in Doflein & Reichenow, 1929
- Bryophyllum** Kahl, 1931
- Myriokaryon** Jankowskii, 1973
- Spathidium** Dujardin, 1841
- S. procerum** Kahl, 1930
 Yorkshire, Plymouth
- Trachelophyllidae** Kent, 1886
- Chaenea** Quennerstedt, 1867
- C. vorax** Quennerstedt, 1867 (TM)
 Baltic, Halland, North Sea, Plymouth, Algeria
- C. minor** Kahl, 19??
 Plymouth
- C. robusta** Kahl, 1930
 Kieler Bucht, Oldesloe
- C. simulans** Kahl, 1930
 Elbe mouth
- C. teres** (Dujardin, 1841)
 = *Trachelius teres* Dujardin, 1841
 Oldesloe brackish water & fresh water
- C. gigas** Kahl, 1933
 Kieler Bucht, Helgoland
- Lagynophrya** Kahl, 1927
- L. costata** Kahl, 1933
 Oldesloe
- L. contractilis** Kahl, 1928
 Oldesloe
- L. halophila** Kahl, 1932
 Kiel
- Trachelophyllum** Claparède & Lachmann, 1859
- T. brachypharynx** Levander, 1894
 Finland, Kieler Bucht in weak brackish water
- PLEUROSTOMATIDA** Schewiakoff, 1896
- Amphileptidae** Bütschli, 1889
- Amphileptus** Ehrenberg, 1830
- Heminotus** Kahl, 1933
- H. caudatus** Kahl, 1933 (TM)
 S Baltic
- Hemiophrys** Wrzesniowski, 1870
 * Possibly a synonym of *Amphilectus* Ehrenberg
- H. rotunda** Kahl, 1930
 Sylt, Genfer See
- H. baltica** Fenchel, 1965
 @ on gills, peraeopods & sideplates of *Gammarus* spp.
 Askö, Öresund, Isefjord
- H. filum** Gruber, 1???
- H. marina** Kahl, 19??
 Kieler Bucht
- Litonotus** Wrzesniowski, 1870
- L. lamella** (Ehrenberg, 1838)
 S Baltic, Öresund
- L. binucleatus** Kahl, 1933
 = *L. pictus* var. *binucleatus* Kahl, 1933
 Kieler Bucht, Öresund, Norway, Barents Sea, Murmansk, White Sea, Sylt
- L. cygnus** (O.F. Müller, 1776?)
 Kieler Bucht, Sylt, Plymouth, Black Sea
- L. fasciola** (Ehrenberg, 18??) Wrzesniowski, 1870?
 * not O.F. Müller the author?
 S Baltic, Öresund, Plymouth
- L. duplostriatus** (Maupas, 1???)
 S Baltic
- Loxophyllum** Dujardin, 1841
- L. helus** (Stokes, 1884)
 Kieler Bucht, Hiddensee, Sylt, White Sea, Roscoff, Mediterranean, Black Sea
- L. helus** v. **multiverrucosum** Kahl, 19??
 Kieler Bucht
- L. helus** v. **rotundatum** Kahl, 19??
 Kieler Bucht
- L. levigatum** Sauerbrey, 1928
 Kieler Bucht, Sylt, Barents Sea, White Sea, Roscoff, Black Sea
- L. vermiforme** Sauerbrey, 1928
 Kieler Bucht, Norway, White Sea, Sylt, Roscoff
- L. variabilis** Dragesco, 1954
 Öresund, White Sea, Yorkshire, W France, Mediterranean
- L. rostratum** Cohn, 18??
 = *meleagris* Fresenius, 18??
 Halland
- L. setigerum** Quennerstedt, 1867
 S Baltic, Halland
- L. setigerum** Quennerstedt, 1867 var. *fibrillatum* Dragesco, 1960
 Sylt, Yorkshire, Roscoff
- L. trichocystum** Hartwig, 1973
 Sylt
- L. uninucleatum** Kahl, 19??
 Plymouth
- L. fasciolatum** Kahl, 19??
 S Baltic
- L. multiplicatum** Kahl, 1932
 S Baltic
- L. perihoplophorum** von Buddenbrock, 19??
 Helgoland
- L. serratum** Kahl, 19??
 Kieler Bucht
- L. simplex** Kahl, 1933
 Kieler Bucht
- L. undulatum** Sauerbrey, 19??
 S Baltic
- L. verrucosum** (Stokes, 1???)
 S Baltic
- L. chaetonotum** Borror, 1965
 New Hampshire
- PHARYNGOPHORIDA** Small & Lynn, in Lee, Hutner & Bovee, 1985
- Actinobolinidae** Kahl, 1930
- Actinobolina** Strand, 1928
- Helicoprorodontidae** Jankowski, in Lee, Hutner & Bovee, 1985
- Helicoprorodon** Fauré-Fremiet, 1950
- H. gigas** (Kahl, 1933)
 Kieler Bucht, Öresund, Sylt, Roscoff, Black Sea, White Sea
- H. minutus** Bock, 1952
 Kieler Bucht, Öresund, Sylt, Roscoff, Black Sea, White Sea
- H. multinucleatum** Dragesco, 1960
 White Sea, French Atlantic, Bermuda, Brazil
- H. barbatus** ??, 19??
 ?
- H. maximus** ?Dragesco, 19??
 France
- Tracheliidae** Ehrenberg, 1838
- Dileptus** Dujardin, 1840
- D. marinus** Kahl, 19??
 S Baltic
- D. estuarinus** Dragesco, 19??
 Öresund
- D. anser** (O.F. Müller, 1786)
 = *Vibrio anser* O.F. Müller, 1786
 Visby
- Paradileptus** Wenrich, 1929
- Teuthophrys** Chatton & Beauchamp, 1923
- Trachelius** Schrank, 1803
- T. gutta** Sarhage, 1???
- Kieler Bucht**
- Acropistium** Perty, 1852
 * genus incertae sedis
- TRICHOSTOMATIA** Bütschli, 1889
- VESTIBULIFERIDA** de Puytorac & al., 1974
- Balantidiidae** Reichenow, in Doflein & Reichenow, 1929
- Balantium** Claparède & Lachmann, 1858
- Isotrichidae** Bütschli, 1889
- Dasytricha** Schuberg, 1888
- Isotricha** von Stein, 1859
- Paraisotrichidae** da Cunha, 1917
- Paraisotricha** Fiorentini, 1890
- Coelosomidae** Corliss, 1961
 * not included in Small & Lynn, 1965
- Coelosomides** Strand, 1928
- C. marina** Anigstein, 1912
 France, Black Sea, Caspian Sea, Bermuda, Brazil
- C. tessieri** Dragesco, 1954
 ?
- Conchostoma** Fauré-Fremiet, 1963

- C. longissimum* Fauré-Fremiet,1963 (TM)
Öresund, Yorkshire, Concarneau
Epimecophrya Kahl,1933
* incertae sedis
- E. cylindrica* Kahl,1933 (TM)
Helgoland
- ENTODINOMORPHIDA Reichenow, in Doflein & Reichenow,1929
- ARCHISTOMATINA de Puytorac & al.,1974
- Buetschliidae Poche,1913
- Alloiozona* Hsiung,1930
- Didesmis* Fiorentini,1890
- Polymorphella* Corliss,1960
- BLEPHAROCORYTHINA Wolska,1971
- Blepharocorythidae Hsiung,1929
- Blepharocorys* Bundle,1895
- Circodinium* Wolska,1971
- Ochoterenaia* Chavarria,1933
- Raabena* Wolska,1967
- ENTODINOMORPHINA Reichenow, in Doflein & Reichenow,1929
- Cycloposthiidae Poche,1913
- Cycloposthium* Bundle,1895
- Triplumaria* Hoare,1937
- Ophryoscolecidae von Stein,1859
- Entodinium* von Stein,1859
- Enoploplastron* Kofoid & MacLennan,1932
- Eremoplastron* Kofoid & MacLennan,1932
- Eudiplodinium* Dogiel,1927
- Ophryoscolex* von Stein,1859
- Ostracodinium* Dogiel,1927
- Spirodiniidae Strelkow,1939
- Cochliatoxum* Gassovsky,1919
- CYRTOPHORA** Small,1976
- PHYLLOPHARYNGEA** de Puytorac & al.,1974
- Silenella* Fenchel,1965
* Systematic position very unclear. May possibly be a larval suctorian
- S. ovoidea* Fenchel,1965 (TM - Askö)
@ on gills & peraeopods of *Gammarus duebeni*
Askö
- PHYLLOPHARYNGIA** de Puytorac & al.,1974
- CYRTOPHORIDA Fauré-Fremiet, in Corliss,1956
- CHLAMYDODONTINA Deroux,1976
- Chilodonellidae Deroux,1970
- Alinostoma* Jankowski, in Lee, Hutner & Bovee,1985
- Chilodonella* Strand,1928
- C. uncinata* (Ehrenberg,18??)
Plymouth
- C. calkinsi* Kahl,19??
?
- C. psammophila* Dragesco,19??
?
- C. cyprini* (Moroff,1902) Strand,1928
@ freshwater & estuarine fish
E Baltic
- Phascolodon* von Stein,1859
- Thigmogaster* Deroux,1976
- Trithigmostoma* Jankowski,1967
- Chitonellidae Small & Lynn, in Lee, Hutner & Bovee,1985
- Chitonella* Small & Lynn, in Lee, Hutner & Bovee,1985
- Chlamydotodontidae von Stein,1859
- Chlamydotodon* Ehrenberg,1835
- C. triquetrus* (O.F. Müller,1776)
W Baltic, Sylt, Barents Sea, Murmansk, White Sea, Yorkshire, Plymouth, France, E USA
- C. obliquus* Kahl,19??
Öresund
- C. mnemosyne* Ehrenberg,18??
Öresund, Halland
- C. cyclops* Entz,18??
?
- Cyrtophoron* Deroux,1975
- Lynchellidae* Jankowski,1968
- Atopochilodon* Kahl,1933
- A. arenifer* Kahl,19??
Helgoland
- Chlamydonella* Deroux,1970
- Coeloperix* Deroux,1976
- Gastronauta* Engelmann,1875
- Lynchella* Kahl,1933
- L. gradata* Kahl,19??
Kieler Bucht, Öresund
- L. aspidisciformis* Kahl,19??
Helgoland
- DYSTERIINA Deroux,1976
- Dysteriidae Claparède & Lachmann,1858
- Dysteria* Huxley,1857
- D. aculeata* Claparède & Lachmann,18??
Plymouth
- D. monostyla* (Ehrenberg,18??) von Stein,18??
?= *Euplates monostylus* Ehrenberg,18??
Halland, Plymouth, New Hampshire
- D. marina* Gourret & Roeser,19??
New Hampshire
- Hartmannulopsis* Deroux & Dragesco,1968
- Microdysteria Kahl,1933
- M. aplanata* Kahl,19??
Kieler Bucht
- Trochilia* DuJardin,1841
- T. fluvialis* Smith,1???
Plymouth
- T. salina* Entz,18??
Plymouth
- T. sp. : Fenchel,1965*
@ between gills and sideplates of *Gammarus* spp.
Öresund, Isefjord
- Hartmannulidae Poche,1913
- Aegyriana* Deroux,1975
- Brooklynella* Lom & Nigrelli,1970
- B. hostilis* Lom & Nigrelli,1970
@ gills of various fish hosts (typically in aquaria)
USA
- Hartmannula* Poche,1913
- Microxysma Deroux,1977
- Orthotrichilia* Deroux,1977
- Trochilioides* Kahl,1931
- T. trivialis* Fenchel,1965
@ *Gammarus* spp.
Askö, Öresund, Isefjord
- T. recta* Kahl,19??
Askö, Öresund
- T. oculata* Kahl,19??
Kieler Bucht
- T. sp. a : Fenchel,1965*
@ on pleopods of *Gammarus duebeni*
Askö rock pools
- Trichopodiella* Corliss,1960
- Plesiotrichopidae Deroux,1976
- Atelepithites* Deroux,1976
- Pithites* Deroux & Dragesco,1968
- Plesiotrichopus* Fauré-Fremiet,1965
- Trochochilodon* Deroux,1976
- HYPOCOMATINA Deroux,1976
- Hypocomidae Bütschli,1889
- Crateristoma* Jankowski,1967
- Hypocoma* Grüber,1884
- H. parasitica* Grüber,1884?
@ preying on colonial peritrichs on *Gammarus oceanicus* Segerstråle; Mediterranean *Zoothamnium*
Askö, Öresund, Isefjord, Mediterranean
- H. zoothamni* (Plate,1888)
= *Acinetoides zoothamni* Plate,1888
@ *Zoothamnion*
? Britain
- H. acinetarum* Collin,1907
@ *Ephelota gemmipara*, *Acineta papillifera* & other free-living suctarians
? Britain
- H. ascidiarum* Collin,1912

- @ e.g. *Ascidella aspersa* & *Botryllus*
?France
- Parahypocoma** Chatton & Lwoff,1939
- P. collini** ?Chatton & Lwoff,1939?
@ *Ciona intestinalis*, *Ascidia mentula* & *Clavelina lepadiformis*
?France
- P. rhamphisokarya** ??Burreson,1973??
@ e.g. *Ascidia callosa*
Pacific
- Rhynchocoma** Jankowski,1975
= *Heterocoma* Chatton & Lwoff,1939 (homonym)
- R. hyperparasitica** (?Chatton & Lwoff,1939?)
= *Heterocoma hyperparasitica* ?Chatton & Lwoff,1939? (TM)
@ *Trichophrya salparum* from *Salpa* sp.
?France
- RHYNCHODIIDA** Chatton & Lwoff,1939
- RHYNCHODINA** Chatton & Lwoff,1939
- Sphenophryidae** Chatton & Lwoff,1921
- Gargarius** Chatton & Lwoff,1934
- G. gargarus** Chatton & Lwoff,1934
= *Rhynchophrya cristallina* Raabe,1935
@ *Mytilus edulis* L.
Gdynia, S France
- Pelecyophrya** Chatton & Lwoff,1922
- P. tapetis** Chatton & Lwoff,1922
@ *Venerupis aureus*, non other *Venerupis* or *Tapes* species
Roscoff
- Sphenophryra** Chatton & Lwoff,1921
- S. dosiniae** (Chatton & Lwoff,1921)
= *S. myae* Mjassnikova,1930
@ *Dosinia exoleta* (L.), *Timoclea ovata* (Pennant), *Corbula gibba* (Oliv), *Spisula solidissima* Dillwyn, *Mya truncata* L. & *Mya arenaria* L.
Askö, Gdynia, Öresund, Gullmarfjord, White Sea, Roscoff, E USA
- S. cardii** Chatton & Lwoff,1950
@ *Cerastoderma edule* (L.) & *Cerastoderma glaucum*
Gdynia, Öresund, Roscoff
- ANCISTROCOMINA** Chatton & Lwoff,1939
- Ancistrocomidae** Chatton & Lwoff,1939
- Ancistrocoma** Chatton & Lwoff,1926
- A. myae** (Kofoid & Busch,1936)
= *Parachaenia myae* Kofoid & Busch,1936
@ *Mya arenaria* L.
Öresund, Gullmarfjord, Woods Hole, California, non Baltic
- A. thorsoni** Fenchel,1965
@ *Abra nitida* (Müller)
Gullmarfjord
- A. pelseneeri** Chatton & Lwoff,1926
@ *Barnea candida*
Wimereux
- A. pholadis** ?Chatton & Lwoff,1926?
@ *Zirfaea crispata*
Wimereux
- A. sp.** : Fenchel,1965
@ *Venerupis aurea*
Gullmarfjord
- Anisocomides** Chatton & Lwoff,1950
- A. zyprhaeae** (?Chatton & Lwoff,1926?)
= *Hypocomides zyprhaeae* ?Chatton & Lwoff,1926?
@ *Zirfaea crispata*
Wimereux
- Colligocineta** Kozloff,1965
- Crebricoma** Kozloff,1946
- C. carinata** (Raabe,1934)
= *Hypocomina carinata* Raabe,1934
@ *Mytilus edulis* L.
Askö, Gdynia, Öresund, Gullmarfjord
- C. kozloffi** ??,19??
@ *Mytilus edulis*
California
- Goniocoma** Chatton & Lwoff,1950
- G. macomae** (Chatton & Lwoff,1926)
= *Hypocomella macomae* Chatton & Lwoff,1926
@ *Macoma balthica* (L.) & *Abra alba* (Wood)
Gullmarfjord, non Askö, nec Öresund
- Heterocinetopsis** Jarocki,1935
- Holocoma** Chatton & Lwoff,1950
- H. primigenius** Chatton & Lwoff,1950
@ *Macoma balthica* (L.)
Askö, non Öresund, ?France
- Hypocomagalma** Jarocki & Raabe,1932
- H. dreissenae** Jarocki & Raabe,1932
@ *Dreissena polymorpha* (Pallas) (in fresh & brackish water)
Baltic & Danish rivers & lakes
- Hypocomatidium** Jarocki & Raabe,1932
- H. jarockii** ?Chatton & Lwoff,1950?
- @ *Nucula nucleus*
Banyuls-sur-mer
- Hypocomella** Chatton & Lwoff,1922
- H. raabei** Chatton & Lwoff,1950
@ *Cerastoderma edule* (L.), *Cerastoderma glaucum* & *Parvicardium exiguum*
Askö, Gdynia, Öresund, Roscoff
- H. cardii** Chatton & Lwoff,1922
@ *Cerastoderma edule* (L.)
Roscoff
- Hypocomides** Chatton & Lwoff,1922
- H. modiolariae** Chatton & Lwoff,1922?
@ *Musculus marmoratus* (Forbes)
Roscoff
- H. elsinora** Fenchel,1965
= *H. modiolariae* : Fenchel,1964, non Chatton & Lwoff,1922?
@ *Musculus niger* (Gray)
Öresund
- H. musculus** Fenchel,1965
@ *Musculus discors* (L.)
Gullmarfjord
- H. astartae** Fenchel,1965
@ *Astarte montagui* (Dillwyn)
Gullmarfjord
- H. hiatellae** Fenchel,1965
@ *Hiatella arctica* (L.) &/or *H. striata* (Fleurieu)
Gullmarfjord
- Hypocomidium** Raabe,1938
- H. fabius** ?Raabe,1938?
@ *Cerastoderma glaucum*
Poland
- Ignotocoma** Kozloff,1961
- Insignicoma** Kozloff,1946
- Isocomides** Chatton & Lwoff,1950
- I. mytili** (Chatton & Lwoff,1922)
= *Hypocomides mytili* Chatton & Lwoff,1922 (TM)
Askö, Öresund, Gullmarfjord, Roscoff, Sète
- Raabella** Chatton & Lwoff,1950
- R. helensis** Chatton & Lwoff,1950
@ *Mytilus edulis* L.
Askö, Gdynia, Öresund, Gullmarfjord
- CHONOTRICHIA** Wallengren,1895
- EXOGEMMIDA** Jankowski,1972
- Chilodochonidae** Wallengren,1895
- Chilodochona** Wallengren,1895
- Vasichona** Jankowski,1972
- Filichonidae** Jankowski,1973
- Aurichona** Jankowski,1973
- Filichona** Jankowski,1973
- Heliochonidae** Jankowski,1972
- Heliochona** Plate,1888
- H. sessilis** Plate,1888
@ gills of *Gammarus locusta* L., *Gammarus oceanicus* Segerstråle, *G. salinus* Spooner
Kieler Bucht, Öresund, Isefjord, non Askö, North Sea
- H. scheuteni** (von Stein,1854)
@ anterior side of pleopod rami of *Gammarus* spp.
Askö, Kieler Bucht, Öresund, Isefjord, Gullmarfjord, Netherlands
- Heterochona** Jankowski,1972
- Lobochonidae** Jankowski,1967
- Eleutherochona** Jankowski,1971
- Lobochona** Dons,1940
- L. limnoriae** Dons,1940 (TM*)
@ on uropod hairs of *Limnoria lignorum*
Trondheim
- Oenophorachona** Matsudo & Mohr,1968
- Toxochona** Jankowski,1972
- Phyllochonidae** Jankowski,1972
- Phyllochona** Jankowski,1972
- Spirochonidae** von Stein,1854
- Cavichona** Jankowski,1973
- Serpentichona** Jankowski,1973
- Spirochona** von Stein,1852
- CRYPTOGEMMIDA** Jankowski,1975
- Actinichonidae** Jankowski,1973

- Actinichona** Jankowski,1973
Carinichona Jankowski,1973
Cristichona Jankowski,1973
Rhizochona Jankowski,1973
Echinichonidae Jankowski,1973
Coronochona Jankowski,1973
Echinichona Jankowski,1973
Eurychona Jankowski,1973
Inversochonidae Jankowski,1973
Ceratochona Jankowski,1973
Chonosaurus Jankowski,1973
Inversochona Jankowski,1973
Kentrochona Rompel,1894
Pleochona Jankowski,1973
Isochonodae Jankowski,1973
Isochona Jankowski,1973
Trichochona Mohr,1948
Stylochonidae Mohr,1948
Armichona Jankowski,1973
Dentichona Jankowski,1973
Eriochona Jankowski,1973
Flectichona Jankowski,1973
Oxychonina Corliss,1979
Spinichona Jankowski,1973
Stylochona Kent,1881
SUCTORIA Claparède & Lachmann,1858
EXOGENIDA Collin,1912
Ephelotidae Kent,1882
Ephelota Wright,1858
E. gemmipara Hertw.,1876
@ different algae & animals
Kieler Bucht, ?Brittan
E. mammilata Dons,1918
@ *Diphasia abietina*
Tromsø
E. cothurnata Dons,1918
@ Phaeophyta, Rhodophyta & hydroids
Tromsø
Podocathus Kent,1882
P. excavatus Dons,1938
Bodø
P. diadema Kent,1882?
?
P. cidaris (Kent,18??)
?
Ophryodendridae von Stein,1867
Loricodendron Jankowski,1973
Ophryodendron Claparède & Lachmann,1859
O. multiramosum Wenzel,1961
@ possibly feeding on content of Sponge cells
?Mediterranean
O. trinacrium Gruber,18??
Greenland
Parapodophryidae Jankowski,1973
Parapodophrya Kahl,1931
Podophryidae Haeckel,1866
Corynophrya Kahl,1934
Kystopus Jankowski,1967
Podophrya Ehrenberg,1838
P. tortuosa Dons,1918
@ small Phaeophyta
Tromsø
P. fixa (O.F. Müller,1786) Ehrenberg,18??
= *Trichoda fixa* O.F. Müller,1786
Halland
?P. patula Claparède & Lachmann,18??
Greenland
Sphaerophrya Claparède & Lachmann,1859
Rhabdophryidae Jankowski,1970
Dendrosomides Collin,1906
D. truncata Dons,1918
@ Phaeophyta & hydroids
Tromsø
Trophogemma Jankowski,1970
Spelaeophryidae Jankowski, in Batisse,1975
Tachyblastonidae Grell,1950
Tachyblaston Martin,1909
T. ephelotensis Martin,1909
= *Dactylophrya roscovia* Collin,1909
@ *Ephelota gemmipara* (host specific)
Engl. Channel
Thecacinetidae Matthes,1956
Thecacineta Collin,1909
T. inclusa (Meunier,1910)
@ usually dorsal side of Halacardidae
N Norway, Barents Sea, Kara Sea
T. halacari Schulz,1933
@ Halacardidae & Ostracods
Kieler Bucht
Urmulidae Fraipont,1878
Metacineta Bütschli,1889
Paracineta Collin,1911
P. oviformis Dons,1918
@ tube of *Spirorbis*
N Norway
P. irregularis Dons,1927
@ setae of *Pherusa plumosa*
W Norway
P. divisa Fraipont,1878
@ *Mysella bidentata* & *Mya arenaria*
Kieler Bucht, North Sea, Mediterranean, Black Sea
Urnula Claparède & Lachmann,1859
ENDOGENIDA Collin,1912
Acinetidae von Stein,1859
Acineta Ehrenberg,1833
A. foetida Eichwald,1847
@ *Gammarus* spp. (most often on distal end of pleopod peduncles)
Askö, Kieler Bucht, Öresund, Ifesfjord
A. sulcata Dons,1927
= *A. benesaeta* Schulz,1933
@ Halacardidae & Ostracoda
Kieler Bucht, N Norway
A. laevis Dons,1918
@ e.g. green algae, *Delesseria* & *Spirorbis*
Tromsø
A. dirisa
?France
A. tuberosa Ehrenberg,18??
@ *Gammarus*, *Idothea* etc.
Kieler Bucht, Halland, Roscoff
A. schulzi Kahl,1934
@ Copepoda
Kieler Bucht
A. amphiasci Precht,1935
@ *Amphiascius typhlops*
Kieler Bucht
A. branchicola Precht,1935
@ gills of *Carcinus maenas* & *Crangon crangon*
Kieler Bucht
A. laomedea Precht,1935
@ *Laomedea*
Kieler Bucht
A. harpacticicola Precht,1935
@ *Nitocra spinipes*
Kieler Bucht
A. papillifera
?France
Acinetopsis Robin,1879
Loricophrya Matthes,1956
Pseudocorynophrya Small & Lynn, in Lee, Hutner & Bovee,1985
Trematosoma Batisse,1972
Dendrosomatidae Fraipont,1878
Dendrosoma Ehrenberg,1838
Lernaeophrya Pérez,1903
Platophrya Gönnert,1935
Trichophrya Claparède & Lachmann,1859
T. piscium
@ fish
E Baltic
T. salparum Entz,1884

- = *Trichophrya ascidiarium* Lachmann,1859 (n. nud.)
 = *Actinobranchion salparum* : Théodorides,1899
 * there may be two species, one in salps and another in ascidians; if so the one in ascidians probably should be named *T. ascidiarium* Chatton & Lwoff,1939
 @ e.g. *Salpa* sp., *Thalia democratica*, *Ciona intestinalis*, *Botryllus*, *Molgula manhattensis*, *Ascidella aspersa* & *Ascidia callosa*
 ?France, Mediterranean, Woods Hole
- T. danæ** Grontved,1951
 * pelagic in surface water
 Denmark Strait, SE Greenland, Davis Strait
- T. pelagica** (von Daday,1888)
 = *Sphaerophrya pelagica* von Daday,1888
 * possibly a young form of *T. salparum*
 Gulf of Naples
- Endosphaeridae Jankowski, in Corliss,1979
- Endosphaera** Engelmann,1876
- Tokophryidae Jankowski, in Lee, Hutner & Bovee,1985
- Choanophrya** Hartog,1902
- Hypophrya** López-Ochoterená,1964
- Multifasciculatum** Goodrich & Jahn,1943
- Pottsiocles** Corliss,1960
 = *Pottsiac* Chatton & Lwoff,1927, non Ryder,1885 (Coelenterata)
 = *Pottsiodes* Corliss,1960
- P. infusorum** (Chatton & Lwoff,1927)
 = *Pottsiac infusorum* Chatton & Lwoff,1927 (TM)
 @ *Folliculinina ampulla*, *F. elegans*, *Cochurnia socialis* & *C. ingenita* & ?other folliculinids
 Monaco, ?Maine
- Pseudogemma** Collin,1909
- P. fraiponti** ?Collin,19??
 @ *Acineta dirisa*
 ?France
- P. pachystyla** ?Collin,19??
 @ *Acineta tuberosa*
 ?France
- P. keppeni** ?Collin,19??
 @ *Acineta papillifera*
 ?France
- Rhyncheta** Zenker,1866
- Tokophrya** Bütschi,1889
- T.** sp. : Precht,1935
 @ *Neomysis vulgaris*
 Kieler Bucht
- EVAGINOCENIDA Jankowski, in Corliss,1979
- Cyathodiniidae da Cunha,1914
- Cyathodinium** da Cunha,1914
- Dendrocometidae Haeckel,1866
- Cometodendron** Swarczewsky,1928
- Dendrocometes** von Stein,1852
- Dendrocometides** Swarczewsky,1928
- Stylocometes** von Stein,1867
- Discophryidae Collin,1912
- Anarma** Goodrich & Jahn,1943
- Discophrya** Lachmann,1859
- Periacineta** Collin,1909
- Prodiscophrya** Kornos,1935
- Squalorophrya** Goodrich & Jahn,1943
- Testudinicola** Jahn, Bovee & Jahn,1979
- Heliophryidae Corliss,1979
- Heliofrya** Saedeleer & Tellier,1930
- NASSOPHOREA** Small & Lynn,1981
- NASSOPHORIA** Small & Lynn,1981
- SYNHYMENIIDA de Puytorac & al.,1974
- Orthodonellidae Jankowski,1968
- Orthodonella** Bhatia,1936
 = *Orthodon* (homonym) = *Rhabdodon* (homonym)
- O. hamatus** (Gruber,1???)
 ?
- Zosterodasys** Deroux,1978
- Nassulopsidae Deroux, in Corliss,1979
- Nassulopsis** Fauré-Fremiet,1959
- Phasmatopsis** Deroux,1978
- Scaphidiomatidae Deroux, in Corliss,1979
- Chilodontopsis** Blochmann,1895
- C. elongata** Kahl,19??
 Øresund
- C. transversa** Kahl,19??
 ?
- C. vorax** (Stokes,1???)
 ?
- C. caudata** Kahl,19??
 S Baltic, Plymouth
- Scaphidiomatidae von Stein,1959
- S. navicula** (O.F. Müller,17??) von Stein,1859
 Sylt, Yorkshire, Mauretania
- Synhymeniidae Jankowski, in Lee, Hutner & Bovee,1985
- Synhymenia** Jankowski,1967
- NASSULIDA Jankowski,1967
- NASSULINA Jankowski,1967
- Nassulidae de Fromental,1874
- Nassula** Ehrenberg,1833
- N. aurea** Ehrenberg,18??
 Plymouth
- N. citrea** Kahl,19??
 ?
- N. halophila** Gurwitsch,1???
- Stomatophrya** Kahl,1933
 * Incertae sedis
- S. singularis** Kahl,1933 (TM)
 Kieler Bucht
- PARAHYMENOSTOMATINA Grain & al.,1976
- Furgasoniidae Corliss,1979
- Furgasonia** Jankowski,1964
- Paranassulidae Fauré-Fremiet,1962
- Enneameron** Jankowski,1964
- Paranassula** Kahl,1931
- P. brunnea** (Fabre-Domergue,18???)
 ?
- MICROTHORACIDA Jankowski,1967
- Discotrichidae Jankowski,1975
- Discotricha** Tuffrau,1954
- D. papillifera** Tuffrau,1954 (TM)
 Øresund
- Microthoracidae Wrzesniowski,1870
- Drepanomonas** Fresenius,1858
- Kreyella** Kahl,1931
- Microrhox** Engelmann,1862
- PROPENICULIDA Small & Lynn, in Lee, Hutner & Bovee,1985
- Leptopharyngidae Kahl,1926
- Leptopharynx** Mermod,1914
 = *Trichopelma* Levander,1900, non Simon,1888 (Arachnida)
- L. torpens** (Kahl,19???)
 = *Trichopelma torpens* Kahl,19???
 Plymouth
- Pseudomicrothorax** Mermod,1914
- PENICULIDA Fauré-Fremiet, in Corliss,1956
- FRONTONIINA Small & Lynn, in Lee, Hutner & Bovee,1985
- Clathrostomatidae Kahl,1926
- Clatrostoma** Penard,1922
- Frontoniidae Kahl,1926
- Disematostoma** Lauterborn,1894
- Didieria** Small & Lynn, in Lee, Hutner & Bovee,1985
- Frontonia** Ehrenberg,1838
- F. arenaria** Kahl,1933
 Kieler Bucht, Øresund, Sylt, Barents Sea, Murmansk, White Sea, Mediterranean Black Sea
- F. marina** Fabre-Domergue,1891
 Askö, W Baltic, Øresund, Hiddensee, Sylt, Barents Sea, Plymouth, French Atlantic, Mediterranean, Black Sea, Kamerun, E USA
- F. algivora** Kahl,19??
 Øresund
- F. microstoma** Kahl,19??
 Askö, Øresund
- F. fusca** var.? Kahl,19??
 Kieler Bucht
- Paraclathrostoma** Small & Lynn, in Lee, Hutner & Bovee,1985
- Schistophrya** Kahl,1933
 * incertae sedis

- S. aplanata* Kahl,1933 (TM)
S Baltic
- Wenrichia* Jankowski,1967
- Lembadionidae Jankowski, in Corliss,1979
- Lembadion* Perty,1849
- Maritujidae Jankowski, in Lee, Hutner & Bovee,1985
- Marituja* Gajewskaja,1928
- Stokesiidae Roque,1961
- Stokesia* Wenrich,1929
- PARAMECIINA Jankowski, in Lee, Hutner & Bovee,1985
- Parameciidae Dujardin,1840
- Paramecium* O.F. Müller,1773
* mainly fresh water species, a few species in the "bursaria" group occur in salt & brackish water.
- P. calkinsi* Woodruff,19??
Öresund
- P. sp. :* Hartwig,1973
Sylt
- Neobursaridiidae Dragesco & Tuffrau,1967
- Neobursaridium* Balech,1941
- Urocentridae Claparède & Lachmann,1858
- Urocentrum* Nitzsch,1827
- HYPOTRICHIA** von Stein,1859
- EUPLOTIDA Small & Lynn, in Lee, Hutner & Bovee,1985
- EUPLOTINA Small & Lynn, in Lee, Hutner & Bovee,1985
- Aspidiscidae Ehrenberg,1838
- Aspidisca* Ehrenberg,1838
- A. angulata* Bock,1952
S Baltic, Öresund
- A. baltica* Kahl,1932
= *A. pulcherrima* var. *baltica* Kahl,1932
Baltic, Sylt
- A. pulcherrima* Kahl,1932
Schlei, Danziger Bucht, Öresund, Sylt, Plymouth, Roscoff
- A. major* Madsen,1931
W Baltic, Öresund, Sylt, White Sea, Yorkshire, French Atlantic, Mediterranean, E USA
- A. lyncker* (O.F. Müller,177?) von Stein,1859
= *Trichoda lyncker* O.F. Müller,17??
Kieler Bucht, Halland, Sylt, Helgoland, White Sea, French Atlantic, Black Sea
- A. sedigita* Quennerstedt,1867
Schlei, Kieler Bucht, Halland, Sylt, White Sea, French Atlantic
- A. fusca* Kahl,1928
Kieler Bucht, Öresund, Sylt, White Sea, Mauretania
- A. irinae* Burkovsky,1970
Sylt, Yorkshire, White Sea
- A. costata* (Dujardin,18??)
Plymouth
- A. hexera* Quennerstedt,1869
Visby, Plymouth
- A. lynceus* Ehrenberg,18??
Visby, Öresund, Plymouth
- A. turrita* (Ehrenberg,18??)
Plymouth
- A. binucleata* Kahl,19??
Kieler Bucht
- A. mutans* Kahl,19??
Kieler Bucht
- A. robusta* Kahl,19??
S Baltic
- A. aculeata* (Ehrenberg,1838)
New Hampshire
- A. polystyla* von Stein,18??
Halland
- A. steini* (von Buddenbrock,19??)
?
- A. crenata* Fabre-Domergue,18??
?
- Euplotaspis* Chatton & Séguela,1936
- E. cionaecola* Chatton & Séguela,1936
@ e.g. *Ciona intestinalis* & *Ascidia callosa*
?
- Euplotidae* Ehrenberg,1838
- Certesia* Fabre-Domergue,1885
- C. quadrinucleata* Fabre-Domergue,1885?
S Baltic
- Diophrys* Dujardin,1841
- D. appendiculata* (Ehrenberg,1838) von Stein,1859
- = *Styloynchia appendiculata* Ehrenberg,1838
- = *Diophrys marina* Dujardin,1841 (TM)
- ?= *Schizopus norvegicus* Claparède & Lachmann,1858
- S Baltic, Öresund, Sylt, Helgoland, White Sea, Yorkshire, Plymouth, French Atlantic, Mediterranean, Black Sea, Mauretania, E USA, Brazil
- D. scutum* (Dujardin,1841) Kahl,1932
= *Ploesconia scutum* Dujardin,1841 = *Styloplotes grandis* Rees,1881
Askö, S Baltic, Danziger Bucht, Öresund, Sylt, Helgoland, N Russia, Plymouth, French Atlantic, Mediterranean, Black Sea, Mauretania, E USA
- D. irmgard* Mansfeld,1923
Kieler Bucht, Sylt, White Sea, French Atlantic, Mediterranean, Florida
- D. kahli* Dragesco,1963
Kieler Bucht, Sylt, Roscoff
- D. multinucleata* Hartwig,1973
Sylt, Yorkshire
- D. tetramacronucleata* Kattar,1970
Sylt, Brazil
- D. oligothrix* Borror,1965
New Hampshire
- D. histrix* von Buddenbrock,1920
?
- D. scutooides* Agamaliev,1967
Caspian Sea
- D. magnus* Raikov & Kovaleva,1969
Japan Sea
- D. peloetes* Borror,1965
Florida
- D. quadricaudatus* Agamaliev,1967
Caspian Sea
- D. kasymovi* Agamaliev,1971
Caspian Sea
- Euplates* Ehrenberg,1830
- E. alatus* Kahl,1932
New Hampshire
- E. balticus* Kahl,1932
Kieler Bucht, Sylt, Helgoland, Mediterranean
- E. charon* (O.F. Müller,1773) Ehrenberg,1832
= *Trichoda charon* O.F. Müller,1773
* very common in rotting sea water
Visby, Halland, New Hampshire
- E. harpa* von Stein,1859
Askö, Öresund, Sylt, Plymouth, Mediterranean, Black Sea
- E. harpa* var. *marinus* Rossolimo,1???
Plymouth
- E. mediterraneus* Fernandez-Leborans,1985
Mediterranean
- E. moebiusi* Kahl,1932
Schlei, Kieler Bucht, Öresund, Hiddensee, Sylt, White Sea, Florida
- E. moebiusi* f. *quadricirratus* Kahl,1932
?
- E. trisulcatus* Kahl,1932
Danziger Bucht, Kieler Bucht, Belt Sea, Sylt, White Sea, France, Florida
- E. eurystomus* (Wrzesniowski,1870) Kahl,1932
Sylt
- E. vannus* (O.F. Müller,1786)
S Baltic, Öresund, Plymouth
- E. crassus* Dujardin,1841
* is forming a group together with *E. vannus* (O.F. M.)
?
- E. elegans* Kahl,1932
Öresund
- E. elegans* f. *litoralis* Kahl,1932
Kieler Bucht
- E. balteatus* (Dujardin,1842) Kahl,1932
@ found on *Strongylocentrotus droebachiensis* by Beers,1952
Belt Sea, White Sea, Yorkshire, French Atlantic, Mediterranean, E USA
- E. bisulcatus* Kahl,1932
Kieler Bucht, New Hampshire
- E. cristatus* Kahl,1932
S Baltic
- E. minuta* Yocom,1930
New Hampshire, Puget Sound
- E. salina* Yocom,1930
Puget Sound
- E. violacea* Kahl,19??
?
- E. quinquecarinatus* Gelei,1950
New Hampshire
- Swedmarkia* Dragesco,1954
- S. arenicola* Dragesco,1954 (TM)
Öresund, Sylt, Yorkshire, French Atlantic, S France
- Gastrocirrhidae Fauré-Fremiet,1961
- Cytharoides* Tuffrau,1975
- Euplotidium* Noland,1937

- Gastrocirrus** Lepsi,1928
- Uronychiidae** Jankowski, in Lee, Hutner & Bovee,1985
- Uronychia** von Stein,1859
- U. transfuga** (O.F. Müller,1786)
= *Trichoda transfuga* O.F. Müller,1786
E & W Baltic, Öresund, Halland, Kristineberg, Sylt, N Russia, Plymouth, Whitstable, French Atlantic, Mediterranean, Black Sea, E USA
- U. bivalvorum** Fenchel,1965
@ *Thysira* spp.
Öresund, Gullmarfjord
- U. uncinata** Taylor,1???
S Baltic
- U. setigera** Calkins,
Plymouth
- DISCOCEPHALINA** Wicklow,1982
- Discocephalidae** Jankowski,1979
- Discocephalus** Ehrenberg, in Hemprich & Ehrenberg,1828
- D. ehrenbergi** Dragesco,1960
Kieler Bucht, Öresund, Sylt, White Sea, Yorkshire, Roscoff, Arcachon
- D. rotatorius** Ehrenberg,1838
S Baltic, Sylt, White Sea, Yorkshire, French Atlantic, Mediterranean, Cape Cod
- D. minimus** Dragesco,1968
?
- D. grandis** Dragesco,1954
?
- Marginotricha** Jankowski,1978
- Prodiscocephalus** Jankowski,1979
- Psammocephalus** Wicklow,1982
- Erionellidae** Wicklow,1982
- Erionella** Jankowski,1978
- OLIGOHYMENOPHOREA** de Puytorac & al.,1974
- HYMENOSTOMATIA** Delage & Hérouard,1896
- HYMENOSTOMATIDA** Delage & Hérouard,1896
- TETRAHYMENINA** Fauré-Fremiet, in Corliss,1956
- Curimostomatidae** Jankowski,1968
@ in haemocoel of arthropods or freeliving
- Curimostoma** Kozloff,1954
- Glaucomidae** Corliss,1971
- Epenardia** Corliss,1971
- Espejoia** Bürger,1908
- Glaucoma** Ehrenberg,1830
- G. vorax** Kidder, Lilly & Claff,1940
Woods Hole
- Glaucomella** Grolieré,1977
- Jaacorliisia** Small & Lynn, in Lee, Hutner & Bovee,1985
- Tetrahymenidae** Corliss,1952
- Deltopyrum** Fauré-Fremiet & Mugard,1946
- Lambornella** Keilin,1921
- Tetrahymena** Furgason,1940
- Turaniellidae** Didier,1971
- Colpidium** von Stein,1860
- Turaniella** Corliss,1960
- OPHYROGLLENINA** Canella,1964
- Ichthyophthiriidae** Kent,1881
- Cryptocaryon** Brown,1951
- C. irritans** Brown,1951 (TM)
= *Ichthyophthirus marinus* Sikama,1961
@ all marine teleosts kept at 20-26° C; may often reach enzootic proportions in aquaria
Marine aquaria, Warmer Seas
- Ichthyophthirius** Fouquet,1876
@ the cause of the "ich" disease of freshwater fish, known in China since Sung dynasty
- I. multifiliis** Fouquet,1876?
@ chiefly fresh water teleosts, but survives salinities lower than 15 ppt on e.g. eels
Unterelbe, Berliner Aquarium
- Ophryogenidae** Kent,1881
- Ophryoglena** Ehrenberg,1831
- ?**O. fusca** (Quennerstedt,1869)
= *Panophrys fusca* Quennerstedt,1869
Visby
- O. sp. : Fenchel,1968**
Askö
- SCUTICOCILIATIDA** Small,1967
- PHILASTERINA** Small,1967
- Ptyssostoma** Hentschel,1927
* Incertae sedis in this suborder
- P. thalassemae** ?Hentschel,1927? (TM)
@ alimentary tract of *Thalassema neptuni*
?Plymouth
- Cinetochilidae** Perty,1852
- Cinetochilum** Perty,1849
- Platynematidae** Kahl,1935
= *Platynema* (homonym)
- P. denticulatum** Kahl,19??
S Baltic, N Öresund, Helgoland
- P. hyalinum** Kahl,19??
S Baltic, Helgoland
- Pseudoplatynematum** Bock,1952
- P. loricatum** Bock,1952?
S Baltic
- P. parvum** Bock,1952?
S Baltic
- Sphenostomella** Jankowski, in Lee, Hutner & Bovee,1985
- Cohnilembus** Kahl,1933
= *Lembus* Cohn,1866, non Guenther,1860 (Pisces)
- C. verminus** (O.F. Müller,1786)
= *Lembus velifer* Cohn,1866 (TM of *Lembus* Cohn,1866 - North Sea Aquarium)
= *Cyclidium elongatum* Claparède & Lachmann,18??
* very common in rotting sea water
Visby, S Baltic, Öresund, Oslofjord
- Cryptochilidae** Berger, in Corliss,1979
- Cryptochilum** Maupas,1883
- C. cuenoti**
@ gut of *Golfingia vulgaris*
Europe
- C. echini** ?Cuenot,1912?
@ gut, coelomic cavity and gonads of *Paracentrotus lividus* & *Psammechinus miliaris*
Arcachon
- Biggaria** Kahl,1934
- Entodiscidae** Jankowski,1973
- Entodiscus** Madsen,1931
- E. boreale** (Hentschel,1924)
= *Cryptochilum boreale* Hentschel,1924 (TM)
@ gut of *Echinus esculentus* & *Strongylocentrotus droebachiensis*
Frederikshavn, Shetlands, Scotland
- E. indomitus** Madsen,1931
@ gut of *Strongylocentrotus droebachiensis*
Frederikshavn
- Entorhipedidae** Madsen,1931
- Entorhipedium** Lynch,1929
- Loxocephalidae** Jankowski,1964
- Balanonema** Kahl,1931
- Cardiostomatella** Corliss,1960
= *Cardiostoma* Kahl,1928, non Sandberger,1872 (Mollusca)
* Incertae sedis
- C. vermiforme** (Kahl,1928)
= *Cardiostoma vermiforme* Kahl,1928
W Baltic, Öresund, Sylt, White Sea, French Atlantic, Mediterranean, Mauretania, E USA
- C. mononucleata** Dragesco,19??
Öresund
- Dexiotricha** Stokes,1885
- Loxocephalus** Eberhard,1862
- Paradexiotricha** Grolieré,1975
- Paraloxocephalus** Small & Lynn, in Lee, Hutner & Bovee,1985
- Paratetrahymena** Thompson,1963
- Paralembidae** Corliss & de Puytorac, in Lee, Hutner & Bovee,1985
- Magnalembus** Small & Lynn, in Lee, Hutner & Bovee,1985
- Mesolembus** Small & Lynn, in Lee, Hutner & Bovee,1985
- Ovolembus** Small & Lynn, in Lee, Hutner & Bovee,1985
- Paralembus** Kahl,1933
- ?
P. digitiformis Kahl,19???
- ?
P. rostratus Kahl,19???
- ?
P. asturianus Fernandez-Leborans, Castro & Zaldumbide,1990
?S Spain
- Paranophryidae** Jankowski, in Lee, Hutner & Bovee,1985
- Anophryoides** de Puytorac Grolieré,1979
- Mesanophrys** Small & Lynn, in Lee, Hutner & Bovee,1985

- Metanophrys*** de Puytorac & al., 1974
- Mugardia*** Small & Lynn, in Lee, Hutner & Bovee, 1985
- Paranophrys*** Thompson & Berger, 1965
- P. maggi*** (Cattaneo, 1???)
= *Anophrys maggi* Cattaneo, 1???
- = *Anophrys sarcophaga* Cohn, 1866
?= *Lembus sarcophagus* : Wallengren, 1918
= *Paranophrys carcinii* Grolier & Leglise, 1977
@ facultative pathogen in e.g. *Carcinus maenas* & *Cancer pagurus*
Öresund, Bretagne
- Parauronematidae*** Small & Lynn, in Lee, Hutner & Bovee, 1985
- Glauconema*** Thompson, 1966
- Miamiensis*** Thompson & Moewus, 1964
- M. avidis*** Thompson & Moewus, 1964
- ? ***Parauronema*** Thompson, 1967
- Potomacus*** Thompson, 1966
- Philasteridae*** Kahl, 1931
* Fenchel, 1968 depicts an undescribed Philasterid, which attack unmetamorphosed *Spisula* larvae, held in culture in Elsinore
- Helicostoma*** Cohn, 1866
- H. notatum*** (Möbius, 18??)
Askö
- H. oblongum*** Cohn, 18??
? in Öresund
- H. buddenbrocki*** Kahl, 1931
@ skin of young *Zaorces viviparus*, *Solea solea* & *Pleuronectes platessa*
* may also be found interstitially
S Baltic
- H.* sp. : Hartwig, 1973
Sylt**
- Philaster*** Fabre-Domergue, 1885
- P. digitiformis*** Fabre-Domergue, 1885?
?
- Philasterides*** Kahl, 1931
- P. armata*** (Kahl, 19??)
?
- Pseudocoelomidae*** Evans & Thomson, 1964
- Anophrys*** Cohn, 1866
* Incertae sedis
- A. arenicola*** Fernandez-Leborans, 1986
?S Spain
- A. sarcophaga*** Cohn, 1866 (TM)
= *Lembus sarcophagus* : Auctt.
Öresund
- A. maggi***
@ pathogenic to *Carcinus maenas*
?
- Pseudocoelomitus*** Evans & Thomson, 1964
- P. pusillus*** (Quennerstedt, 1869)
= *Lembus pusillus* Quennerstedt, 1869 = *Cohnilembus pusillus* : Kahl, 1930-35
* very common in rotting sea water
Visby, Öresund, Halland, Plymouth
- P. cantabricus*** Fernandez-Leborans, Castro & Zaldumbide, 1990
?S Spain
- P. antoniensis*** Fernandez-Leborans, 1986
?S Spain
- P. portuensis*** Fernandez-Leborans, 1986
?S Spain
- P. marinus*** Thompson, 1966
Virginia
- Thyrophylacidae*** Berger, in Corliss, 1961
- Flagiopyliella*** Poljansky, 1951
- Thyrophylax*** Berger, in Corliss, 1961
- Thigmophryidae*** Chatton & Lwoff, 1926
- Conchophyllum*** Raabe, 1936
- Thigmophryna*** Chatton & Lwoff, 1923
@ in mantle cavity of marine bivalves (with one exception)
- T. saxicavae*** Fenchel, 1964
@ *Hiatella arctica* &/or *H. striata*
Gullmarfjord
- T. macomaee*** Chatton & Lwoff, 1926
@ *Macoma balthica*
Askö, Gdynia, Wimereux, non Öresund
- T. annella*** Fenchel, 1965
@ intestine of *Malacobdella grossa* in *Arctica islandica*
Gullmarfjord
- T. cf. T. bivalvorum*** Chatton & Lwoff, 1923 : Fenchel, 1965
* *T. bivalvorum* C. & Lw. is known from *Venerupis pullastra* & *Mactra* (= *Spisula*) *solida* from the French channel coast
@ *Spisula elliptica*
Öresund
- T.* sp. : Fenchel, 1965
@ *Phoxas pellucidus*
Öresund**
- T.* sp. : Fenchel, 1965
@ *Cerastoderma ovale*
Öresund**
- T. tapetis*** Chatton & Lwoff, 1926
@ *Venerupis pullastra*
Wimereux, non Gullmarfjord
- Uronematidae*** Thompson, 1964
- Homologastridae*** Kahl, 1926
- Urocyclidae*** Small & Lynn, in Lee, Hutner & Bovee, 1985
- Uronema*** Dujardin, 1841
- U. marinum*** Dujardin, 1841
@ this bacteriophage may cause high mortalities in oyster hatcheries & may also cause enzootics & mortalities among different aquarium fish species
* very common in rotting sea water
Visby, Kieler Bucht, Öresund, Limfjorden, Frederikshavn, Sylt, Helgoland, Barents Sea, White Sea, Plymouth, Whitstable, E USA
- U. filiforme*** Kahl, 1932
Plymouth, New Hampshire
- U. acutum*** von Buddenbrock, 1920
New Hampshire, Gulf of Mexico
- U. acutum* v. *multistriatum*** Madsen, 1931
Frederikshavn
- U. elegans*** Maupas, 1???
- Öresund
- U. pluricaudatum*** Noland, 1937
?
- U. constantiana*** Tucalesco, 1962
?
- U. botuliniformis*** Wenzel, 1961
* probably a member of Philasteridae sensu Borror, 1965
?
- U. castellonensis*** Fernandez-Leborans, Castro & Zaldumbide, 1990
?S Spain
- Uropedalium*** Kahl, 1928
- U. pyriforme*** Kahl, 19??
S Baltic
- Urozonidae*** Grolier, 1975
- Urozona*** Schewiakoff, 1889
- PLEURONEMATINA*** Fauré-Fremiet, in Corliss, 1956
- Calyptotrichidae*** Small & Lynn, in Lee, Hutner & Bovee, 1985
- Calyptotricha*** Phillips, 1882
- Conchophthiridae*** Kahl, in Doflein & Reichenow, 1929
- Conchophthirus*** von Stein, 1861
@ in mantle cavity of freshwater bivalves
- Ctedectomatidae*** Small & Lynn, in Lee, Hutner & Bovee, 1985
- Compososomella*** Small & Lynn, in Lee, Hutner & Bovee, 1985
- Ctedectoma*** Stokes, 1884
- Hippocomos*** Czapik & Jordan, 1977
- Paractedotoma*** Small & Lynn, in Lee, Hutner & Bovee, 1985
- Cyclidiidae*** Ehrenberg, 1838
- Cristigera*** Roux, 1899
- C. media*** Kahl, 19??
Oldesloe, Öresund, Plymouth
- C. phoenix*** Penard,
Plymouth
- C. minuta*** Kahl, 19??
Kieler Bucht
- C. setosa*** Kahl, 19??
S Baltic
- C. sulcata*** Kahl, 19??
Kieler Bucht
- C. cirrifera*** Kahl, 1928
Frederikshavn
- C. consticta*** Madsen, 1931
Frederikshavn
- C. vestita*** ??, 1???
- Oldesloe
- Cyclidium*** O.F. Müller, 1786
- C. citrullus*** Cohn, 1865
* very common in rotting sea water
Visby, Öresund, Kattegatt, Limfjorden, Frederikshavn
- C. candens*** Kahl, 19??
S Baltic, ?Plymouth
- C. glaucoma*** O.F. Müller, 1786
Frederikshavn, Plymouth

- C. elongatum* Schewiakoff,19??
S Baltic
- C. veliferum* Kahl,19??
Kieler Bucht
- C. plouneouri* Dragesco,1963
New Hampshire
- C. marinum* Borror,1963
?
- Echinocyclidium* Jankowski, in Lee, Hutner & Bovee,1985
- Paracyclidium* Small & Lynn, in Lee, Hutner & Bovee,1985
- Pseudocyklidium* Small & Lynn, in Lee, Hutner & Bovee,1985
- Dragescoidae* Jankowski, in Lee, Hutner & Bovee,1985
- Dragescoa* Jankowski,1974
- Histiobalanitidae de Puytorac & Corliss, in Corliss,1979
- Histiobalanum* Stokes,1886
- H. marinum* (Kahl,1933)
= *Histiobalanitidum marinum* Kahl,1933
Kieler Bucht, Öresund, White Sea, Yorkshire, Roscoff, Concarneau
- H. marinum* v. *majus* (Kahl,19??)
Kieler Bucht
- Peniculostomatidae Fenchel,1965
- Echinosociella* Berger, in Lee, Hutner & Bovee,1985
- Peniculistoma* Jankowski,1964
= *Kidderia* Raabe,1934 (homonym) = *Morgania* Kahl,1934 (homonym)
- P. mytili* (DeMorgan,1925)
= *Conchophithirus mytili* DeMorgan,1925 (TM - Plymouth)
= *Kidderia mytili* : Raabe,1934 = *Morgania mytili* : Kahl,1934
@ mantle cavity (especially the foot) of *Mytilus edulis*
Askö, Gdynia, Öresund, Bohusl., Frederikshavn, Murmansk, Plymouth, Roscoff, E USA
- Pleuronematidae* Kent,1881
- Pleurocoptes* Wallengren,1896
- Pleuronema* Dujardin,1836
- P. marinum* Dujardin,1841
W Baltic, Öresund, Halland, Sylt, White Sea, Yorkshire, Plymouth, French Atlantic, Mediterranean, Black Sea, Mauretanian, E USA
- P. coronatum* Kent,1881
Kieler Bucht, Öresund, Hiddensee, Sylt, Helgoland, White Sea, Yorkshire, French Atlantic, Mediterranean, Black Sea, Kamerun, E USA, Brazil
- P. crassum* Dujardin,18??
Plymouth
- P. arenicola* Dragesco,1960
- P. oculata* Dragesco,1960
?
- P. simplex* Dragesco,1960
?
- Schizocalyptra* Dragesco,1968
- Thigmocomidae Kazubski,1958
- Thigmocoma* Kazubski,1958
- THIGMOTRICHINA Chatton & Lwoff,1922
- Ancistridae Issel,1903
- Ancistrum* Maupas,1883
- A. caudatum* Fenchel,1964
@ *Musculus niger* (Gray) & *Modiolus modiolus* (L.)
Öresund, Gullmarfjord
- A. dosiniae* Chatton & Lwoff,1926
@ *Dosima exoleta* (L.)
Gullmarfjord, Baie de Morlaix
- A. arcopagiae* Fenchel,1965
@ *Arcopagia crassa* (Pennant)
Gullmarfjord
- A. crassum* Fenchel,1965
?= *A. tapepis* Chatton & Lwoff,1926 (nomen nudum - in *Venerupis pullastra*)
@ *Venerupis aurea* (Gmelin) & *Venerupis pullastra* (Montagu)
Gullmarfjord
- A. venerupis* Fenchel,1965
@ *Venerupis aurea* (Gmelin)
Gullmarfjord
- A. nuculae* Fenchel,1965
@ *Nucula turgida* Leckebny & Marshall
Öresund, Gullmarfjord
- A. abrarium* Fenchel,1965
@ *Abra nitida* (Müller)
Gullmarfjord
- A. macomae* Chatton & Lwoff,1949
= *A. tellinae* : Raabe,1936, non Issel,1903
@ *Macoma balthica* (L.) & *Tellina tenuis* da Costa
Askö, Gdynia, Öresund, Roscoff
- A. hydriobiae* Raabe,1936
@ *Hydrobia ulvae* (Pennant) & *Hydrobia ventrosa* (Montagu)
Gdynia, Öresund, Frederikshavn
- A. mytili* (Quennerstedt,1867)
- = *Opalina mytili* Quennerstedt,1867 (TM)
= *Anoplophrya mytili* : Kent,1882
@ *Mytilus edulis* L. (& *Modiolus modiolus* (L.))
Askö, Gdynia, Öresund, Halland, Bohusl., Frederikshavn, Plymouth, Roscoff, Mediterranean, E USA
- A. isseli* Kahl,19??
@ *Modiolus barbatus* & *Modiolus modiolus*
Mediterranean, NE America
- Ancistrumina* Raabe,1959
- A. kofoidi* (Busch,1937)
= *Ancistrina kofoidi* Busch,1937
@ *Petricola pholadiformis*
California
- Protophyra* Kofoed,1903
- P. ovicola* Kofoed,1903 (TM - New Port - in *Littorina saxatilis*)
= *Isselina intermedia* Cépéde,1910
@ mantle cavity of *Littorina* spp.
Gullmarfjord, Frederikshavn, Plymouth, Boulogne
- Hemispeiridae* König,1894
- Ancistrospira* Chatton & Lwoff,1926
- A. venera* Chatton & Lwoff,1926
@ *Astarte elliptica*, *Astarte sulcata*, *Astarte fusca*, *Venus fasciata*
Öresund, Gullmarfjord, S France
- Plagiospira* Issel,1903
- P. crinita* Issel,1903
@ *Loripes lacteus* (Poli), *Cardita calyculata* Brugière, *Divaricella divaricata* (L.), *Nucula nucleus* (L.), *Nucula turgida* Leckebny & Marshall, *Thyasira flexuosa* (Montagu), *Thyasira sarsi* (Philippi) & *Thyasira* sp.
Öresund, Gullmarfjord, Roscoff, Mediterranean
- Hemispeiroidea* Fabre-Domergue,1888
Hemispeiroips König,1894
- H. asteriasi* Fabre-Domergue,1888 (TM - Concarneau)
@ *Astieria rubens* & *Mariasterias glacialis*
Gullmarfjord, Frederikshavn, Concarneau, S France
- H. antedonis* (Cuénnot,1891)
= *Trichodina antedonis* Cuénnot,1891 = *Hemispeiroips comataulae* König,1894
@ *Anedon mediterranea* Lamarck & *Antedon petasus* Dübén & Koren
Gullmarfjord, Concarneau
- Boveria* Stevens,1901 (TM *B. subcylindrica* Stevens,1901 - ?California, in waterlings of a holothurian)
- B.* sp. : Fenchel,1965
@ *Monia patelliforis*
Læsø, Bohusl.
- B. teredinidi* Nelson,1923
@ *Teredo navalis*
?
- Proboveria* Chatton & Lwoff,1936 (TM *P. loripedes* ?Chatton & Lwoff?,1936? - Roscoff & Sète, in *Loripes lacteus* (Poli))
- P. thyasirae* Fenchel,1965
@ *Thyasira sarsi* & *T. sp.*
Gullmarfjord
- P. triangularis* Fenchel,1965
@ *Thyasira* sp.
Gullmarfjord
- Hysteroconitidae Diesing,1866
- Hysteroconita* Diesing,1866
- Kozloffia* de Puytorac,1968
- Ptychostomum* von Stein,1860
- PERITRICHIA** von Stein,1859
- SESSILIDA** Kahl,1933
- Astylozoidae** Kahl,1933
- Astylozoon* Engelmann,1862
- Hasatella* Erlanger,1890
- Ellobiophryidae Chatton & Lwoff,1929
- Ellobiophrya* Chatton & Lwoff,1923
- E. donaci* Chatton & Lwoff,1923 (TM)
@ *Donax vittatus*
Roscoff
- Epistylididae* Kahl,1933
- Campanella* Goldfuss,1820
- Epistylis* Ehrenberg,1830
- E. gammaris* Precht,1935
@ on antennae of *Gammareus oceanicus*, *G. salinus* & *G. zaddachi*
Askö, Kieler Bucht, Öresund, Isefjord
- E. arenicola* (Fabre-Domergue,1888)
= *Rhabdosyllis arenicola* Fabre-Domergue,1888
@ gills of *Arenicola marina*
Kieler Bucht, Roscoff, Concarneau
- E. carcinii* Precht,1935
@ gill cavity of *Carcinus maenas*
Kieler Bucht
- E. nitocrae* Precht,1935
@ *Nitocra spinipes*
Kieler Bucht
- E. jaerae* Precht,1935
@ gills of *Jaera* sp.
Kieler Bucht

- E. harpacticola* Kahl,1933
@ *Halicyclops magniceps*
Kieler Bucht
- Heteropolaria* Foissner & Schubert,1977
- Nidula* Gajewska,1933
- Opisthostyla* Stokes,1886
- O. sertularium* (Kent,1881?)
Plymouth
- Rhabdostyla* Kent,1881
- R. variabilis* (Dons,1918)
= *Scyphidia variabilis* Dons,1918
@ *Scoloplos armiger* & *Phyllodoce laminata*
Kieler Bucht, Tromsø
- R. commensalis* Möbius,1888
@ body cuticula of *Capitella* sp. & tentacles of *Terebellides stroemi*
Kieler Bucht
- R. nereicola* Precht,1935
Platynereis sp.
Kieler bucht
- R. arenaria* Cuénot,1891
@ *Asterias rubens*, *Leptosynapta*, *Astropecten*, etc.
Kieler Bucht, Roscoff
- R. putrina* (O.F. Müller,1776) Warren,1986
= *Vorticella putrina* O.F. Müller,1776
* marine?
?
- Lagenophryidae Bütschli,1889
- Circolagenophrys* Jankowski,1980
- Lagenophrys* von Stein,1851
- L. rotunda* Precht,1935
@ Ostracode
Kieler Bucht
- L. lunatus* Ima,19??
@ brackish water palaemonid shrimps
- L.* sp. : Fenchel,1965
@ *Gammarus locusta* L. & sometimes *G. oceanicus* Segerstråle
Öresund
- Operculariidae Fauré-Fremiet, in Corliss,1979
- Ampullaster* Jankowski,1976
- Opercularia* Goldfuss,1820
- Orbiculariaria* Lust,1950
- Propyxidium* Corliss,1979
- Ophrydiidae Ehrenberg,1838
- Ophrydium* Bory de St. Vincent,1826
- Opisthonectidae Foissner,1976
- Opisthonecta* Fauré-Fremiet,1906
- Telotrochidium* Kent,1881
- Rovinjellidae Matthes,1972
- Rovinjella* Matthes,1972
- Scyphidiidae Kahl,1933
- Ambiphryra* Raabe,1952
- A. miri* Raabe,1952
@ *Nerophis ophidion*
Baltic
- Apiosoma* Blanchard,1885
- A.* sp. : Calenius & Bylund,1980
@ fish
E Baltic
- Paravorticella* Kahl,1933
- P. terebellae* (Fauré-Fremiet,1920)
@ frontoventral on *Terebellides stroemi* & on skin of *Terebella lapidaria*
Stoller Grund, France
- Scyphidia Dujardin,1841
- S. hydriobiae* Kahl,1933
@ *Hydrobia* & *Littorina*
Kieler Bucht
- S. spinicola* Precht,1935
@ Spinoids, e.g. *Pygospio elegans*
Kieler Bucht
- S. terebellidis* Precht,1935
@ gills & tentacles of *Terebellides stroemi*
Kieler Bucht
- S. gasterosteii* (Fauré-Fremiet,1905)
= *Epistylis gasterosteii* Fauré-Fremiet,1905
@ gills & chest of *Gasterosteus aculeatus*
Kieler Bucht
- S. adunconucleata* MacKenzie,1969
@ *Pleuronectes platearessa*
Scotland
- S. physarum* Lachmann,1856
@ *Neritina fluviarialis*
Baltic (Visby)
- Termitophryidae Lom, in Corliss,1979
- Termitophrya* Noiro-Timothée,1959
- Vaginicolidae de Fromentel,1874
- Cothurnia* Ehrenberg,1831
- C. gammari* Precht,1935
@ inside of sideplates of *Gammarus locusta* L., *G. oceanicus* Segerstråle & *G. duebeni*
Kieler Bucht, Öresund, Isefjord, non Askö
- C. plectostyla* Stokes,
Plymouth
- C. socialis* Gruber,1879
Finland, Trondheim, North Sea, Vancouver
- C. nebuliae* Dons,1927
@ telson of *Nebalia* sp.
Trondheim, Tromsø
- C. limnoriae* Dons,1927
@ *Limnoria lignorum*
Varanger
- C. pedunculata* Dons,1918
= *C. nodosa* : Mereschkowsky,1879 (p.p.), non Claparède & Lachmann,1858
@ algae & different animals
Kieler Bucht, Trondheim, Tromsø, Iceland, White Sea
- C. grandis* (Perty,1???)
@ hydroids
Oslofjord, Tromsø, Pacific
- C. cypridicola* Kahl,1933
@ algae & different animals
Kieler Bucht
- C. acuta* Levander,1915
@ setae of *Harmonia* spp.
Tvärrimme, Kieler Bucht
- C. cordylophorae* Kahl,1933
@ *Cordylophora caspia*
Kieler Bucht, Bremerhaven
- C. recurva* Claparède & Lachmann,1858
@ Harpacicoids
Kieler Bucht, Helgoland
- C. angusta* Kahl,1933
@ Ostracods
Kieler Bucht
- C. ceramicola* Kahl,1933
Kieler Bucht
- C. harpactici* Kahl,1933
@ *Cletocampus confluentus* & *Mesochra lilljeborgi*
Kieler Bucht
- C. nitocrae* Precht,1935
@ furcal chaetae of *Nitocra spinipes*
Kieler Bucht
- C. peloscolicis* Precht,1935
@ setae of *Pherusa plumosa*
Kieler Bucht
- C. complanata* Precht,1935
@ setae of *Pherusa plumosa*
Kieler Bucht
- C. styloroididis* Precht,1935
@ setae of *Pherusa plumosa*
Kieler Bucht
- C. nereicola* Precht,1935
@ *Hediste diversicolor*
Kieler Bucht
- C. halacaricola* Precht,1935
@ rear extremities of *Copidognathus fabricius*
Kieler Bucht
- C. triangula* Precht,1935
@ *Copidognathus* & ostracods
Kieler Bucht
- C. maritima* Ehrenberg,18??
= *C. nodosa* Claparède & Lachmann,1858
@ algae
Visby
- Platycola* Kent,1882
- P. circularis* Dons,1940
@ uropods of *Limnoria lignorum*
Norway
- P. dilatata* Kent,1882?
= *P. donsi* Kahl,1933
* on varying substrates
Norway
- Pyxicola* Kent,1882
- P. socialis* (Gruber,1879)
@ algae, hydroids, molluscs etc.
Finland, Kieler Bucht, White Sea
- Thuricola* Kent,1881
- Vaginicola* de Lamarck,1816
- V. doliolum* Penard,
Plymouth

- Vorticellidae Ehrenberg,1838
- Carchesium** Ehrenberg,1830
- C. duplicatum** Precht,1935
@ *Gammarus oceanicus*
Askö, Kieler Bucht, Öresund, Isefjord
- C. gammari** Precht,1935
@ *Gammarus*
Kieler Bucht
- C. jaerae** Precht,1935
@ legs of *Jaera*
Kieler Bucht
- C. steinii** (Wrzesniowski,1877)
= *Epistylis steinii* Wrzesniowski,1877
@ *Gammarus*
* chiefly a fresh water species
Kieler Bucht
- C. abrae** Precht,1935
@ siphon bases of *Abra alba*
Stoller Grund
- C. sp. a :** Fenchel,1965
@ gills & pleopods of *Gammarus duebeni*
Askö
- Haplocaulus** Precht,1935
- H. nicoleae** Precht,1935 (TM)
@ *Nicolea zostericola*
Kieler Bucht
- H. furcellariae** Precht,1935
@ *Furcellaria*
Kieler Bucht
- H. amphiorae** (Cuénnot,1891) Warren,1986
= *Vorticella amphiorae* Cuénnot,1891
@ *Amphiura*
?
- H. claparedei** (Andrussowa,1886) Warren,1986
= *Vorticella claparedei* Andrussowa,1886
* marine?
?
- H. elegans** (Dons,1915) Stiller,1971
= *Vorticella elegans* Dons,1915 = *Intranstylum elegans* : Nenninger,1948
@ operculum of a *Spirorbis* sp.
Tromsø
- H. fusca** (Biernacka,1963) Warren,1986
= *Vorticella fusca* Biernacka,1963, non Precht,1935
Danziger Bucht
- Pseudocarchesium** Sommer,1951
- P. brevistyla** (d'Udekem,1864) Stiller,1971
= *Vorticella brevistyla* d'Udekem,1864
* marine?
?
- Pseudovorticella** Foissner & Schiffmann,1974
- P. difficilis** (Kahl,1933)
= *Vorticella difficilis* Kahl,1933
@ several brackish water animal species
Kieler Bucht
- P. micata** (Kahl,1933)
= *Vorticella micata* Kahl,1933
?
- P. punctata** (Dons,1918) Warren,1986
= *Vorticella punctata* Dons,1918 = *Vorticella perlata* Kahl,1933
@ several brackish water & marine animal species
Kieler Bucht, Norway
- Vorticella** Linnaeus,1767
- V. aequilata** Kahl,1939
- V. annulata** (Gourret & Roeser,1888) Kahl,1935
= *V. brevistyla* var. *annulata* Gourret & Roeser,1888
?
- V. anomala** Gourret & Roeser,1886
?
- V. biddulphae** Stiller,1939
= *V. variabilis* Stiller,1939
@ on *Biddulphia chinensis*
North Sea
- V. calciformis** Kahl,1933
?
- V. campanula** Ehrenberg,1831
= *V. macrostoma* Schmarda,1854 = *V. cylindrica* Dons,1918
@ *Diphasia abietina*
Tromsø
- V. campanulata** Sramek-Husek,1948
= *V. constricta* Kahl,1933, non Fromentel,1874
?
- V. communis** Fromentel,1874
= *V. subsphaerica* Dons,1918
@ *Dynamena pumila*
Tromsø
- V. convallaria** Linnaeus,1758
= *V. cyclopis* Kahl,1933 = *V. hyalina* Stiller,1940
?
- V. cratera** Kent,1881
= *V. dudekemi* Kahl,1933
?
- V. elongata** Fromentel,1874
= *V. subcylindrica* Ghosh,1922
?
- V. fusca** Precht,1935
@ *Enteromorpha*
Kieler Bucht
- V. granulata** Kahl,1933
?
- V. hamata** Ehrenberg,1831
= *V. delicatulata* Biernacka,1963 ?= *V. plicata* Gourret & Roeser,1886
?
- V. infusionum** Dujardin,1841
?
- V. jaerae** Precht,1935
@ attached to *Jaera marina*
Kieler Bucht
- V. lima** Kahl,1933
= *Pseudovorticella lima* (Kahl,1933) Jankowski,1976
?
- V. longiseta** Dietz,1964
?
- V. longitricha** Gajewska,1933
?
- V. marina** Greeff,1870
= *V. minuta* Precht,1935
@ e.g. *Ceranium*, *Furcellaria*, *Delesseria* & *Pallene*
Kieler Bucht
- V. mayeri** Fauré-Fremiet,1920
= *Pelagovorticella mayeri* (Fauré-Fremiet,1920) Jankowski,1980
* Pelagic
?
- V. mortensi** Dons,1922
* nomen dubium
?
- V. nutans** O.F. Müller,1773
?
- V. obconica** Kahl,1935
= *V. conica* Dons,1918, non Stokes,1887
@ *Spirorbis* sp.
Tromsø
- V. ovum** Dons,1918
= *V. incerta* Nenninger,1948
- V. patellina** O.F. Müller,1776
= *V. formicata* Dons,1915
@ e.g. *Carcinus*, *Mytilus*, *Gammarus*, *Laomedea* & *Laminaria*
Kieler Bucht, North Sea, Tromsø, W France, Woods Hole
- V. pulchra** Kahl,1933
?
- V. pyriforme** Stiller,1939
= *V. intermissa* Nenninger,1948
- V. robusta** Dons,1922
* nomen dubium
?
- V. striata** Dujardin,1841
= *V. latestriata* Sommer,1951 = *V. minima* Stiller,1939 = *V. suboctava* Sommer,1951
= *V. striatula* Dons,1918 = *V. oceanica* Zacharias,1906
@ many different vertebrates & invertebrates
Kieler Bucht, Tromsø
- V. subprocumbens** Ghosh,1922
= *V. nana* Kahl,1933 = *V. solitaria* Stiller,1935
?
- V. telescopica** Kent,1861, non Kahl,1935
= *V. telescopiformis* : Kahl,1935
- V. teninucleata** Dons,1922
* nomen dubium
?
- V. undulata** (Dons,1918) Kahl,1935
= *Vorticellopsis undulata* Dons,1918
* nomen dubium
Tromsø
- V. verrucosa** Dons,1918
= *V. singularis* Kahl,1933
@ *Laomedea* sp.
Tromsø
- V. sp. :** Fenchel,1965
@ *Gammarus duebeni*
Askö, non Isefjord
- Zoothamniidae** Sommer,1951
- Craspedomyoschiston** Precht,1935
= *Craspedomyoschiston* Jankowsky, in Lee, Hutner & Bovee,1985
- C. sphaeroma** Precht,1935
@ gills of *Sphaeroma rugicauda*
Kieler Bucht
- Intranstylum** Fauré-Fremiet,1904
- I. duplicatum** Precht,1935
@ inside of sideplates of *Gammarus duebeni* & *Gammarus* sp.
Askö rock pool, Kieler Bucht
- I. coniferum** Precht,1935
@ first pereiopods of *Gammarus locusta*
Kieler Bucht

- I. brachymyon** Precht,1935
@ between chaete of rear pereiopods of *Gammarus locusta*
Kieler Bucht
- I. longum** Precht,1935
@ between antennae of *Gammarus locusta*
Kieler Bucht
- I. sp. a :** Fenchel,1965
@ on oostegites of *Gammarus locusta* L.
Öresund
- Myoschiston** Precht,1935
- M. duplcatum** Precht,1935
@ *Carcinus maenas* & *Idothea* spp.
Kieler Bucht
- M. neomysidis** Precht,1935
@ *Neomysis vulgaris*
Kieler Bucht
- M. careini** Precht,1935
@ gills of *Carcinus maenas*
Kieler Bucht
- M. centropagidarum** Precht,1935
@ *Centropages hamatus* & *Eurytemora hirundoides*
Kieler Bucht
- M. balanorum** (Mereschkowsky,1877)
= *Epistylis balanorum* Mereschkowsky,1877
@ "gills of *Balanus improvisus*
Kieler Bucht
- M. cypridicola** Precht,1935
@ legs of *Cyprideis littoralis*
Kieler Bucht
- Zoothamnium** Bory de St. Vincent,1826
- Z. hiketes** Precht,1935
@ *Gammarus* spp. & *Jaera*
Askö, Kieler Bucht, Öresund, Isefjord
- Z. duplicatum** Kahl,1933
@ *Gammarus oceanicus* or on many other invertebrate species or freelifing
Askö, Kieler Bucht
- Z. rigidum** Precht,1935
@ on sideplates & pereiopods of *Gammarus oceanicus*
Kieler Bucht, Öresund, Isefjord
- Z. nanum** Kahl,1933
@ *Gammarus* spp. & various other crustacea
Kieler Bucht, Öresund, Isefjord
- Z. intermedium** Precht,1935
@ *Palaemon*, *Crangon*, *Microdeutopus*, *Ciona*, etc.
Kieler Bucht
- Z. laomedae** Precht,1935
@ *Laomedea loveni*
Kieler Bucht
- Z. abrae** Precht,1935
@ *Abra alba*
Stoller Grund
- Z. alternans** Precht,1935
@ *Crisia eburnea*
Kieler Bucht
- Z. vermicola** Precht,1935
@ *Eteone longa*, *Nereimyra punctata*, *Lagis koreni*, *Nephthys* sp., *Ophiura albida*
Stoller Grund, Kieler Bucht
- Z. sp. a :** Fenchel,1965
@ on antennae & sometimes pereiopods of *Gammarus locusta* L.
Öresund
- MOBIILIDA** Kahl,1933
- Leiotrochidae** Johnston,1938
- Leiotrocha** Fabre-Domergue,1888
- L. patellae** (Cünnot,1891)
= *Trichodina patellae* Cünnot,1891
@ gills of *Patella* spp., & a number of bivalves, e.g. *Thyasira sarsi* (Philippi), *Thyasira flexuosa* (Montagu),
Abra nitida (Miller) & *Abra prismatica* (Montagu)
* sibling species complex
Öresund, Læsø, Gullmarfjord, Roscoff
- Polyvyclidae** Poljansky,1951
- Polyycyla** Poljansky,1951
- Trichodinidae** Claus,1874
- Trichodina** Ehrenberg,1830
= *Cyclochaeta* Jackson,1875 = *Cyclocyrrha* Fabre-Domergue,1888
= *Anhymina* Fabre-Domergue,1888
@ on fish, especially on the gills
- T. macomarum** Raabe & Raabe,1959
@ *Macoma balthica* (L.)
Askö, Gdynia, Öresund
- T. sp. 1 :** Fenchel,1965
?= *Cyclochaeta cardii* Delphy,1938 (nomen nudum)
* *Cyclochaeta cardii* Delphy is from Arcachon
@ *Cerastoderma edule* (L.)
Öresund
- T. sp. 2 :** Fenchel,1965
?= *Cyclochaeta cardii* Delphy,1938 (nomen nudum)
@ *Cerastoderma edule* (L.)
Öresund
- T. sp. :** van Banning,1979
@ *Ostrea edulis*
Netherlands
- T. polandiae** Fenchel,1965
= *T. domergui* f. *cardii* Raabe & Raabe,1959
@ *Cerastoderma glaucum*
Askö, Poland, non Öresund, nec Gullmarfjord
- T. scolopontis** Precht,1935
@ *Scoloplos armiger*
Kieler Bucht
- T. cardiorum** Raabe & Raabe,1959
@ small *Cerastoderma glaucum*
Poland
- T. baltica** Quennerstedt,1869
@ *Hydrobia ventrosa* & *Neritina fluviatilis*
Visby, Polish Baltic, Kieler bucht
- T. terebellididis** Precht,1935
@ *Terebellides stroemi*
Stoller Grund
- T. serpularum** (Fabre-Domergue,1888)
= *Cyclochaeta serpularum* Fabre-Domergue,1888
@ *Serpula* spp., *Nephthys* sp., *Phyllodoce* sp., etc
Kieler Bucht, Concarneau
- T. astericola** (Precht,1935)
= *Cyclochaeta astericola* Precht,1935
@ *Asterias rubens*
Kieler Bucht
- T. myicola** Uzmann & Stickney,1954
@ *Mya arenaria* L.
E USA, non Baltic, nec SW Scandinavia
- T. halli** ??,1??
@ several fish species
New Jersey
- T. labrorum** Chatton,1910
@ Labrid fish gills
?France
- T. oviducti** Polyansky,1955
@ in urogenital tract of 5 sp. of gen. *Raja*
Atlantic Ocean
- T. puytoraci** Lom,1962
@ *Mugil* spp.
Black Sea
- T. rectuncinata** Raabe,1958
@ >15 fish species
Adriatic, Black Sea, Caribbean
- T. domerguei** Wallengren,18??
@ sticklebacks & other fresh or brackish water teleosts
* euryhaline fresh water species
Kieler Bucht
- Vauchomia** Mueller,1938
- Trichodinopsidae** Kent,1881
- Trichodinopsis** Claparède & Lachmann,1858
- Urecolariidae** Dujardin,1840
- Urecolaria** von Stein,1854
- U. elongata** Fenchel,1965
@ *Nuculana minuta* (Müller) & *Nuculana pernula* (Müller)
Öresund, Gullmarfjord
- ASTOMATIA** Schewiakoff,1896
@ often in lumen of digestive tracts of many annelids & some molluscs
- ASTOMATIDA** Schewiakoff,1896
- Anoplophryidae** Cépède,1910
- Almophrya** de Puytorac & Dragesco,1969
- Anoplophrya** von Stein,1860
@ endosymbionts of terrestrial oligochaetes
- A. nodulata** ??,1??
@ e.g. *Phallodrilus monospermathecus*, *Thalassodrilus prostatus*, *Marionina* sp.
Schlei
- A. polydorae** ??,1??
@ e.g. *Lumbricillus lineatus* & *Marionina* sp.
Schlei, Sylt
- A. mytili** Delphy,1938
@ mantle cavity & intestines of *Mytilus edulis*
* Dubious species.
Arcachon
- O. stellarum** Cépède,1907 (TM)
@ *Asterias rubens* A. forbesi & *A. vulgaris* (affects asteroid gonads and may cause total castration in male hosts)
Netherlands, Belgium, Engl. Channel, E USA
- Buetschliidae** de Puytorac, in Corliss,1979
- Buetschliida** Awerinzew,1908
- Clausilocolidae** de Puytorac, in Corliss,1979
- Clausilocola** Lom,1959
- Haptophryopsis** de Puytorac,1971
- Contophryidae** de Puytorac,1972

- Contophrya*** de Puytorac & Dragesco, 1968
- Dictonophrya*** de Puytorac & Dragesco, 1968
- Haptophryidae*** Cépède, 1910
- Cepedietta*** Kay, 1942
- Haptophrya*** von Stein, 1867
- Steinella*** Cépède, 1910
- Hoplitophryidae*** Cheissin, 1930
- Buetschliellopsis*** de Puytorac, 1954
- Delphyella*** de Puytorac, 1969
- Hoplitophrya*** von Stein, 1859
- Juxtaradiophrya*** de Puytorac, 1954
- Radioiphryoides*** Lom, 1956
- Intosellinidae*** Cépède, 1910
- Intosellina*** Cépède, 1910
- Maupasellidae*** Cépède, 1910
- Maupasiella*** Cépède, 1910
- Radiophryidae*** de Puytorac, 1972
- Acanthodiophrya*** de Puytorac & Dragesco, 1968
- Heletta*** Kaczanowski, 1961
- Metaracoelophrya*** de Puytorac & Dragesco, 1969
- Metaradiophrya*** Heidenreich, 1935
- Mrazeckiella*** Kijenskij, 1925
- Radiophrya*** Rossolimo, 1926
- Archiaستوماتidae*** Foissner, 1976
* Family incertae sedis
- Archiaستoma*** Foissner, 1976
- APOSTOMATIA*** Chatton & Lwoff, 1928
- APOSTOMATIDA*** Chatton & Lwoff, 1928
- Colliniidae*** Cépède, 1910
- Metacollinia*** Jankowski, 1980
- Cyrtocaryidae*** Cortiss, 1979
* Not included in Small & Lynn, 1985
@ lateral ceca of digestive tube in polychaetes
- Cyrtocaryum*** Fauré-Fremiet & Mugard, 1949
- Spirobuetschliella*** Hovasse, 1950
* Incertae sedis in this family
- S. chattoni*** Hovasse, 1950 (TM)
@ *Pomatoceros triquier*
Banyuls-sur-mer
- Foettingeriidae*** Chatton, 1911
- Foettingeria*** Caulley & Mesnil, 1903
- F. actiniarium*** ?Caulley & Mesnil, 1903? (TM)
@ various actinarians, e.g. *Metridium senile*, *Adamsia palliata* & *Sagartia troglodytes*
along French coasts
- Gymnodynionoides*** Minkiewicz, 1912
- G. inkystans*** Minkiewicz, 1912?
@ host tissue feeder in many Malacostraca, e.g. *Gammarus* spp.
Askö, Öresund, ?sefjord
- G.*** sp.
@ *Sagartia parasitica* from *Pagurus prideauxi* shells
Roscoff
- Hyalophysa*** Bradbury, 1966
- Ophiuraespira*** Chatton & Lwoff, 1930
@ Ophiuroids
- Pericaryon*** Chatton, 1911
@ Ctenophora
- P. cesticola*** Chatton, 1911 (TM - French Mediterranean)
@ trophonts in *Cestus veneris*; phoronts probably in copepods
Mediterranean
- Phitorophrya*** Chatton & Lwoff, 1930
@ other Apostomatines
- Phoretophysa*** Chatton & Lwoff, 1930
- Spirophrya*** Chatton & Lwoff, 1924
- S. subparasitica*** Chatton & Lwoff, 1924 (TM)
@ phoront attached to harpacticoid *Idya furcata*; trophonts in gastric cavity of *Cladonema radiatum*
Banyuls-sur-mer
- ASTOMATOPHORIDA*** Jankowski, 1966
- Opalinopsisidae*** Hartog, 1906
- Chromidina*** Gonder, 1905
@ Beside cephalopods, there is a first crustacean (Euphausiid, Copepod or other appropriate crustacean) host stage
- C. coronata*** (?Foettinger, 1881?)
- @ renal or renal-pancreatic coela of e.g. *Octopus vulgaris*, *Eledone cirrosa*, *Sepiola rondeletii* & *Illex coindetii*
Engl. Channel, Mediterranean
- C. elegans*** (?Foettinger, 1881?)
@ renal or renal-pancreatic coela of *Sepia elegans*, *S. orbignyiana* & *Illex coindetii*
Mediterranean
- Opalinopsis*** Foettinger, 1881
- O. sepioiae*** ?Foettinger, 1881?
@ midguts & digestive glands of *Sepiola rondeletii*, *Sepia elegans* & *S. officinalis*; probably is the same species observed from *Sepiola atlantica*, *Sepiella oceaniana* & *Rossia macrosoma*
Engl. Channel & Mediterranean
- O.*** sp.: Mohr (unpubl.), Hochberg, 1971
@ *Alloteuthis subulata*
Engl. Channel
- PILISUCTORIDA*** Jankowski, 1966
- Conidophryidae*** Kirby, 1941
- Ascophrys*** Campillo & Deroux, 1974
- Askoella*** Fenchel, 1965
- A. janssoni*** Fenchel, 1965 (TM - Askö)
@ *Gammarus* spp.
Askö, Öresund
- Conidophrys*** Chatton & Lwoff, 1934
- C. pilisuctor*** Chatton & Lwoff, 1934
@ host tissue feeder in many Malacostraca, e.g. *Gammarus* spp., *Limnoria*, *Corophium*, *Idothea*
Öresund, Isefjord, France, California
- PLAGIOPYLIA*** Small & Lynn, in Lee, Hutner & Bovee, 1985
- PLAGIOPYLIDA*** Small & Lynn, in Lee, Hutner & Bovee, 1985
- Plagiopylidiae*** Schewiakoff, 1896
- Lechiropyla*** Lynch, 1930
- Plagiopyla*** von Stein, 1860 (TM *P. nasuta* von Stein, 1860 - fresh water)
- P. cucullio*** Quennerstedt, 1867
= *Paramaecium cucullio* Quennerstedt, 1867
= *Plagiopyla nasuta* v. *marina* Gourret & Roeser, 1886
* among rotting green algae
Öresund, Halland, Mediterranean
- P. frontata*** Kahl, 19??
Öresund, Isefjord, Kattegatt, Limfjorden
- P. ovata*** Kahl, 19??
S Baltic
- P. vestita*** Kahl, 19??
?
- P. minuta*** ??, 1??
@ found on *Strongylocentrotus droebachiensis* by Beers, 1952
- Pseudoplagiopyla*** Small & Lynn, in Lee, Hutner & Bovee, 1985
- Sonderidae*** Small & Lynn, in Lee, Hutner & Bovee, 1985
- Parasonderia*** Fauré-Fremiet, 1973
- Sonderia*** Kahl, 1928
- S. sinuata*** Kahl, 19??
Öresund
- S. schizostoma*** Kahl, 19??
Askö, Öresund
- S. vorax*** Kahl, 19??
Askö, Öresund, Isefjord
- S. cyclostoma*** Kahl, 19??
Öresund
- S. labiata*** Fauré-Fremiet & Tuffrau, 1955
Concarneau
- S. mira*** Kahl, 19??
?
- S. vestita*** Kahl, 19??
?
- S. macrochilus*** Kahl, 19??
?
- S. pharyngea*** Kirby, 1934?
- COLPODEA*** de Puytorac & al., 1974
- CYRTOLOPHOSIDIDA*** Foissner, 1978
- Cyrtolophosididae*** Stokes, 1888
- Cyrtolophosis*** Stokes, 1885
- Grossglockneriidae*** Foissner, 1980
- Grossglockneria*** Foissner, 1980
- Woodruffiidae*** von Gelei, 1954
- Enigmostoma*** Jankowski, 1975
- Platypophrya*** Kahl, 1926
= *Telostoma Grandori*, 1935, non Ørsted, 1843 (Plathelm.). nec von Graff, 1882 (Plathelm.)
- P. ferroii*** (R & L. Grandori, 1935?)
= *Telostoma ferroii* R & L. Grandori, 1935?
Plymouth

Rhyposophrya Kahl,1933
* incertae sedis

R. aplanata Kahl,1933
Helgoland

Rostrophrya Njiné,1978

Woodruffia Kahl,1931

BRYOPHRYIDA de Puytorac & al.,1979

Bryophryidae de Puytorac & al.,1979

Bryophrya Kahl,1931

Puytoraciella Njiné,1978

COLPODIDA de Puytorac & al.,1974

Colpodidae Ehrenberg,1838

Bresslaua Kahl,1931

Colpoda O.F. Müller,1773

Tillina Gruber,1879

Marynidae Poche,1913

Maryna Gruber,1879

Mycterothrix Lauterborn,1898

BURSARIOMORPHIDA Fernández-Galiano,1978

Bursariidae Dujardin,1840

* predominantly fresh water forms

Bursaria O.F. Müller,1773