

NEAT (North East Atlantic Taxa):

South Scandinavian marine "Aschelminth" (except Nematoda, which are treated separately) & some other species poor phyla Check-List.

compiled at TMBL (Tjärnö Marine Biological Laboratory) by:

Hans G. Hansson 1988-11-16 / small revisions until Sep. 1996, when it - somewhat revised (and again May 1997) - was republished as a pdf file on Internet.

The following phyla are treated in this list: **Placozoa, Gnathostomulida, Tardigrada, Entoprocta, Cyclophora, Priapulida, Kinorhyncha, Loricifera, Mesozoa, Acanthocephala, Nematomorpha, Gastrotricha, Rotifera, Echiura, Sipuncula, Pogonophora, Hemichordata, Chaetognatha & Xenoturbellida.**

Citation suggested: Hansson, H.G. (Comp.), 1997. NEAT (North East Atlantic Taxa): South Scandinavian marine "Aschelminth" (excl. Nematoda) & other species poor phyla Check-List. Internet pdf Ed., May 1997. [<http://www.tmbi.gu.se>].

Denotations: (TM) = Genotype @ = Associated to * = General note

N.B.: This is one of several preliminary check-lists, covering S. Scandinavian marine animal (and partly marine protist) taxa. Some financial support from (or via) NKMB (Nordiskt Kollegium för Marin Biologi), during the last years of the existence of this organisation (until 1993), is thankfully acknowledged. The primary purpose of these checklists is to facilitate for everyone, trying to identify organisms from the area, to know which species that earlier have been encountered there, or in neighbouring areas. A secondary purpose is to facilitate for non-experts to find as correct names as possible for organisms, including names of authors and years of description. So far these checklists are very preliminary. Due to restricted access to literature there are (some known, and probably many unknown) omissions in the lists. Certainly also several errors may be found, especially regarding taxa like Plathelminthes and Nematoda, where the experience of the compiler is very rudimentary. The credibility of some other lists, containing taxa like e.g. Porifera, where, at least in certain subtaxa, taxonomic confusion seems to prevail, is certainly also suboptimal so far.

Species found in South Scandinavia, as well as from neighbouring areas, chiefly the British Isles, have been considered, as some of them may show to have a slightly more northern distribution, than what is known today. However, species with a typical Lusitanian distribution, with their northern distribution limit around France or Southern British Isles, have as a rule been omitted here, albeit a few species with probable northern limits around the British Isles are listed here until distribution patterns are better known. The compiler would be very grateful for every correction of presumptive lapses and omissions an initiated reader could make.

N.B. This document is intended to be completely unavailable regarding nomenclature within the meaning of ICZN.

PLACOZOA

Treptopanax Monticelli,1893

T. reptans Monticelli,1896

* Possibly a synonym of **Trichoplax adhaerens** von Schultze
Aquarium tanks of the Plymouth laboratory, Mediterranean

Trichoplax F.E. von Schultze,1883

T. adhaerens F.E. von Schultze,1883

Mediterranean

GNATHOSTOMULIDA Ax,1956

FILOSPERMOIDEA Sterrer,1972

Haplognathiidae Sterrer,1972

Haplognathia Sterrer,1970

H. ruberrima (Sterrer,1966)
Öresund, Gullmarfjord

H. filum (Sterrer,1966)
Öresund, Gullmarfjord, Sylt

H. simplex (Sterrer,1966)
Öresund, Gullmarfjord, Sylt

H. rubromaculata (Sterrer,1969)
Gullmarfjord, Sylt

H. rosea (Sterrer,1969)
Gullmarfjord, Sylt

H. gubbarnorum (Sterrer,1969)
Gullmarfjord, Sylt

H. lunulifera (Sterrer,1969)
Gullmarfjord

Pterognathiidae Sterrer,1972

Pterognathia Sterrer,1966

P. swedmarki Sterrer,1966
Öresund, Gullmarfjord, Sylt

P. atros Sterrer,1966
Gullmarfjord

P. meixneri Sterrer,1969
Gullmarfjord

P. sorex Sterrer,1969
Gullmarfjord, Sylt

Pterognathia spp.
* 3 more species known from Scandinavia?

BURSOVAGINOIDEA Sterrer,1972

SCLEROPERALIA Sterrer,1972

Agnathiliidae Sterrer,1972

Agnathiliella Sterrer,1971
Florida

Clausognathiidae Sterrer,1992

Clausognathia Sterrer,1992 (TM)**Clausognathia suicauda** Sterrer,1992 - Belize

Mesognathariidae Sterrer,1972

Mesognatharia Sterrer,1966
Bohuslän?, Ireland

Labidognathia Riedl,1970
Florida, Panama

Tenuignathia Sterrer,1976
Florida

Gnathostomariidae Sterrer,1972

Gnathostomaria Ax,1956
Öresund, Mediterranean

Rastrognathiidae Kristensen & Nørrevang,1977

Rastrognathia Kristensen & Nørrevang,1977

R. macrostoma Kristensen & Nørrevang,1977
S Kattegatt

Problognathidae Sterrer & Farris,1975

Problognathia Sterrer & Farris,1975
Bermuda

Onychognathiidae Sterrer,1972

Onychognathia Riedl,1971
Florida

Valvognathia Kristensen & Nørrevang,1978

V. pogonostoma Kristensen & Nørrevang,1978
Öresund

Nanognathia Sterrer,1973
Florida

Gnathostomulidae Sterrer,1972

Gnathostomula Ax,1956

G. paradox Ax,1956
Öresund, Bohuslän, Kiel, Helgoland, Sylt, Bergen

Semaeognathia Riedl,1970
Florida

Corculognathia ?,19??
?

CONOPHORALIA Sterrer,1972

Austrognathiidae Sterrer,1972

Austrognathia Sterrer,1965
Öresund, Atlantic coast of France, Mediterranean

Austrognatharia Sterrer,1971
Ireland, Mediterranean

TARDIGRADA

= **IL TARDIGRADO** Spallanzani,1777

* Not all terrestrial species from the marine splash zone are included.

HETEROTARDIGRADA Marcus,1927

ARTHROTARDIGRADA Marcus,1927

Halechiniscidae Thulin,1928

Archechiniscinae

Archechiniscus Schulz,1953

A. marci Schulz,1953
Caribbean, S & E Pacific

Tanartinae Renaud-Mornant,1980

Tanarctus Renaud-Debyser,1959

T. arborepinosus Lindgren,1971
W Atlantic

T. longisetosus Grimaldi de Zio, D'Addobbo Gallo, Morone De Lucia, Vaccarella & Grimaldi,1982
Mediterranean

T. ramazzottii Renaud-Mornant,1975
Bretagne

T. tauricus Renaud-Debyser,1959
Mediterranean, Florida

T. gracilis Renaud-Mornant,1980
Koster area (R.M. Kristensen pers. comm.), North Carolina

T. dendriticus Renaud-Mornant,1980
North Carolina

T. heterodactylus Renaud-Mornant,1980
North Carolina

Actinarctus Schulz,1935*A. doryphorus doryphorus* Schulz,1935 (TM - Helgoland)@ on body surface of *Echinocyanus pusillus*, but a common habitat is - according to Kristensen (pers. comm.) - probably the only connection between the tetrigrade and the urchin.
Gulmarfjord area (Bonden), Koster area (Kalkgrund), Helgoland*A. doryphorus ocellatus* Renaud-Mornant,1971
Roscoff, Brest*A. neretus* Grimaldi de Zio, D'Addobbo Gallo, Morone De Lucia, Vaccarella & Grimaldi,1982
Mediterranean*A. physophorus* Grimaldi de Zio, D'Addobbo Gallo, Morone De Lucia, Vaccarella & Grimaldi,1982
Mediterranean*Zioella* Renaud-Mornant,1987*Florarctinae* Renaud-Mornant,1982*Florarctus* Delamare Deboutteville & Renaud-Mornant,1965*F. antillensis* van der Land,1968
Caribbean*F. acer* Renaud-Mornant,1990
France*F. cinctus* Renaud-Mornant,1976
Roscoff, Mediterranean*F. hulingsi* Renaud-Mornant,1976
Mediterranean*Ligarctus* Renaud-Mornant,1982*Wingstrandarctus* Kristensen,1984 (TM *W. corallinus* Kristensen,1984 - E Australia)*W. intermedius* (Renaud-Mornant,1967) Kristensen,1984
= *Halechiniscus intermedius* Renaud-Mornant,1967
Mediterranean, tropical seas*W. sp. :* Kristensen,1984
Tromsø*W. sp. :* Kristensen,1984
Roscoff*Halechiniscinae* Thulin,1928*Chrysoarctus* Renaud-Mornant,1984*Halechiniscus* Richters,1908*H. flabellatus* Grimaldi de Zio, D'Addobbo Gallo, Morone De Lucia, Vaccarella & Grimaldi,1982
Mediterranean*H. guiteli* Richters,1908 (TM - Bretagne)
@ *Ostrea edulis*.
* A species incerta!
Helgoland, Engl. Channel, Bretagne*H. greveni* Renaud-Mornant & Deroux,1976
Roscoff*H. perfectus* Schulz,1955
North Sea, Brest at 170m, Scotland*H. remanei* Schulz,1955
Gulmarfjord area (Bonden), Brest at 130m, Florida*H. subterraneus* Renaud-Debyser,1959
Mediterranean*Paradoxipus* Kristensen & Higgins,1989 (TM *Paradoxipus orzeliscooides* Kristensen & Higgins - Florida)*Orzeliscinae**Orzeliscus* du Bois-Reymond Marcus,1952*O. belopus* du Bois-Reymond Marcus,1952 (TM)
Scotland (Firemore Bay within Loch Ewe), Arcachon*O. septentrionalis* Schulz,1953
German North Sea coast*Opydorscus* Renaud-Mornant,1989*Styraconyxinae* Renaud-Mornant & Kristensen,1983*Angura* Pollock,1979*Bathyechiniscus* Steiner,1926 (TM *Bathyechiniscus tetronyx* Steiner,1926 - Antarctic Seas)*Lepoarctus* Kristensen & Renaud-Mornant,1983 (TM *Pleocola conifera* Renaud-Mornant,1975)
* Correct subfamily?*Paratanarctus* D'Addobbo Gallo, De Zio Grimaldi, Morone D Lucia & Troccoli,1992*Pleocola* Cantacuzène,1951*P. limnoriae* Cantacuzène,1951 (TM)
@ Sometimes associated with *Limnoria lignorum*.
Roscoff*Raiarctus* Renaud-Mornant,1981*R. colorus* Renaud-Mornant,1981
Florida, Caribbean Sea*Rhomboarctus* Renaud-Mornant,1984*Styraconyx* Thulin,1942*S. hallasi* Kristensen,1977
W Greenland warm springs, Mediterranean*S. haploceros* Thulin,1942 (TM - St. Malo)@ *Lichina*-tufts.*S. kristensenii* Renaud-Mornant,1981
Mediterranean, tropical seas*S. nanoqsunguak* Kristensen & Higgins,1984
off Disko Island (W Greenland) in 100-110 m, Mediterranean*S. givitoq* Kristensen & Higgins,1984
@ found on a Bryozoan colony
Disco Bay (W Greenland) in 70 m, Mediterranean*S. sargassi* Thulin,1942
= *Bathyechiniscus tetronyx* : Marcus,1936, non Steiner
@ on the invasive seaweed *Sargassum* spp.
Mediterranean, Azores*S. paulae* Robotti,1971
?*S. craticulus* (Pollock,1983)
= *Bathyechiniscus craticulus* Pollock,1983*Tetraekentron* Cuénot,1892*T. synaptae* Cuénot,1892 (TM - Roscoff)
@ Parasite on tentacles of *Leptosynapta gallienae*
Roscoff*Tholoarctus* Kristensen & Renaud-Mornant,1983
* Correct subfamily?*T. natans* Kristensen & Renaud-Mornant,1983 (TM - Roscoff)

* Semi-pelagic on "Dentalium -sand"

Roscoff

Euclavarciniae Renaud-Mornant,1983*Clavarctus* Renaud-Mornant,1983*Euclavaretus* Renaud-Mornant,1975 (TM *Euclavaretus thieli* Renaud-Mornant,1975 - deep Indian Ocean)*Exoclavaretus* Renaud-Mornant,1983*Moebjergarctus* Bussau,1992*Parmursa* Renaud-Mornant,1984*Proclavaretus* Renaud-Mornant,1983*Batillipedidae* Riggan,1962*Batillipes* Richters,1909*B. acaudatus* Pollock,1971
Yorkshire (Robin Hood's Bay)*B. annulatus* De Zio,1962
Mediterranean*B. bullicaudatus* McGinty & Higgins,1968
Scotland (Loch Ewe) (see McIntyre & Murison, 1973), Florida, Virginia, Massachusetts*B. carnonensis* Fize,1957
Mediterranean, Indian Ocean*B. dicrocercus* Pollock,1970
Woods Hole*B. friaufi* Riggan,1962
W Atlantic*B. littoralis littoralis* Renaud-Debyser,1969
Arcachon, Loch Ewe? (May have been the other subspecies)*B. littoralis submersus* d'Hondt,1970
Roscoff*B. mirus* Richters,1909 (TM - Kieler Bucht)
* Old distribution data regarding this species should probably be considered
with some care. It may have been confused with species described later.
Kieler Bucht - Bohuslän, W. Denmark, Bergen, Tärminne??, Britain, cosmopolite*B. noerrevangi* Kristensen,1978
Denmark (Nivå Bay, E Sjælland)*B. pennaki* Marcus,1946
Arcachon, Mediterranean, Massachusetts*B. phreaticus* Renaud-Debyser,1959
Elbe estuary, Britain (e.g. Yorkshire), Scilly, Arcachon*B. roscoffensis* Kristensen,1978
@ *Leptosynapta gallienae*.
Roscoff*B. similis* Schulz,1955
Mediterranean*B. tubernatis* Pollock,1971
Elbe estuary, Yorkshire, Scotland*Stygarctidae* Schulz,1951*Stygarctinae* Schulz,1951*Megastygarctides* McKirdy & al.,1976 (TM *M. orbiculatus* McKirdy & al.,1976)*M. setolosa* Morgan & O'Reilly,1988
NW Scotland*Mesostygarctus* Renaud-Mornant,1979

M. intermedius Renaud-Mornant,1979
Mediterranean

Parastygarcus Renaud-Debyser,1965 (TM *Parastygarcus higginsi* Renaud-Debyser,1965 - Madagascar)

P. sterri Renaud-Mornant,1970
Adriatic Sea, Florida

Pseudostygarcus McKirdy & al.,1976

Stygarctus Schulz,1951

S. bradypus Schulz,1951 (TM - Sylt)
Sylt, Roscoff

S. granulatus Pollock,1970
Woods Hole

S. gourbaultae Renaud-Mornant,1981
Florida, Caribbean Sea

Neoractinae

Neoractus Grimaldi de Zio, D'Addabbo Gallo & Morone De Lucia,1992

N. primigenius Grimaldi de Zio, D'Addabbo Gallo & Morone De Lucia,1992 (TM - Tyrrhenian Sea)
Mediterranean

Renaudarctidae Kristensen & Higgins,1984

Renaudarctus Kristensen & Higgins,1984 (TM *Renaudarctus psammocryptus* Kristensen & Higgins,1984 - Florida)

Neostygartinae? ,19??

Neostygarcus Grimaldi de Zio, D'Addabbo Gallo & Morone De Lucia,1982

N. acanthophorus Grimaldi de Zio, D'Addabbo Gallo & Morone De Lucia,1982 (TM - Mediterranean)
Mediterranean (in marine cave)

Coronarctidae Renaud-Mornant,1974

Coronarctus Renaud-Mornant,1974 (TM *Coronarctus tenellus* Renaud-Mornant,1974 - deep Indian Ocean)

C. stylisetus Renaud-Mornant,19??
Faroe Bank in 104-209 m, Cape Hatteras in 439 m

C. laubieri Renaud-Mornant,1987
Biscay

Trogloarctus Villora-Moreno,1996

T. trionyches Villora-Moreno,1996 (TM - Mediterranean submarine cave)
Mediterranean

ECHINISCOID(E)A Marcus,1927

Echiniscoididae Kristensen & Hallas,1980
Bohuslän, Bergen, Helgoland

E. sigismundi galliensis Kristensen,1980
Eng. Channel

E. sigismundi groenlandicus Kristensen,1980
Greenland

E. sigismundi hispaniensis Kristensen,1980
Biscay

Anisonyches Pollock,1975 (TM *Anisonyches diakidius* Pollock,1975 - Bahamas)

Echiniscoidea Plate,1889

E. sigismundi sigismundi (Max Schultze,1865)
= *Echiniscus sigismundi* Max Schultze,1865 (TM)

@ Associated with *Semibalanus balanoides* and *Mytilus edulis*.

E. sigismundi mediterranicus Kristensen,1980
W Mediterranean

E. hoepneri Kristensen,1980
@ *Semibalanus balanoides*
Greenland

Echiniscidae

Echiniscus C.A.S. Schultze,1840

E. quadrispinosus Richters,1902
* A moss waterfilm species recorded once in the sea.
Arcachon

Bryodelphax Thulin,1928 (TM *Echiniscus intermedius* Murray,1910)

B. parvulus Thulin,1928
@ May be found supralitorally on the lichen *Anaptychia*
?

EUTARDIGRADA Marcus,1927

PARACHELA

Macrobiotidae Thulin,1928

Macrobiotus Schultze,1834 (TM *Macrobiotus Hufelandi* Sigismund Schultze,1834)

M. echinogenitus Richters,1903
= *M. crenulatus* Richters,1903
@ May be found supralitorally on the lichen *Anaptychia*
?Bohusl.

Hypsibiidae Pilato,1969

Hypsibiinae Pilato,1969

Halobiotus R.M. Kristensen,1982

* This is the only really marine genus of Eutardigrada

H. crispae R.M. Kristensen,1982 (TM)
Bohuslän, Disco Island, Arctic Sea

H. stenostomus (Richters,1908)

= *Isohypsistius stenostomus* (Richters,1908)

= *Hypsibius stenostomus* (Richters,1908)

= *Macrobiotus stenostomus* Richters,1908

Fehmarn-Belt, Schleimünde, Black Sea

H. appelloefi (Richters,1908)

* according to Kristensen (pers. comm.) is this probably only a form of *H. stenostomus* (Richters)

= *Isohypsistius appelloefi* (Richters,1908)

= *Hypsibius appelloefi* (Richters,1908)

= *Macrobiotus appelloefi* Richters,1908

Bohuslän, Öresund, Bergen, Kieler Föhrde

H. geddesi (Hallas,1971)

= *Isohypsistius geddesi* (Hallas,1971)

= *Hypsibius (Isohypsistius) geddesi* Hallas,1971

Norway (Tromsö), Denmark (Frederikshavn)

H. arcturulus Crisp & R.M. Kristenssen,1983

Greenland

Hypsibius Ehrenberg,1848

H. dujardini (Doyère,1840)

= *Macrobiotus Dujardini* Doyère,1840

@ may be found in the upper littoral spray zone on the maritime rock moss *Schistidium maritimum* ?

Isohypsistius Thulin,1928

I. prosostomus Thulin,1928

@ May be found supralitorally on the lichens *Anaptychia*, *Ramalina* & *Xanthoria*
Bohusl.

Ramazzottius Binda & Pilato,1986

R. Oberhaeuseri (Doyère,1840)

= *Hypsibius (Hypsibius) oberhaeuseri* (Doyère,1840)

= *Macrobiotus Oberhaeuseri* Doyère,1840
@ May be found supralitorally on the lichens *Anaptychia*, *Ramalina* & *Xanthoria*
Britain

APOCHELA

Milnesiidae

Milnesium Doyère,1840

M. tardigradum Doyère,1840

@ May be found supralitorally on the lichens *Anaptychia*, *Ramalina* & *Xanthoria*
Baltic - Bohusl., Europe

ENTOPROCTA Nitsche,1869

= KAMPTOZOA Cori,1929

Loxosomatidae Hincks,1880

Loxosoma Keferstein,1863, emend. Mortensen,1911

= *Cyclatella* van Beneden & Hesse,1864

L. saltans Assheton,1912
@ in the tube of *Leiochone clypeata*.
Scotland (Isle of Skye)

L. loxalina Assheton,1912

@ in an unidentified Maldanide tube
W Scotland

L. singulare Keferstein,1862 (TM)

@ *Notomastus latericeus*.

English Channel

L. annelidicola (van Beneden & Hesse,1864)

= *Cyclatella annelidicola* van Beneden & Hesse,1864

@ *Nicomache lumbricalis* & *Petaloprotus terricola*.

Eng. Channel, Wales

L. clapedii Bobin & Prenant,1953

@ *Notomastus latericeus*.

N Kattegatt, Roscoff

L. monensis Eggleston,1965

@ inside unidentified arenaceous worm-tube

Isle of Man

L. pectinaricola Franzén,1962

@ *Pectinaria belgica*.

Öresund - Koster Area - Oslofjord - Skagerrak - Bergen, Cullercoats

L. rhodinicola Franzén,1962

@ *Rhodine loventi* (chiefly on the anterior segments)

S Kattegatt - Koster Area - Oslofjord - Skagerrak - Bergen, North Sea

"*L.*" *loricatum* Harmer,1915

* As the type specimens are described from the Indonesian bryozoan *Canda*, the European material, found by Bobin & Prenant, 1953, probably belongs to another species, possibly a

Loxosomella.

@ *Hermonia hystrix*.

Roscoff

Subgen. *Loxomorpha* Nielsen,1964

L. (Loxomorpha) significans Nielsen,1964

@ on the anterior end of *Achyris biceps*.

Kattegatt - Koster Area - Oslofjord - Bergen

L. (Loxomorpha) agile Nielsen,1964
@ in the tubes of *Nicomache trispinata*.
Kattegatt - Bohuslän - Bergen

Loxomespilon Bobin & Prenant,1953
* Sensu Emschermann, 1971 is this genus closely related to subgenus
Loxomitra within *Loxosomella*.

L. perezi Bobin & Prenant,1953 (TM)
@ on *Sthenelais boa*.
Plymouth, Roscoff

Loxosomella Mortensen,1911
= *Loxocalyx* Mortensen,1911

L. antedonis Mortensen,1911
* The Swansea record by Ryland & Austin,1960 is probably not the same species as Mortensen found on a crinoid from Greenland.
@ on *Poliomethra prolixa*, Swansea, on settling panels.
Greenland

L. atala (Barrois,1877)
* Doubtful species
@ on a sponge, *Dysidea*?
English Channel

L. brumpti (Nilus,1909)
@ *Phascolion spitzbergense*.
Kolafjord

L. fagei Bobin & Prenant,1953
@ on ventral side of *Hermonia hystrix*.
Roscoff

L. elegans Nielsen,1964
@ in the tubes of *Praxillella praetermissa*, *Rhodine gracilior*, *Heteroclymene robusta* &
Nicomache lumbinalis
Öresund - Koster Area, Bergen

L. similis Nielsen,1964
@ in the tubes of *Nicomache trispinata*.
Kattegatt - Bohuslän, Bergen

L. polita Nielsen,1964
@ in the tubes of *Petaloprotus tenuis* var. *borealis*, *Nicomache lumbinalis* and (more rarely)
Rhodine gracilior.
Öresund - Koster Area, Bergen

L. ornata Nielsen,1964
@ in the tubes of *Thelepus cincinnatus*
Öresund - Bohuslän

L. marsypos Nielsen & Ryland,1961
@ in the tubes of *Eunice pomata* and *Chaetopterus* sp..
Skagerrak, Bergen, Isle of Man

L. harmeri (Schultz,1895)
@ *Gattyana cirrosa* and *Harmothoe* spp.
Öresund - Kattegatt, Culvercoats, White Sea

L. atkinsae Bobin & Prenant,1953
= *L. bouxi* Bobin & Prenant,1953
=? *L. cuenoti* Bobin & Prenant,1953
= *L. arvayae* Bobin & Prenant,1953
= *L. fungiformis* Bobin & Prenant,1953
@ on *Phascolion strombi* and in it's tube
Kattegatt - Bohuslän, Isle of Man, English Channel, France

L. nitschei (Vigelius,1882)
= *Loxosoma nitschei* Vigelius,1882
@ on different Bryozoaans, Hydroids, Algae and stones
Öresund - Bohuslän, Bergen, France, Culvercoats, Arctic Seas

L. murmanica (Nilus,1909)
= *Loxosoma murmanica* Nilus,1909
@ on *Phascolion strombi* and in it's shelter and inside the oldest parts of *Amphicteone auricomata*
tubes.
Öresund - Skagerrak, Kolafjord, Engl. Channel, W France, Isle of Man

L. phascolosomata (Vogt,1876)
@ on posterior end of *Golfingia vulgare* & *Golfingia elongata* and on their commensal bivalves
Mysella bidentata & *Epilepton clarkei*
Kattegatt - Koster Area, W Norway, Helgoland, Plymouth, SW Ireland

L. fauveli Bobin & Prenant,1953
@ *Aphrodita aculeata* & *Laemonice filicornis* & *Hermonia hystrix*.
N Kattegatt, Roscoff, England

L. teissieri (Bobin & Prenant,1953)
@ *Dysidea fragilis*.
Roscoff

L. compressa Nielsen & Ryland,1961
@ dorsal notopodial setae of various Polynoidae, e.g. *Gattyana cirrosa* & *Lagisca extenuata*
Kattegatt - Koster Area, Bergen, Isle of Man

L. nordgaardi Ryland,1961
@ different Bryozoans.
Kattegatt - Koster Area, Bergen, Culvercoats

L. glandulifera Franzén,1962
@ inside the tube of *Panthalis oerstedi*
Kattegatt - Bohuslän

L. discopoda Nielsen & Ryland,1961
@ *Amphilepis norvegica*
Bergen, Norwegian Skagerrak Coast

L. scaura Nielsen,1964
@ *Nephrys paradoxo*, *N. incisa* & *N. hombergi*.
Kattegatt - Koster Area

L. varians Nielsen,1964
@ various Nephthyidae.
W Baltic - Bohuslän in >20m, Bergen, Iceland

L. aripes Nielsen,1964
@ *Laonice cirrata*.
Bergen

L. obesa (Atkins,1932)
= *Loxosoma obesum* Atkins,1932
@ *Aphrodita aculeata*.
Bergen, Plymouth, Isle of Man

L. brucei Eggleston,1965
@ inside unidentified arenaceous polychete tube.
Isle of Man

L. claviformis (Hincks,1880)
= *Loxosoma claviformis* Hincks,1880
@ *Aphrodita & Laemonice*.
Isle of Man, Roscoff, Monaco

L. crassicauda (Salensky,1877)
= *Loxosoma crassicauda* Salensky,1877
@ on a diversity of invertebrates, e.g. *Phyllochaetopterus*, in tubes of *Sabella*, *Branchiomma*,
Bispira, on *Ciona*, *Zoothryon*, etc.
Plymouth, Naples

Subgen. **Loxomitra** Nielsen,1964

L. (Loxomitra) kefersteinii (Claparède,1867)
= *Loxosoma kefersteini* Claparède,1867
@ on *Bugula* and settling panels.
Swansea, Naples, Red Sea, Florida

Pedicellinidae Smit,1867

Pedicellina M. Sars,1835

P. nutans Dalyell,1848
@ no specific substrata demanded.
Frederikshavn, Swedish Westcoast, S North Sea, Plymouth, W Norway, Wales, Scotland

P. hispida Ryland,1965
@ on *Terebraella*
Plymouth, France, on *Bowerbankia*, New Zealand

P. cernua (Pallas,1771)
= *Brachionus cernuus* Pallas,1771
= *Hydra coronata* Fleming,1828
= *Pedicellina echinata* M. Sars,1835 (TM)
= *Pedicellina glabra* Dalyell,1848
= *Crinomorpha belgica* van Beneden,1845
= *Pedicellina americana* Leidy,1855
@ no specific substrata demanded.
W Baltic, Isefjord, Jutland, Bohuslän, Bergen, "Cosmopolitic"

Barentsiidae
= *Urnatellidae* Annandale,1915

Barentsia Hincks,1880
= *Pseudopedicellina* Toriumi,1951

B. gracilis (Sars,1835)
= *Pedicellina gracilis* M. Sars,1835
= *Crinomorpha belgica* van Beneden,1845
@ on inanimate substrata or e.g. on ventral side of *Aphrodita aculeata*, on upper surface of elytrae of *Lepidonotus squamatus*, on shells inhabited by *Phascolion* and on several other invertebrates.
W Baltic - Koster Area, Norway, "Cosmopolitic"

B. benedeni (Foëtinger,1886)
= *Pedicellina benedeni* Foëtinger,1886
W Baltic, Ostende, Hull, Mediterranean, Cape Cod, Japan

B. elongata Julian & Calvet,1903
@ on *Flustra* and various hard substrata.
Skagerrak, W North Sea, Helgoland, Mediterranean, Newfoundland

B. matsuoshimana Toriumi,1951
@ on various living or non-living substrata.
W Baltic, N Öresund - Skagerrak, Isle of Man, U.S. West coast, Japan

B. ramosa (Robertson,1900)
Puget Sound - California, Harbour of Doel (near Antwerpen) 1962, but not found afterwards
there or in any other place in Europe.

B. major Hincks,1888

Spitzbergen

B. bulbosa Hincks,1880 (TM)

Barents Sea

B. variarticulata K.A. Andersson,1902
Franz-Josef's Fjord

B. laxa Kirkpatrick,1890

* This species has frequently been reported from Northern Europe, but all identifications have been found to be wrong, so it should really vanish from lists like this.

B. mutabilis (Toriumi,1951)

* Doubtful species. Sensu P. Emschermann may odd growth forms of other species be hidden behind this name.

Wales, Japan

B. stiria Jullien & Calvet,1903

* An unrecognizable species.

Gulf of Gascone in 135 m.

Urnatella Leidy,1851 (TM) *U. gracilis* Leidy,1851 - fresh water

* Fresh water taxon.

Loxokalyptidae Emschermann,1972

Loxokalypus Emschermann,1972 (TM) *L. socialis* Emschermann,1972 - N Pacific

CYCLIOPHORA P. Funch & R. Møbjerg Kristensen,1995

EUCYCLIOPHORA P. Funch & R. Møbjerg Kristensen, 1995SYMBIIDA P. Funch & R. Møbjerg Kristensen, 1995Symbiidae P. Funch & R. Møbjerg Kristensen, 1995Symbion P. Funch & R. Møbjerg Kristensen, 1995S. pandora P. Funch & R. Møbjerg Kristensen, 1995 (TM)@ mouth parts & their setae of *Nephrops norvegicus*
Kattegatt, Gullmarfjord, FaeroesS. sp. : R.M. Kristensen (pers.comm.)@ *Homarus gammarus* & sporadically on *Nephrops norvegicus** This species has almost the double size of *S. pandora*.

Scandinavia

S. sp. : R.M. Kristensen (pers.comm.)@ *Homarus americanus*E North America (& on *H. americanus* transferred to European seas)**PRIAPULIDA** Théel, 1906PRIAPULA (4 Scandinavian spp.)PRIAPULIMORPH(II)A Salvini-Plawen, 1974Priapulidae Gosse, 1855Priapulus de Lamarck, 1816 (TM)= *Priapus* Linnaeus, 1758 (p.p.)

Bohuslän - S Baltic, all Norway, Britain

Priapulopsis Koren & Danielssen, 1875P. bicaudatus (Danielssen, 1868)= *Priapulus bicaudatus* Danielssen, 1868 (TM)

Norwegian Sea, N Troms - NE Finnmark, Arctic region

Tubiluchidae van der Land, 1970Tubiluchus van der Land, 1968 (TM *T. corallicola* van der Land, 1968 - Curaçao)T. arcticus Adrianov, Malakhov, Tchesunov & Tzetlin, 1989

* Described from females only.

The mouth of the White Sea

T. sp.* Only females found so far. According to Kristensen (pers. comm) the specimens from this population look more like animals from Bermudas than those from the White Sea. Furthermore Parra & Benito 1994 found larvae of this genus in Biscay.
Bohuslän, in shell-gravel (Kalkgrund off Nord-Koster)HALICRYPTOMORPH(HI)A Adrianov & Malakhov, 1995Halicyptidae Salvini-Plawen, 1974Halicyptus v. Siebold, 1849H. spinulosus v. Siebold, 1849 (TM)

Baltic - Öresund, Kungsbackafj., Iddefjorden, Trondheim, The Arctic

SETICORONARI(D)A Salvini-Plawen, 1974

* Not represented in the area.

KINORHYNCHA Reinhard, 1887ECHINODERA Gosse, 1864

* Several larval forms were once described as separate species. Many of those have not yet been correlated with adults, and perhaps never will. Names of such larvae are included here for the sake of completeness.)

CYCLORHAGIDA Zelinka, 1896CYCLORHAGAE Zelinka, 1896Echinoderidae Bitschli, 1876Echinoderes Claparède, 1863E. borealis Greeff, 1869 * Description based on immature stage. Species indeterminatus
BelgiumE. dujardini Claparède, 1863 (TM)
Baltic, Bergen, Netherlands, Britain, Roscoff, Ireland, etc.E. elongatus (Nyholm, 1947)
Gullmarfjord, BritainE. ferrugineus Zelinka, 1928
N EuropeE. gracilis (Zelinka, 1928) * Description based on immature stage.

Gullmarfjord, Mediterranean

E. greeffi (Zelinka, 1928) * Description based on immature stage.

Helgoland

E. krishnaswamyi Higgins, 1985
PlymouthE. kristenseni Higgins, 1985
RoscoffE. lanuginosa Greeff, 1869 * Description based on immature stage.
BelgiumE. levanderi Karling, 1954
BalticE. minimus Zelinka, 1928 * Description based on immature stage.

Gullmarfjord, Mediterranean

E. monocercus Claparède, 1863 * Description based on immature stage.
Bergen, France, MediterraneanE. setiger Greeff, 1869
Bergen, Belgium, BritainE. worthingtoni Southern, 1914
Plymouth, Roscoff, IrelandZelinkaderidae Higgins, 1990Zelinkaderes Higgins, 1990 (TM *Z. floridensis* Higgins, 1990 - Florida)Z. submersus (Gerlach, 1969)
= *Cateria submersa* Gerlach, 1969
North SeaCentroderidae Zelinka, 1896Campylocleres Zelinka, 1913 (TM *C. vanhoeffeni* Zelinka, 1913 - Antarctic)C. adhérensis Nyholm, 1947
Gullmarfjord areaC. macquariae Johnston, 1938
North Sea, W Atlantic, S PacificCondyloderes Higgins, 1969C. multispinosus (McIntyre, 1962)
= *Centroderes multispinosus* McIntyre, 1962 (TM)
ScotlandCentroderes Zelinka, 1907
* Nyholm, 1947 states that this genus is represented on the Swedish West Coast.C. spinosus (Reinhard, 1881)
Br. IslesDracoderidae Higgins, 1990

* No species in the area.

CONCHORHAGAE Zelinka, 1896Semnoderidae Remane, 1936Semnoderes Zelinka, 1907S. armiger Zelinka, 1928 (TM)
Gullmarfjord area, ScotlandCRYPTORHAGAE Higgins, 1968

* No species in the area.

HOMALORHAGIDA Zelinka, 1896HOMALORHAGAE Zelinka, 1896Neocentrophidae Higgins, 1969Paracentrophyes Higgins, 1983 (TM *P. praedictus* Higgins, 1983 - Belize)P. quadridentatus (Zelinka, 1928)
= *Pycnophyes quadridentatus* Zelinka, 1928
= *Pycnophyes flagellatus* Zelinka, 1928
W Sweden, MediterraneanPycnophyidae Zelinka, 1896Pycnophyes Zelinka, 1907P. biserratus (Zelinka, 1928) * This is a larval stage (*Centrophyes*).
W Sweden, MediterraneanP. calmani Southern, 1914
= *P. communis* Nyholm, 1947 & Lang, 1949; non Zelinka
Öresund, Britain, IrelandP. curvatus (Zelinka, 1928) * This is a larval stage (*Centrophyes*).
W Sweden, MediterraneanP. communis Zelinka, 1908
= *Centrophyes longistatus* Zelinka, 1928
Baltic, Gullmarfjord, MediterraneanP. dentatus (Reinhard, 1881)
Kiel, Ireland, Netherlands, Britain,P. denticulatus (Zelinka, 1928) * This is a larval stage (*Centrophyes*).
Britain (Fladden), MediterraneanP. flaveolatus Zelinka, 1928
Gullmarfjord, Öresund, MediterraneanP. kielensis Zelinka, 1928
= *Hyalophyes pelticidus* (Reinhard, 1881)
Kiel, ÖresundP. maximus Reimer, 1963
Baltic (Kadetrinne)P. moderatus (Zelinka, 1928) * This is a larval stage (*Centrophyes*).
W Sweden, MediterraneanP. rectilineatus (Zelinka, 1928) * This is a larval stage (*Centrophyes*).
W Sweden, MediterraneanP. zeliniae Southern, 1914 (TM)
Ireland, Britain (Fladden)Kinorhynchus Sheremetevskij, 1974 (TM *K. paraneapolitanus* Sheremetevskij, 1974 - Black Sea)K. giganteus (Zelinka, 1928)
= *Trachydemus giganteus* Zelinka, 1928

Kiel, Mediterranean

LORICIFERA Kristensen,1983

Kristensen,1983

NANALORICIDA

Nanalicidae Kristensen,1983

Nanalicus Kristensen,1983

N. mysticus Kristensen,1983 (TM)

Engl. Channel, Faeroe Banks, Mediterranean, Öresund? (If the Loricifera reported from Scandinavia are identical with N. mysticus is not known yet. The genus is however known from larvae in Öresund)

Pliciloricidae Higgins & Kristensen,1986

Rugiloricus Higgins & Kristensen,1986 (R. cauliculus Higgins & Kristensen,1986 - South Carolina)

R. sp.

Bohuslän (so far only larvae observed according to Kristensen (pers.comm.))

Plicilorus Higgins & Kristensen,1986 (TM P. enigmaticus Higgins & Kristensen,1986 - North Carolina)

MESOZOA van Beneden,1877

* This taxon should perhaps best be treated as unnatural, separating the two groups involved into two different phyla.

RHOMBOZOOA

@ Parasites in excretory organs of chiefly non-pelagic Cephalopoda

* Dicyemids have also been found in Sagittatus sagittatus & Loligo vulgaris, but not in L. forbesi from Helgoland.

DICYEMIDA

Dicyema von Kölliker,1849

D. paradoxum von Kölliker,1849

= Dicyema clausianum ?,1???

@ Octopus vulgaris, Octopus macropus

Engl. Channel, Mediterranean

D. typus van Beneden,1876

@ Octopus vulgaris

Engl. Channel, Mediterranean

D. moschatum Whitman,1883?

@ e.g. Sepiola rondeleti, ? Eledone moschata

Mediterranean

D. macrocephalum van Beneden,1876

@ e.g. Sepiella oveniana, Sepiola steenstrupiana, Sepia elegans (S. biserialis)

Mediterranean

D. microcephalum Whitman,1883

@ e.g. Sepiola steenstrupiana

Mediterranean

D. schulzianum ?van Beneden,1876?

@ e.g. Sepia elegans

Mediterranean

D. rondeletiolas ?Nouvel,1944?

@ e.g. Sepiella oveniana

Mediterranean

Pseudicyema Nouvel,1933

P. truncatum (Whitman,1883)

@ e.g. Rossia macrosoma & Sepia officinalis

Bohuslän, Engl. Channel, Mediterranean

Dicyemina Whitman,1883?

D. eleodina (Wagener,1857)

= D. muelleri Claparède,18??

@ Eledone cirrhosa, E. moschata

Bohuslän, Norway, Engl. Channel, Mediterranean

D. lameerei Nouvel,1932

@ Eledone cirrhosa & Octopus vulgaris

Engl. Channel, Mediterranean

D. gracile (?Wagener,1857)

= Dicyemina koellikeriana ?,1???

@ e.g. Sepia officinalis

Engl. Channel

HETEROCYEMIDA

Conocymidae Stunkard,1937

= Heterocyemidae

Conocyema van Beneden,1882

C. polymorpha van Beneden,1882?

@ Octopus vulgaris

Mediterranean

Microcyema van Beneden,1882

M. vespa van Beneden,1882?

@ Sepia officinalis

Engl. Channel, Mediterranean

ORTHONECTIDA

PLASMODIGENEA

Rhopaluridae Stunkard,19??

= Orthonectidae Hartmann,1???

Rhopalura Giard,1877

R. philinæ Lang,1954

@ Philine scabra

Bohuslän

R. granosa Atkins,1933

@ Heteranomia squamula

Britain

R. paraphanostomea Westblad,1942

@ Paraphanostoma macroposthium & P. brachyposthium

Gullmarfjord

R. ophiocomae Giard,1877

* Mortensen,1924 is using the name R. giardi Metschnikoff as the name of the parasite in Axiognathus mysticus @ gonads of Axiognathus squamatus, Ophiothrix fragilis & Ophiura albida Bergen, Brittan

R. julini Caullery & Mesnil,1901?

@ in the coelom of Malacoboceros fuliginosus ?

R. pterocirri de Saint-Joseph,1896

@ Pterocirrus macroceros

?

R. metschnikovi Caullery & Mesnil,1901?

@ in the coelom of Spiro martinensis and in mesenchyma and gut wall of Tetrasemma flavidum & Amphiporus ochraceus ?

R. pelseneeri Caullery & Mesnil,1901

@ in mesenchyma and gut wall of Tetrasemma flavidum & T. vermiculus

Vauville (France)

R. pelseneeri var. vermiculica Caullery,1914

@ Tetrasemma vermiculus

Wimereux (France)

R. intoshi Metschnikoff,1881

@ Lineus lacteus

Mediterranean

R. litoralis Shtein,1954

@ Rissoa ocellata & Columbella rosacea

Barents Sea

R. major Shtein,1954

@ Lepeta caeca, Natica clausa & Solariella sp.

Barents Sea

?R. sp. : Punnett,1901

@ Lineus gesserensis

Britain

Intosha Giard,1877

I. leptoplanae Giard,1877

@ In gut of Leptoplana tremellaris

Roscoff

I. linei Giard,1877

@ In skin and gut epithelium of Lineus ruber

Roscoff

Stoecharthrum Caullery & Mesnil,1899

S. giardi Caullery & Mesnil,1899

@ In the coelom of Scoloplos armiger

?

S. sp. : Foster,1982

@ Mytilus edulis

NE Pacific (Washington)

Rhopalura ? sp. Fauvel,1897

@ Ampharete acutifrons

?

Rhopalura ? sp. Arwidsson,1903

@ Nicomache limbricalis

?

Heteronectidae

Pelmatosphaera Caullery & Mesnil,1904

P. polycirri Caullery & Mesnil,1904

@ In the coelom of Polycirrus haematodes

E & W Scotland

ACANTHOCEPHALA Kohlreuther,1771

EOACANTHOCEPHALA Van Cleave,1936

NEOECHINORHYNCHIDA Southwell & Macfie,1925

Neoechinorhynchidae Van Cleave,1919

Neoechinorhynchus Hamann, in Stiles & Hassall,1905

= Neorhynchus Hamann,1892, non Sclater,1869 (Aves) nec Milne-Edwards,1879 (Crustacea)

N. rutili (Müller,1776) (TM)

@ in freshwater fishes and eels.

Baltic

N. agilis (Risso,17??)

@ in Mugil spp. and salmon?

Mediterranean, Scotland?, North Sea?

Tenuisentidae Van Cleave,1936

Paratenuisentis ??,19??

P. ambiguus (Van Cleave,1921)

@ Anguilla anguilla & Gammarus tigrinus (only intermediate host)

introduced from N America to Germany & Netherlands

PALAEACANTHOCEPHALA Meyer,1931

ECHINORHYNCHIDA Southwell & Macfie,1925

Rhadinorhynchidae Travassos,1923

Rhadinorhynchus Lühe,1911

=*Nipporynchus* Chandler,1934

=*Echinosoma* Porta,1907 (p.p.)

R. pristis (Rudolphi,1802) (TM)

=?*R. alosae* (Hermann,1782)

=?*Echinorhynchus subulatus* Zeder,1803

@ in e.g. *Thynnis*, *Scomber*, *Belone*, *Box*, *Brama*, *Gadus* & *Alosa* spp.

Atlantic, Mediterranean

Leptorhynchoides Kostylev,1924

L. plagicephalus (Westrumb, 1821) (TM)

@ in *Acipenser* spp.

Baltic?, Black Sea, Caspian Sea

Pomphorhynchidae Yamaguti,1939

Pomphorhynchus Monticelli,1905, emm. Lühe,1911

P. laevis (Zoega, in Müller,1776) (TM)

=*Echinorhynchus gobii* Viborg,1795

=*Echinorhynchus proteus* Westrumb,1821 (p.p.)

@ in many fishes of both seawater and freshwater origin and exceptionally in dolphins.

Baltic, White Sea, Arctic Seas

P. tereticolle (Rudolphi,1809)

=*Echinorhynchus doublae* Schrank,1790 (p.p.)

=*Echinorhynchus attenuatus* Müller,1779 (p.p.)

=*Echinorhynchus piscinus* Zeder,1800 (p.p.)

=*Echinorhynchus longicollis* Pallas,1781 (p.p.)

@ in e.g. *Bleennius viviparus*, *Cottus scorpius*, *Cyprinus*, *Acerina*, *Platessa*, *Zoarces*, *Gadus* & *Acipencer*

Denmark, Ireland, "Europe"

Echinorhynchidae Cobbold,1879

Echinorhynchinae Travassos,1920

Acanthocephalus Koelreuter,1771, emm. Lühe,1911

A. anguillae (O.F. Müller,1780) (TM)

=*Echinorhynchus globulosus* Rudolphi,1809

@ chiefly in *Tinca tinca*, but also in e.g. eels and flounders.

Baltic

A. lucii (O.F. Müller,1777)

=*Echinorhynchus angustatus* Rudolphi,1809

@ chiefly predatory freshwater fishes, but also e.g. *Hippoglossoides platessoides*, *Platichthys flesus* & *Clupea harengus*

Baltic

Echinorhynchus Zoega, in Müller,1776, emm. Lühe,1911

E. salmonis Müller,1776 (TM of *Metechinorhynchus* Petrotchenko,1956)

@ chiefly freshwater fishes, but also e.g. eel, flatfishes, salmon, cod & *Zoarces*; intermediate host: *Pontoporeia affinis*

Baltic

E. clavula Dujardin,1845

@ freshwater fishes and eel; intermediate host: *Gammarus pulex*

Baltic

E. gadi Zoega, in Müller,1776 (TM)

=*E. acus* Rudolphi,1802

@ in a spectrum of both saltwater and freshwater fishes; intermediate hosts: e.g. *Amphithoe rubricata*, *Calliopius laeviusculus*, *Gammarus locusta*, *Pontoporeia femorata*

Baltic - Bohuslän, North Sea, Britain, Barents Sea, White Sea, etc.

E. truttae Schrank,1788

=*Metechinorhynchus truttae* : Auctt.

@ *Salmo fario*, *Thymallus thymallus*, *Esox lucius*

North Sea

E. serpentulus Grimm,1870

@ *Cottus quadrimaculatus*

White Sea

E. cestodicola von Linstow,1905

@ on "Bothrioccephalus nigropunctatus" in *Sebastodes norvegicus*

Murmansk

POLYMPHIDIA

Polympiphidae Meyer,1931

Arhythmorhynchus Lühe,1911

A. longicole (Villot,1875)

@ *Larus ridibundus* L. canus, *Mergus merganser*, *Arenaria interpres*, *Erolia maritima*

Britain, Europe, Barents Sea, Arctic

A. invaginabile (von Linstow,1890)

@ in *Numenius arquatus*

Europe

A. invaginabile sueicum Lundström,1942

@ *Anthus spinolella littoralis*

Sweden

A. frassoni (Molin,1858) (TM)

@ in *Numenius* spp., *Merganser serrator* & *Larus marinus*

Europe

A. roseum (Molin,1858)

@ *Cantharus vulgaris*; 1st interm. host: *Palaemon squilla*; 2nd interm. host: *Platessa passer*, *Leuciscus* sp.

Europe

Corynosoma Lühe,1905

C. strumosum (Rudolphi,1802)

=*Echinorhynchus strumosus* Rudolphi,1802 (p.p.) (TM)

=*Echinorhynchus ventricosus* Rudolphi,1809

=*Echinorhynchus hystris* Bremser,1824 (p.p.)

=*Echinorhynchus striatus* Villot,1875

=*Echinorhynchus gibber* Olsson,1894 (p.p.)

@ adults in intestine of mammals, e.g. *Phoca Phocaena*, *Halichoerus*, *Mustela* or birds, e.g. *Phalacrocorax*, *Larus*, *Pelecanus* ; intermediate hosts: marine amphipoda; larvae encysted in fishes (transport hosts), e.g. *Cottus*, *Clupea*, *Pleuronectes*, *Anguilla*

Bothnian Bay (in *Phoca hispida*), Kattegatt - Skagerrak area (in *Phoca vitulina*), Europe

C. semerne (Forssell,1904) Lühe,1911

=*Echinorhynchus semerne* Forssell,1904

=*Echinorhynchus strumosum* Rudolphi,1802 (p.p.)

=*Echinorhynchus hystris* Bremser,1824 (p.p.)

@ adults in seals, e.g. *Phoca Phocaena*, *Cystophora*, *Odobenus*, *Otaria*, whales (*Phocaena phocaena*) or birds : e.g. *Phalacrocorax* & *Mergus* ; intermediate hosts: amphipoda ; transport hosts: fishes, e.g. *Myoxocephalus*, *Clupea*, *Pleuronectes*, *Rombus*, *Gasterosteus*

Europe, e.g. Bothnian Bay (in *Phoca hispida*), Greenland, Arctic region

C. reductum (von Linstow,1905)

=*Echinorhynchus reductus* von Linstow,1905

@ *Phoca hispida*

Arctic region

C. mergi Lundström,1941

@ *Mergus serrator*, *Cephus grylle*, *Phalacrocorax carbo*

Europe

C. tunitae (Weiss,1941)

@ *Sula bassana*, *Phalacrocorax* spp.

Shetlands, Britain, Tunisia

Bolbosoma Porta,1908

=*Bolborhynchus* Porta,1906, non Bonaparte,1857 (Aves)

B. balaenae (Gmelin,1790)

=*Sipunculus lendix* Phipps,1774 (nomen oblitum?)

=*Echinorhynchus balaenae* Gmelin,1790

=*Echinorhynchus porrigens* Rudolphi,1819

@ intestine of *Balaenoptera*, *Megaptera*, *Balaena* & *Hyperoodon*

Iceland, Australia

B. capitatum (von Linstow,1880)

=*Echinorhynchus capitatus* von Linstow,1880

@ intestine of *Pseudorca crassidens*, *Globicephalus melas*, *Physeter macrocephalus*, *Steno rostratus* & *Megaptera boops* ?

Iceland, Faeroes, Mediterranean

B. turbinella (Diesing,1851) (TM)

=*Echinorhynchus turbinella* Diesing,1851

=*Echinorhynchus balanoccephalus* Owen,1803

=*Echinorhynchus ruber* Collett,1886

@ intestine of e.g. *Balaenoptera* spp. & *Hyperoodon rostratus*

Subarctic - N & S Atlantic, Pacific

B. brevicolle (Malm,1867)

=*Echinorhynchus brevicolle* Malm,1867

@ in *Balaenoptera* spp. & *Physeter macrocephalus*

North Atlantic, e.g. Sweden (in the "Malmiska hvalen"), S Africa

B. vasculosum (Rudolphi,1819)

=*Echinorhynchus vasculosus* Rudolphi,1819

=*Echinorhynchus aurantiacus* Risso,1826

=*Echinorhynchus pellucidus* Leuckart,1828

=*Echinorhynchus annulatus* Molin,1856

=*Echinorhynchus serranti* Linton,1888

=*Echinorhynchus bifasciatus* Lühe,1904

@ intestine of *Delphinus delphis* & *Mesoplodon bidens* (larvae in teleosts)

Atlantic, Mediterranean

B. heteracanthis (Heitz,1918)

@ Body cavity & pyloric appendages of *Salmo salar*

Europe

Polymorphus Lühe,1911

P. (Polymorphus) boschadis (Schrank,1788) (TM)

@ adults in birds, e.g. *Larus*, *Uria*, *Cygnus*, *Anas*, *Cygnus*, *Uria*, *Harelda* ; larval development in e.g. *Gammarus*, *Gobius*, *Cyclopterus*, *Anguilla* may be intermediate hosts

Europe

P. minutus (Goeze,1782)

@ Egypt, in different "waterbirds", e.g. *Somateria*, *Anas*, *Cygnus*, *Uria*, *Harelda* ; larval development in e.g. *Gammarus*, *Potamobius*

Murmansk

P. phippsi Kostylev,1922

@ in *Somateria mollissima*

Murmansk

P. magnus Skrjabin,1913

@ in *Fulgula rufina* and *Bucephala clangula*

Russia

P. (Profiliocillus) botulus (Van Cleave,1916)

@ *Somateria mollissima* ; Larvae in *Carcinus maenas*, *Hyas araneus* & *Pagurus pubescens*

Europe, e.g. Waddenze, Britain, Arctic Seas

Filicollis Lühe,1911

F. anatis (Schrank,1788) (TM)

@ e.g. *Larus*, *Somateria*, *Tadorna*, *Cygnus*, *Mergus*, *Anser* ; development in *Asellus aquaticus*

Europe

Plagiorhynchidae Meyer,1931

Porrorhinae

Oligoterorhynchus Monticelli,1914

O. campylurus (Nitzsch,1857) (TM)

@ e.g. *Stercorarius scura*

N Europe

Plagiorhynchinae Meyer,1931

Plagiorhynchus Lühe,1911

P. crassicollis (Villot,1875) (TM)
= *P. lanceolatus* (von Linstow,1876)
@ in different birds, e.g. oystercatcher
Europe, Greenland

P. lineare (Westrumb,1821)
@ *Sterna sandvicensis*, *Rissa tridactyla*
Europe

P. odhneri Lundström,1942
@ in different birds, e.g. oystercatcher
Sweden

NEMATOMORPHA Vejdovsky,1886

NECTONEMATOIDEA Rauther,1930

NECTONEMATIDA

Nectonematidae

Nectonema Verrill,1879

N. munidae Brinkmann,1930
@ in *Munida*, *Pagurus pubescens* & *Pontophilus norvegicus*
Bergen

N. svensksundi Bock,1908
@ ?
Spitzbergen

N. agile Verrill,1879 (TM)
@ in pagurides, *Palaeon serratulus*, *Liocarcinus pusillus* & *Palaemonetes vulgaris*
Roscoff, Massachusetts

GASTROTRICHA Metchnikoff,1864

MACRODASYIDA Remane,1924

Dactylopodidae Strand,1929

Dactylopodida Strand,1929
= *Dactylopodalia* Remane, in Kükenthal & Krumbach,1929
= *Dactylopodella* Remane,1926, non G.O. Sars,1905 (Crustacea)

D. baltica (Remane,1926) (TM)
Sweden

D. typhele (Remane,1927)
= *D. wellsi* d'Hondt,1965
Helgoland, Arcachon

D. cornuta cornuta (Swedmark,1956)
= *Dactylopodalia cornuta* Swedmark,1956
Helgoland, Engl. Channel, Roscoff

D. cornuta brevis d'Hondt,1966
Arcachon

D. roscovita (Swedmark,1967)
Roscoff

Dendrodasys Wilke,1954

D. gracilis Wilke,1954 (TM)
Scotland

Xenodasys Swedmark,1967
= *Chordodasys*

X. sanctigoulini Swedmark,1967
Roscoff

X. riedli Schöpfer-Sterrer,1969
Scotland

Lepidodasyidae Remane,1927

Lepidodasys Remane,1926

L. martini Remane,1926 (TM)
Sweden, Baltic, Bergen, North Sea

L. platyrurus Remane,1927
Baltic, Irish Sea, Mediterranean

Cephalodasys Remane,1926

C. maximus Remane,1926 (TM)
Kiel, Sylt, Bergen

C. turbanelloides (Boaden,1960)
Sweden, Arcachon, Irish Sea

C. cambriensis (Boaden,1963)
Wales, Roscoff

C. littoralis Renaud-Debyser,1964
Scotland, Arcachon

Dolichodasys Gagne,1977

D. elongatus Gagne,1977 (TM)

D. carolinensis Ruppert & Shaw,1977
Scotland

Megadasys Schmidt,1974

= *Thiodasys* Boaden,1974

M. sternerri (Boaden,1974)

= *M. minor* Kisielewski,1987
Bergen, Scotland

Mesodasys Remane,1951

M. laticaudatus Remane,1951
= *M. lobocerus* (Boaden,1960)
Sweden, Kiel, Bergen, Sylt

M. littoralis Remane,1951
Arcachon, Napoli

Paradasys Remane,1934

P. subterraneus Remane,1934 (TM)
Helgoland, Baltic, Tromsö, Engl. Channel

P. hexadactylis Karling,1954
Sweden, Arcachon, Irish Sea

Pleurodasys Remane,1927

P. helgolandicus Remane,1927 (TM)
= *P. megasoma* (Boaden,1963)
Poland, Helgoland, Wales, Arcachon, Engl. Channel

Macrodasyidae Remane,1927

Macrodasys Remane,1924

M. buddenbrocki Remane,1924 (TM)
Sweden, Baltic, Bergen, ?Tromsö, North Sea, Irish Sea

M. cephalatus Remane,1927
Bergen, Helgoland

M. caudatus Remane,1927
Helgoland, Arcachon, Roscoff, Irish Sea

M. affinis Remane,1936
Sylt, Engl. Channel

M. balticus Roszczak,1939
Poland

M. remanei Boaden,1963
Wales

M. sp. Remane,1963
Helgoland

M. sp. Boaden,1963
Wales

Urodasys Remane,1926

U. mirabilis Remane,1926 (TM)
= *U. roscovensis* (Kisielewski,1987)
Öresund, Baltic, North Sea

U. sp. :?19??
Arcachon

U. sp. :?19??
Bergen

Chordodasys Schoepfer-Sterrer,1969 (TM) *Chordodasys riedli* Schoepfer-Sterrer,1969 -?
* Correct family? Published in Cah. de Biol. Marin 10:391-404.

Planodasyidae Rao & Clausen,1970

Crasiella Clausen,1968

C. diplura Clausen,1968
Bergen, Scotland

C. oceanica d'Hondt,1974
Roscoff

Planodasys ?19??
*No species within the area?

Taumastodermatidae Remane,1926

Taumastodermatinae Remane,1926

Platydasys Remane,1927 non Butler,1892 (Lepidoptera)
* The name should be changed

P. maximus maximus Remane,1927 (TM)
Poland, Bergen, Irish Sea

P. maximus celticus Swedmark,1955
Roscoff

P. rarus Forneris,1961
Helgoland

P. mastigurus Clausen,1965
Bergen

P. ocellatus Clausen,1965
Bergen

P. brachycephalus Lévi,1954
Roscoff, N Ireland

P. ruber Swedmark,1956
W France

P. schultzi Remane,1940 (nomen nudum)
* Not mentioned in Renaud-Mornant,1986 (Stygo. Mundi)
Baltic

Pseudostomella Swedmark,1956

- P. roscovitae** Swedmark,1956 (TM - Roscoff)
North Sea, Engl. Channel, Roscoff
- Ptychostomella** Remane,1926
- P. ommatophora** Remane,1927
Sweden, Denmark, Germany
- P. pectinata** Remane,1926 (TM)
Denmark, Kiel
- P. helena** Roszczak,1939
Poland
- P. bergenensis** Clausen,1996
Bergen
- P. sp.** Forneris,1961
Helgoland
- Tetranchyroderma** Remane,1926
- T. hystrix** Remane,1926 (TM)
Kiel, Wales, France
- T. apum** Remane,1927
Helgoland, Engl. Channel
- T. polycanthum** (Remane,1927)
= *Echinodasys polycanthus* Remane,1927
Bergen, Helgoland, Engl. Channel, Irish Sea
- T. megastomum** (Remane,1927)
= *Echinodasys megastoma* Remane,1927
= *T. suecicum* Boaden,1960
Kiel, Sweden, Sylt, Engl. Channel, Roscoff
- T. cirrophora** Lévi,1950
Baltic, Sweden, Engl. Channel, Roscoff, Irish Sea
- T. coeliopodium** Boaden,1963
Ireland
- T. tribulosum** Clausen,1965
Bergen, Helgoland
- T. dragescoei** Swedmark,1967
Roscoff
- T. antennatum** Luporini & al.,1970
French Atlantic coast, Mediterranean
- T. massiliense** Swedmark,1956
European atlantic, Mediterranean
- T. pugetensis** Wieser,1957
Bergen
- T. thysanogaster** Boaden,1965
Bergen
- T. norvegicum** Clausen,1996
Bergen, Trondheim
- Thaumastoderma** Remane,1926
- T. heideri** Remane,1926 (TM)
Sweden, Baltic, Bergen, North Sea
- T. bifurcatum** Clausen,1991
Bergen, Trondheim, Tromsø
- T. truncatum** Clausen,1991
Bergen
- T. swedmarki** Lévi,1959
Roscoff
- T. cantacuzeni** Lévi,1958
Bretagne
- T. arcassonense** d'Hondt,1965
Arcachon
- T. ramuliferum** Clausen,1965
Engl. Channel, Mediterranean
- T. renaudae** Kisielewski,1987
France (north or west coast)
- Diplodasyinae
- Acanthodasys** Remane,1927
- A. aculeatus** Remane,1927 (TM)
Sweden, Baltic, North Sea, Irish Sea, Roscoff
- A. arcassonensis** Kisielewski,1987
Arcachon
- A. sp.** Forneris,1961
Sylt
- Diplodasys** Remane,1927 (TM) *D. platydasyooides* Remane,1927 - Mediterranean
- D. minor** Remane,1927
Helgoland
- D. ankeli ankeli** Wilke,1954
Sweden, North Sea, Bergen
- D. caudatus** Kisielewski,1987
W France
- D. swedmarki** Kisielewski,1987
W France
- Hemidasys** Claparède,1867
- Turbanellidae Remane,1925
- Turbanella** M. Schultze,1853
- T. hyalina** M. Schultze,1853 (TM)
= *T. varians* Maguire,1976 (p.p.)
Sweden, Baltic, North Sea
- T. (Zelinika) plana** (A. Giard,1904)
France
- T. cornuta** Remane,1925
= *T. varians* Maguire,1976 (p.p.)
Sweden, Bergen, "cosmopolite"
- T. subterranea** Remane,1934
Sylt, Baltic
- T. ambronensis** Remane,1943
= *T. digitifera* d'Hondt,1965
Sylt, European atlantic
- T. lutheri** Remane,1952
Stockholm, Bergen, North Sea
- T. remanei** Forneris,1961
Kiel, North Sea
- T. otti** Schrom,1???
= *T. mustela* Wieser,1957
Arcachon
- T. bocqueti** Kaplan,1958
Engl. Channel
- T. multidigitata** Kisielewski,1987
W France
- Desmodasys** Clausen,1965
- D. phocooides** Clausen,1965 (TM)
Bergen
- Dinodasys** Remane,1927
- D. mirabilis** Remane,1927 (TM)
Sweden, Baltic, North Sea, Roscoff
- Paraturbanella** Remane,1927
- P. dohrni** Remane,1927 (TM)
Kiel, Sylt
- P. scanica** Clausen,1996
Bergen, Trondheim, Tromsø
- P. teissieri** Swedmark,1954
Sylt, Engl. Channel, Wales
- P. armoricana** (Swedmark,1954)
Roscoff
- P. intermedia** Wieser,1957
N Ireland
- P. pallida pallida** Luporini & al.,1971
England
- P. cuanensis** Maguire,1976
Ireland, England
- P. eireanna** Maguire,1976
N Ireland
- P. sp. : Schmidt,1972**
Tromsø
- Pseudoturbanella** d'Hondt,1968
- P. stylifera** d'Hondt,1968 (TM)
France (Landes)
- CHAETONOTIDA** Remane,1924
- MULTITUBULATINA** d'Hondt,1971
- Neodasyidae Remane,1929
- Neodasys** Remane,1927
- N. chaetonotoideus** Remane,1927 (TM)
Sylt, Baltic, Arcachon, Irish Sea
- N. uchidai** Remane,1961
Sylt, Mediterranean
- PAUCITUBULATINA** d'Hondt,1971
- Xenotrichulidae Remane,1927
- Draculiciterinae
- Draculiciteria** Hummon,1974
- D. tessellata** (Renaud-Mornant,1968)
=? *Xenotrichuloides mirabilis* d'Hondt,1967
Mediterranean, Arcachon?
- Xenotrichulinae** Remane,1927
- Heteroxenotrichula** Wilke,1954
- H. pygmaea** (Remane,1934)
= *Xenotrichula flandrensis* d'Hondt,1968

- Öresund, Schilksee, Arcachon
- H. affinis** (Remane, 1934)
Öresund, Arcachon, North Sea
- H. subterranea** (Remane, 1934)
Öresund, Arcachon, North Sea
- H. arcassonensis** Ruppert, 1979
Brittany, Arcachon
- H. wilkeae** Ruppert, 1979
Arcachon, German North Sea coast
- H. squamosa** Wilke, 1954 (TM)
North Sea, Mediterranean
- Xenotrichula** Remane, 1927
- X. velox** Remane, 1927
Baltic, Öresund, Irish Sea
- X. intermedia** Remane, 1934
= *X. beauforti* Lévi, 1950
=? *X. punctata* Wilke, 1954
* Renaud-Mornant, 1986 is keeping *X. punctata* as a separate sp., known from European Atlantic & the Mediterranean.
Öresund, Tromsö, Schilksee, Amrum, Roscoff, "cosmopolite"
- X. cornuta** Wilke, 1954
S North Sea, Flanders Coast.
- X. variocirrata** d'Hondt, 1966
Arcachon
- X. gymnocephala** Remane, 1940 (nomen nudum)
Baltic
- X. micracantha** (Remane, 1926)
= *Chaetonothus micracanthus* Remane, 1926 (TM)
Helgoland
- X. bispina** Roszczak, 1979
Poland, Black Sea
- X. simplex** Mock, 1979
Sylt
- X. carolinensis** Ruppert, 1979 *syltensis* Mock, 1979
Sylt
- X. quadrifurcata** Kisielewski, 1988
W or N France
- X. punctata** Wilke, 1954
* Possibly a synonym of *X. intermedia* Remane, 1934. See above
Bergen?
- Chaetodontidae** Zelinka, 1889
- Ichthydium** Ehrenberg, 1830 (TM) *Cercaria podura* O.F. Müller, 1786 - fresh water
- I. hummoni** Ruppert, 1977
Arcachon, Sylt
- I. rostrum** Roszczak, 1968
Poland
- I. rupperti** Mock, 1979
Sylt
- I. supralittoralis** Mock, 1979
Sylt
- Musellifer** Hummon, 1969
- M. profundus** Vivier, 1974
Scotland
- Halichaetonotus** Remane, 1936
- H. aculifer aculifer** (Gerlach, 1953)
Kiel, Engl. Channel, Mediterranean
- H. arenarius** (d'Hondt, 1971)
Roscoff
- H. atlanticus** Kisielewski, 1988
W France
- H. balticus** Kisielewski, 1974
Baltic
- H. batillifer** (Luporini & al., 1972)
France
- H. dubius** (Remane, 1926)
=? *H. decipiens* (Remane, 1929)
Kiel, Helgoland
- H. jucundus** (d'Hondt, 1971) emend. Mock, 1979
Sylt, Ille Calot
- H. lamellatus** Kisielewski, 1974
Baltic
- H. littoralis** (d'Hondt, 1971)
Roscoff
- H. paradoxus** (Remane, 1927)
Arcachon, Mediterranean
- H. pleuracanthus** (Remane, 1926)
Baltic, Engl. Channel, Ireland
- H. schromi** Kisielewski, 1974
=? *H. jucundus* Schrom, 1972 non (d'Hondt, 1971)
Baltic, Sylt
- H. spinosus** Mock, 1979
Baltic, Sylt, France
- H. tentaculatus** (d'Hondt, 1971)
Arcachon
- Chaetonotus** Ehrenberg, 1830
- C. marinus** Giard, 1904
* Not mentioned in Renaud-Mornant, 1986 (Stygof. Mundi)
Sweden, France
- C. balticus** Remane, 1926
* Not mentioned in Renaud-Mornant, 1986 (Stygof. Mundi)
Kiel
- C. dispar** Wilke, 1954
Sweden, Roscoff, Mediterranean
- C. atrox** Wilke, 1954
Sylt, Bretagne, Wales
- C. condensus** Mock, 1979
Sylt
- C. lacunosus** Mock, 1979
Sylt
- C. somniculosus** Mock, 1979
Denmark, Sylt
- C. tempestivus** Mock, 1979
Sylt
- C. variosquamatus** Mock, 1979
Sylt
- C. parthenopeius** Wilke, 1954
Engl. Channel, European Atlantic, Mediterranean
- C. oceanides** d'Hondt, 1971
* Not mentioned in Renaud-Mornant, 1986 (Stygof. Mundi)
Ille Calot
- C. inaequidentatus** Kisielewski, 1988
Engl. Channel
- C. larus** (O.F. Müller, 1773)
England
- C. linguaformis** Voigt, 1904
Netherlands
- C. maximus** Ehrenberg, 1831
?
- C. neptuni** Wilke, 1954
Belgium, France
- C. persetosus** Zelinka, 1889
* Fresh-water species found in the Baltic
Poland
- C. hystrix** Metsch.,
* Not mentioned in Renaud-Mornant, 1986 (Stygof. Mundi) * Fresh-water species found in the Baltic
Poland
- Aspidiophorus** Voigt, 1904
- A. bisquamatus** Mock, 1979
Sylt
- A. marinus** Remane, 1926
Sweden, "cosmopolite"
- A. mediterraneus** Remane, 1927
Engl. Channel, Mediterranean
- A. ornatus** Mock, 1979
Sylt
- A. squamulosus** (Roszczak, 1935)
Poland
- A. paramediterraneus** Hummon, 1974
?
- Heterolepidoderma** Remane, 1926
- H. axi** Mock, 1979
Sylt
- H. grandiculum** Mock, 1979
Sylt, Denmark, France
- H. marinum** Remane, 1926
Kiel, North Sea, Roscoff
- H. (?) dubium** Remane, 1927
* Not mentioned in Renaud-Mornant, 1986 (Stygof. Mundi)
?
- H. foliatum** Renaud-Mornant, 1967
* Fresh-water species found in the Baltic
Arcachon
- H. ocellatum** Metchnikoff,
* Not mentioned in Renaud-Mornant, 1986 (Stygof. Mundi) * Fresh-water species found in the Baltic
Baltic
- H. arenosum** Kisielewski, 1988
Belgium, Engl. Channel
- H. armatum** Schrom, 1966
Netherlands

H. clipeatum Schrom,1972
Belgium

Lepidodermella Blake,1933 (TM) *Chaetonotus squammatus* Dujardin,1841 - fresh water)

L. limogena Schrom,1972
Scotland

L. punctatum (Greuter,19??)
* Not mentioned in Renaud-Mornant,1986 (Stygof. Mundi) *Fresh-water species found in the Baltic?
L. squamatum psammicum Roszczak,1968
=? *L. squamatum* (Dujardin,1841)
*Fresh-water species found in the Baltic
Poland

ROTIFERA Cuvier,1798
= **ROTATORIA** Ehrenberg,1838

SEISONIDEA Bartos,1959
= **PARAROTATORIA** Sudzuki,1964
SEISONIDA Remane,1933

Seisonidae

Seison Grube,1859

S. nebaliae Grube,1859 (TM)
Roscoff, Brest, on *Nebalia*

S. annulatus Claus,1876
Roscoff, W North Sea, on *Nebalia*

BDELLOIDEA Hudson & Gosse,1886
= **DIGONONTA** Wesenberg-Lund,1899

BDELLOIDA

Phiodinidae Bryce,1910

Rotaria Scopoli,1777
= Rotifer Cuvier,1798

R. macroura (Schrank,1803)
= *Rotifer macrourus* Schrank,1803
= *Esechelina gracilicauda* Bory de St. Vincent,1826
Europe (in brackish water)

R. citrina (Ehrenberg,1838)
= *Rotifer citrinus* Ehrenberg,1838
= *Rotifer vulgaris* Schrank,
Europe (in the Sea and in brackish water)

R. rotatoria rotatoria (Pallas,1766)
= *Brachionus rotatorius* Pallas, 1766
= *Rotifer vulgaris* Schrank, 1801
Cosmopolitan, in limnic - marine waters

R. sp. (Remane,1949)
Helgoland, German North Sea part

Zelinkiella Harring,1913
= *Discopus* Zelinka,1888 non Thomson,1864 (Coleoptera)

Z. synaptae Lankester,1868
@ On *Lepiosynapta* spp, including *L. inhaerens*
Engl. Channel, North Sea?

Philodina Ehrenberg,1830 emend. Bryce,1910

P. citrina Ehrenberg,1832
E Baltic, Bergen

P. roseola Ehrenberg,1832
Tvärminne, (in pools containing weak brackish water)

MONOGONONTA Plate,1889
= **EUROTATORIA** Bartos,1959

PLOIMA Remane,1933
= **PLOIMIDA** Hudson & Gosse,1886

Colurellidae Bartos,1959

Lepadella Bory de St. Vincent,1826

L. psammophila Tschaschel,1979
Sylt, German North Sea beaches

L. oblonga (Ehrenberg,1834)
Tvärminne, in fresh-water rock-pools

L. patella (O.F. Müller,1786)
= *Brachionus patella* O.F. Müller,1786 (TM)
Tvärminne

L. similis (Lucks,1912)
= *L. persimilis* De Ridder,1961
Tvärminne

L. quadrarinata (Stenroos,1898)
Tvärminne

L. triptera (Ehrenberg,1830)
Tvärminne

L. acuminata (Ehrenberg,1834)
Tvärminne, in fresh-water rock-pools

Colurella Bory de St. Vincent,1824

C. unicaudata Godske Eriksen,1968

Bergen, Tvärminne, Belgium

C. adriatica Ehrenberg,1831
W & N Norway, S Finland, Roscoff, Zuiderzee

C. colurus (Ehrenberg,1830)
Tvärminne, W & N Norway

C. dicentra (Gosse,1887)
Bergen

C. halophila Wulfert,1942
W Norway

C. uncinata (O.F. Müller,1773)
= *Brachionus uncinatus* O.F. Müller,1773 (TM)
Tvärminne, Bergen

C. geophila Donner,1951
Tvärminne?

Lophocaris Ehrenberg,1838

L. salpina (Ehrenberg,1834) (TM)
Roscoff

Brachionidae Wesenberg-Lund,1899

Brachionus Pallas,1766 (TM) *Tubipora urceus* Linnaeus,1758 - probably synonom. to *B. urceolaris* Müller

B. plicatilis O.F. Müller,1786
Tvärminne, Bergen, Roscoff

B. quadridentatus Hermann,1783
Tvärminne

B. rubens Ehrenberg,1838

Tvärminne, on *Daphnia magna* & *D. longispina*

B. urceolaris (O.F. Müller,1773)

Tvärminne, Bergen

Notholca Gosse,1886

N. acuminata (Ehrenberg,1832)
Tvärminne, Zuiderzee, W Norway, in fresh-water rock-pools

N. marina Focke,1961
=? *N. thalassia* Gosse, in Hudson & Gosse,1886
Malmö, Tvärminne, W Norway

N. striata (O.F. Müller,1786)
= *Brachionus striatus* O.F. Müller,1786
W & N? Norway, Tvärminne, N Germany, Zuiderzee

N. bipalium (Müller,1786)
= *Brachionus bipalium* O.F. Müller,1786
W Norway, Tvärminne

N. liepetterseni Godske Björklund,1972
W Norway

N. psammarina Buchholz & Rühmann,1956
German Baltic & North Sea coasts, Öresund, W Norway

N. caudata Carlin,1943
Tvärminne

N. labis Gosse, in Hudson & Gosse,1887
Tvärminne

N. salina Focke,1961
W Norway, German Baltic & North Sea coasts

N. squamula (O.F. Müller,1786)
= *Brachionus squamula* O.F. Müller,1786 (TM)
Belgium, France

Pseudonotholca Marukawa,1928

P. japonica Marukawa,1928 (TM)
N & W Norway

Argonotholca Gillard,1948

A. foliacea (Ehrenberg,1838) (TM)
Tvärminne

Keratella Bory de St. Vincent,1822

K. cruciformis (Thompson,1892)
W Norway, planktonic, Roscoff

K. eichwaldi (Levander,1894)
Tvärminne, Zuiderzee

K. cochlearis (Gosse,1851)
Tvärminne, Zuiderzee

K. quadrata (O.F. Müller,1786)
= *Brachionus quadridentatus* O.F. Müller,1786 (TM)
Tvärminne, Zuiderzee

Euchlanis Ehrenberg,1832

E. dilatata Ehrenberg,1832 (TM)
Tvärminne

Tripleuchlanis Myers,1930

T. plicata (Levander,1894)
= *Euchlanis plicata* Levander,1894 (TM)
Tvärminne

Kellicottia Ahlstrom,1938

- K. longispina** (Kellcott,1879)
 = *Anuraea longispina* Kellcott,1879 (TM)
 Europe, freshwater-species, sometimes found in brackish waters
- Platyiota** Herring,1913
- P. quadricornis** (Ehrenberg,1832) (TM)
 Belgium, Spain
- Mytilina** Bory de St. Vincent,1836
- M. mucronata** (O.F. Müller,1773)
 = *Brachionus mucronatus* O.F. Müller,1773 (TM)
 Tvärrminne
- M. ventralis** (Ehrenberg,1832)
 Tvärrminne
- Trichotria** Bory de St. Vincent,1827
- T. pocillum** (O.F. Müller,1776)
 = *Trichoda pocillum* O.F. Müller,1776 (TM)
 Tvärrminne
- Lecanidae** Bartos,1959
- Lecane** Nitzsch,1827
 = *Monostyla* Ehrenberg,1830
- L. ohioensis** (Herrick,1885)
 Tvärrminne
- L. closterocerca** (Schmarda,1859)
 Tvärrminne
- L. luna** (O. F. Müller,1776)
 = *Cercaria luna* O.F. Müller,1776 (TM)
 Tvärrminne
- L. nana** (Murray,1913)
 Tvärrminne
- L. inermis** (Bryce,1892)
 Tvärrminne, in fresh-water rock-pools
- L. lunaris** (Ehrenberg,1832)
 Tvärrminne, in fresh-water rock-pools
- Proalidae** Bartos,1959
- Proales** Gosse,1886
- P. germanica** Tschaschel,1979
 Sylt, German North Sea coast
- P. halophila** (Remane,1929)
 Tvärrminne, German Baltic & North Sea coasts, W Norway
- P. oculata** Tschaschel,1979
 Sylt, German North Sea coast
- P. syltensis** Tschaschel,1979
 Sylt, German North Sea coast
- P. paguri** Thane-Fenchel,1966
 Denmark, Roscoff
- P. gonothyraea** Remane,1929
 Roscoff, in the theca of *Gonothyraea loveni* Allman
- P. reinhardtii** (Ehrenberg,1834)
 Roscoff, W & N Norway, Plymouth, Öresund, S Finland
- P. similis** De Beauchamp,1908
 Tvärrminne
- P. decipiens** (Ehrenberg,1832)
 = *Notommata decipiens* Ehrenberg,1832 (TM)
 Tvärrminne, in fresh-water rock-pools
- Notommatidae** Remane,1933 (p.p.)
- Notommata** Ehrenberg,1830 (TM) *Vorticella aurita* O.F. Müller,1786 - fresh & brackish water species
- N. glyphura** Wulfert,1935
 Tvärrminne, in fresh-water rock-pools
- N. sp.?** d'Hondt,1970
 Roscoff
- Cephalodella** Bory de St. Vincent,1826
- C. auriculata** (O.F. Müller,1773)
 = *Vorticella auriculata* O.F. Müller,1773
 Tvärrminne
- C. delicata** Wulfert,1937
 Tvärrminne
- C. gracilis** (Ehrenberg,1832)
 Tvärrminne
- C. megalcephala** (Glascott,1893)
 Tvärrminne
- C. pachydactyla** Wulfert,1937
 Tvärrminne, Bergen
- C. wrighti** Wulfert,1960
 Tvärrminne
- C. gibba** (Ehrenberg,1838)
 Roscoff
- C. ventripes** (Dixon-Nuttall,1901)
 Tvärrminne
- C. gibba microdactyla** Koch-Althaus,1963
 Tvärrminne, Bergen
- C. catellina** (O.F. Müller,1786)
 = *Cercaria catellina* O.F. Müller,1786 (TM)
 Roscoff
- C. sp.** d'Hondt,1970
 Roscoff
- Pleurotrocha** Ehrenberg,1830 (TM) *P. petromycon* Ehrenberg,1830 - fresh & brackish water species
- P. atlantica** Myers,1936
 American Atlantic coast, Mediterranean
- Lindidae** Remane,1933
- Halolindia** Remane,1933
- H. gravita** (Lie-Pettersen,1905)
 = *Lindia tecusa* Herring & Myers,1922 (TM)
 Bergen, Roscoff
- Lindia** Dujardin,1841 (TM) *L. torulosa* Dujardin,1841 - fresh water
- L. sp. nov. ? (aff. pallida)** d'Hondt,1970
 Roscoff
- Trichocercidae** Remane,1933
- Trichocerca** de Lamarck,1801
- T. pediculus** Remane,1949
 Amrum, Kieler Bucht, German Baltic & North Sea coasts
- T. taurcephala** (Hauer,1931)
 Tvärrminne, German Baltic & North Sea coasts
- T. marina** (Daday,1890)
 Baltic, North Sea, Bergen, Roscoff, Plymouth
- T. rattus** (O.F. Müller,1776)
 = *Trichoda rattus* O.F. Müller,1776 (TM)
 Tvärrminne
- T. arctica** (Voigt,1957)
 = *Diurella minuta* Olofsson,1918, non *Coelopus minutus* Gosse
 Tvärrminne, Spitsbergen
- T. brachyura** (Gosse,1851)
 Tvärrminne
- T. longiseta** (Schrank,1802)
 Tvärrminne, in fresh-water rock-pools
- Gastropodidae** Remane,1933
- Ascomorpha** Perty,1850 (TM) *A. ecaudis* Perty,1850 - fresh water
- A. saltans** Bartsch,1870
 Europe
- Chromogaster** Lauterborn,1893
- C. ovalis** (Bergendal,1892)
 = *C. testudo* Lauterborn,1893 (TM)
 Europe
- Dicranophoridae** Remane,1933
- Aspelta** Herring & Myers,1928 (TM) *Diglena circinator* Gosse,1886 - fresh water
- A. harringtoni** Remane,1929
 German & Danish Baltic beaches, Scandinavian Atlantic coast, Roscoff
- A. reibischii** (Remane,1929)
 German & Danish Baltic & North Sea beaches, W Norway
- A. clydona** Herring & Myers,1928
 E Sweden, German Baltic, Bergen, South Finland
- A. clydona europaea** Hauer,1939
 E Baltic
- Encentrum** Ehrenberg,1838
- E. algente** Herring,1921
 Bergen, in rock-pool, Kieler Bucht
- E. bidentatum** (Lie-Pettersen,1906)
 Bergen, in rock-pool, Kieler Bucht
- E. cruentum** Herring & Myers,1928
 W Norway
- E. enteromorphae** Otto,1936
 Kieler Bucht
- E. flexilis** Godske-Eriksen,1968
 Bergen, in rock-pool
- E. ambiguum** Tschaschel,1979
 Sylt, German North Sea coast
- E. axi** Tschaschel,1979
 Sylt, German North Sea coast
- E. eulitorale** Tschaschel,1979
 Sylt, German North Sea coast
- E. kostei** Tschaschel,1979
 Sylt, German North Sea coast
- E. listense** Tschaschel,1979
 Sylt, German North Sea coast

<i>E. longirostrum</i> Tschaschel,1979 =? <i>Wigrella amphora</i> Remane,1929 Sylt, German North Sea coast, Baltic?	<i>S. curvata</i> Lie-Pettersen,1905 Bergen, Baltic, North Sea
<i>E. marinum</i> (Dujardin,1841) =? <i>E. similimum</i> Remane,1929 = <i>Furcularia marina</i> Dujardin,1841 (TM) Tvärminne, W & N Norway, Roscoff, Sylt	<i>S. gyrina</i> Hood,1887 Tvärminne, Bergen, North Sea
<i>E. matthesi</i> Remane,1949 German & Danish Baltic Sea	<i>S. litoralis</i> Rousselet,1902 W Sweden, Bergen, Baltic, Scotland, Plymouth, Zuiderzee
<i>E. eristes</i> Herring & Myers,1928 Kieler Bucht	<i>S. neapolitana</i> Rousselet,1902 W Sweden, Baltic, North sea, Bergen
<i>E. mustela</i> (Milne,1885) Cosmopolitan, in fresh & brackish waters	<i>S. oblonga</i> Ehrenberg,1831 Tvärminne
<i>E. obesum</i> Tschaschel,1979 Sylt, German North Sea coast	<i>S. tremula</i> (O.F. Müller,1786) Tvärminne, in fresh-water rock-pools
<i>E. pachidum</i> (Gosse,1887) W & N Norway, Baltic, North Sea	<i>S. triophthalma</i> Lauterborn,1894 W Sweden, Baltic, Bergen, Roscoff, Plymouth, Zuiderzee
<i>E. pachypus</i> Wulfert,1936 German Baltic & North Sea coasts	<i>S. vorax</i> Rousselet,1902 W & N Norway, Tvärminne, North Sea, Roscoff, Plymouth
<i>E. permutandum</i> Tschaschel,1979 Sylt, German North Sea coast	<i>S. tavina</i> Hood,1883 Baltic, North Sea, Roscoff
<i>E. psammophilum</i> Althaus,1957 Sylt, German North Sea coast	<i>S. pectinata</i> Ehrenberg,1832 (TM) Europe, in brackish water
<i>E. remanei</i> Voigt,1957 Helgoland, German Baltic & North Sea coasts	<i>S. monopus</i> Plate,1889 Baltic, North Sea
<i>E. rousseleti</i> (Lie-Pettersen,1905) Bergen, Roscoff, Kiel, Scotland	<i>S. grimppei</i> Remane,1929 Baltic, England
<i>E. sacculiforme</i> Tschaschel,1979 Sylt, German North Sea coast	<i>S. atlantica</i> Zelinka,1907 North Sea, North Atlantic
<i>E. spattatum</i> Wulfert,1936 Poland, Germany	<i>S. johanseni</i> Herring,1921 Baltic
<i>E. striatum</i> Althaus,1957 Sylt, German North Sea coast	<i>S. fennica</i> Rousselet,1909 Baltic
<i>E. subterraneum</i> (Remane,1949) Kieler Bucht, German Baltic coast	<i>S. razelmi</i> Rudescu,1960 =? <i>S. spec.</i> Remane,1937 Black Sea, Kiel?
<i>E. tectipes</i> Remane,1949 German Baltic & North Sea coasts	<i>S. bicornis</i> Smith,1904 Zuiderzee
<i>E. villosum</i> Herring & Myers,1928 Baltic rock-pools	<i>Polyartha</i> Ehrenberg,1834 (TM <i>P. vulgaris</i> Carlin,1943 - fresh water)
<i>E. wierzejskii</i> Tschaschel,1979 Sylt, Helgoland, Kieler Bucht, German Baltic & North Sea coasts	<i>P. remata</i> (Skorikov,1896) =? <i>P. platyptera</i> Ehrenberg,1834? Bohuslän?
<i>E. myersi</i> (Wulfert,1936) Kieler Bucht	<i>Ploesoma</i> Herrick,1885 (TM <i>P. lenticulare</i> Herrick,1885 - fresh water)
<i>Wigrella</i> Wiszniewski,1932 (TM <i>W. depressa</i> Wiszniewski,1932 - fresh water)	<i>P. truncatum</i> (Levander,1894) Bohuslän
<i>W. amphora</i> (Remane,1929) =? <i>Encentrum longirostrum</i> Tschaschel,1974 Kieler Bucht	<u>FLOSCULARIACEAE</u> Remane,1933
<i>Wierzejskella</i> Wiszniewski,1934 (TM <i>Encentrum velox</i> Wiszniewski,1932 - fresh water)	<u>Testudinellidae</u> Bartos,1959
<i>W. marina</i> Remane,1949 Helgoland, Kieler Bucht	<i>Testudinella</i> Bory de St. Vincent,1826
<i>W. subterranea</i> Remane,1949 Kieler Bucht	<i>T. clypeata</i> (O.F. Müller,1786) = <i>Brachionus clypeatus</i> O.F. Müller,1786 (TM) Tvärminne, Bergen, Roscoff, Zuiderzee
<i>Dicranophorus</i> Nitzsch,1827	<i>T. obscura</i> Althaus,1957 Roscoff
<i>D. forcipatus</i> (O.F. Müller,1773) = <i>Cercaria forcipata</i> O.F. Müller,1773 (TM) Cosmopolitan, in fresh & brackish water	<i>T. patina</i> (Hermann,1783) Tvärminne, in fresh-water rock-pools
<i>Albertia</i> Dujardin,1838 (TM <i>A. vermiculus</i> Dujardin,1838 - fresh water)	<i>Pompholyx</i> Gosse,1851 (TM <i>P. complanata</i> Gosse,1851 - fresh water)
<i>A. crystallina</i> Schultze,1851 Baltic, in the intestine of the oligochaete <i>Paranais littoralis</i>	<i>P. sulcata</i> (Hudson,1885) England?, in fresh & brackish water
<i>A. naidis</i> Bousfield,1886 =? <i>A. intrusor</i> Gosse,1886 Cosmopolitan?, in the intestine of <i>Nais barbata</i> , <i>N. elinguis</i> , <i>N. variabilis</i> , <i>Stylaria lacustris</i> & <i>Ripistes parasita</i> .	<i>Filinia</i> Bory de St. Vincent,1824
<i>A. soyeri</i> Coineau & Kunst,1964 France?, in the intestine of <i>Nais elinguis</i>	<i>F. cornuta</i> (Weisse,1847) Cosmopolitan, in fresh & brackish waters
<i>Erignatha</i> Herring & Myers,1928 (TM <i>Diglena clastopis</i> Gosse,1886 - fresh water)	<i>F. brachiata</i> (Rousselet,1901) Cosmopolitan, in fresh & brackish waters
<i>E. thienemanni</i> Remane,1929 Kieler Bucht, Roscoff	<u>Hexarthridae</u> Bartos,1959
<u>Asplanchnidae</u> Herring & Myers,1926	<i>H. mira</i> (Hudson,1871) = <i>Pedalion mirum</i> Hudson,1871 (TM) ?, in fresh & brackish waters
<i>Asplanchna</i> Gosse,1850	<i>H. fennica</i> (Levander,1892) Tvärminne, Zuiderzee, Cosmopolitan
<i>A. priodonta</i> Gosse,1850 (TM) Europe, in fresh & brackish waters	<i>H. oxyuris</i> (Sernov,1903) E Baltic
<u>Synchaetidae</u> Remane,1933	<u>Flosculariidae</u> Herring,1913
<i>Synchaeta</i> Ehrenberg,1834	<i>Floscularia</i> Cuvier,1798 (TM <i>Serpula ringens</i> Linnaeus,1758 - fresh water)
<i>S. baltica</i> Ehrenberg,1834 Sweden, Tvärminne, Bergen, North Sea	<i>F. melicerta</i> (Ehrenberg,1832) Cosmopolitan, in fresh & brackish waters
<i>S. cecilia</i> Rousselet,1902 W Sweden, North Sea, Tvärminne, Bergen, Roscoff	<u>Conochiliidae</u> Remane,1933
	<i>Conochilus</i> Ehrenberg,1834

C. hippocrepis (Schrank,1830)
Europe, in fresh & brackish waters

COLLOTHECACEAE Remane,1933

Collotheidae Bartos,1959

Collothea Herring,1913

C. ornata (Ehrenberg,1832)
Europe, in fresh & brackish waters

C. campanulata (Dobie,1849)
= *Floscularia campanulata* Dobie,1849 (TM)
Europe, in fresh & brackish waters

C. mutabilis (Hudson,1885)
Europe, in fresh & brackish waters

ECHIURA Sedgwick,1898 (10-11 European Atlantic spp.)

XENOPNEUSTIDA Fisher,1946

* Not represented in the area.

HETEROMYOTIDA Fisher,1946

* Not represented in the area.

BONELLIIDA

Bonelliidae Baird,1868

Bonella Rolando,1821

B. viridis Rolando,1821 (TM)
Kattegatt - Bohuslän - Skagerrak - Trondheimsfjorden, S & W coasts, off Ireland, Mediterranean

Maxmuelleria Bock,1942 (TM *Thalassema gigas* Max Müller,1852 - Mediterranean)

M. lankesteri (Herdman,1898)

= *Thalassema lankesteri* Herdman,1898

* May possibly be conspecific with the Mediterranean *M. gigas* (M. Müller,1852)
Kattegatt - Bohuslän - Skagerrak (off Arendal), W Scotland, Irish Sea, NW Spain

M. faex (Selenka,1885)

= *Thalassema faex* Selenka,1885

Norwegian coast from Sogn - Trondheimsfjorden

Hamingia Danielssen & Koren,1881

H. arctica Danielssen & Koren,1881 (TM)

Norwegian coast from Stavanger and northwards

Amalosoma Fischer,1948 (TM *Acanthohammingia paradoxa* Fisher,1946 - Japan)

A. eddysonense Stephen,1956

Plymouth, Biscay

ECHIURIDA

Echiuridae de Blainville,1827

Echiurinae Monro,1927

Echiurus Guérin,1831

* Guérin did not put -Méneville to his last name until 1836.

E. echiurus (Pallas,1767)

= *Lumbricus echiurus* Pallas,1767 (TM)

Bohuslän - Öresund, all Norway, W North Sea

E. abyssalis Skorikow,1906

West coast of Ireland in deep water, Capri

Thalasmatinae Monro,1927

Thalassema Lamarck,1801

T. thalasseum (Pallas,1766)

= *Lumbricus thalasseum* Pallas,1766 (TM)

= *Thalassema neptuni* Gærtner, in Pallas,1774

Plymouth, S Wales, Ireland, France, Mediterranean

T. arcassonense Cuénot,1902

Bay of Arcachon

Anellassorhynchus Annandale,1922 (TM *Thalassema branchiorhynchus* Annandale & Kemp,1915

- India)

A. indivisus (Sluiter,1900)

Off W coast of Spain in 90 m. dept.

Species inquirenda(i.e.: it is questionable if this even is an Echiurid)

Sactosoma Danielssen & Koren,1881

S. vitreum Danielssen & Koren,1881 (TM)

Between Faeroes and Norway in 2222 m

SIPUNCULA Delle Chiaje,1823

* Ribosomal 18S DNA sequences indicate a close relationship with Annelida.

PHASCOLOSMATIDEA

ASPIDOSIPHONIDA

= **ASPIDOSIPHONIFORMES** Cutler & Gibbs,1985

Aspidosiphonidae Baird,1868, emend. Stephen & Edmonds,1972

Aspidosiphon Diesing,1851

A. (Aspidosiphon) muelleri Diesing,1851 (TM)

= *A. clavatus* : Auctt. & possibly (de Blainville,1827)

= *Sipunculus (Phascolosoma) scutatus* J. Müller,1844

= *A. mirabilis* Théel,1875

= *A. jukestii* Baird,1873

= *A. armatum* Danielssen & Koren,1880

= *Phascolosoma radiata* Alder,1860

Bohuslän, Bergen - Trondheimsfjorden, Shetlands, Mediterranean, N & W Africa, Azores, Canaries

A. (Akrikos Cutler & Cutler,1989) *zinni* Cutler,1969
Biscay in 600-3000m

PHASCOLOSMATIDAE

= **PHASCOLOSMATIFORMES** Cutler & Gibbs,1985

Phascolosomatidae Stephen & Edmonds,1972

Phascolosoma Leuckart,1828

= *Phymosomum* de Quatrefages,1865

= *Phymosoma* Selenka & al.,1883

= *Physcosoma* Selenka,1897

P. granulatum Leuckart,1828 (TM)

= *Physcosoma granulatum* : Auctt.

Bergen area, Shetlands, Hebrides, Orkneys, W Ireland, Mediterranean, Azores, Cape Verde

SIPUNCULIDEA

SIPUNCULIDA

Sipunculidae Rafinesque,1814

Sipunculus Linnaeus,1766

= *Phallosooma* Levinsen,1883

S. nudus Linnaeus,1766 (TM)

Bergen area, S North Sea, Plymouth, Ireland, France

S. norvegicus norvegicus Danielssen,1868

= *Sipunculus priapuloides* Koren & Danielssen,1875

= *Phallosooma priapuloides* : Levinsen,1883.

Kattegatt - Lofoten, N North Sea

Siphonosoma Spengel,1912 (TM *Phascolosoma australe* Keferstein,1865 - Australia)

S. arcassonense (Cuénot,1902)

Acahorn Bay

GOLFINGIDA

Golfingidae Stephen & Edmonds,1972

Golfingia Lankester,1885

G. elongata (Keferstein,1863)

= *Phascolosoma elongatum* Keferstein,1863

= *Phascolosoma charcoti* Héribel,1906

= *Phascolosoma derjugini* Gadd,1911

Kattegatt - Bohuslän - Skagerrak - Sogn, Plymouth, Iberian Peninsula, Mediterranean

G. margaritacea margaritacea (M. Sars,1851)

= *Sipunculus margaritaceus* M. Sars,1851

= *Phascolosoma margaritacea fimmarchica* Théel,1905

= *Phascolosoma margaritaceum* forma *sibirica* Théel,1905

= *Phascolosoma hanoni* Danielssen & Koren,1881

= *Phascolosoma nordenskjöldi* Théel,1911

= *Phascolosoma trybomi* Théel,1905

= *Golfingia cantabrensis* Edmonds,1960

= *Phascolosoma profundum* Roule,1898

Kattegatt - Bohuslän - Skagerrak - all Norway, W Scotland, SW Britain

G. vulgaris vulgaris (Blainville,1827)

= *Sipunculus vulgaris* de Blainville,1827 (TM)

= *Phascolosoma vulgare* var. *astuta* (Selenka,1885)

= *Phascolosoma kolense* Gadd,1911

Kattegatt - Bohuslän - Skagerrak, all Norwegian coast, North Sea, Britain, Mediterranean

G. muricauddae (Southern,1913)

= *Phascolosoma muricatum* Southern,1913

Off SW Ireland in 475-1088m

G. iniqua (Sluiter,1912)

= *Phascolosoma iniquum* Sluiter,1912

= *Phascolosoma mutabile* Southern,1913

= *Phascolosoma rugosum* Southern,1913

SW Ireland in 914-1242m, NE Atlantic in 500-1800m

G. murinæ Cutler,1969

Biscay in ca 1000-4800m

Nephosoma Pergament,1940

N. minuta (Keferstein,1863)

=? *Sipunculus johnstoni* Forbes,1841

= *Phascolosoma minutum* Keferstein,1863

= *Golfingia (Phascoloides) minuta* (Keferstein)

= *Phascolosoma sabellariae* Théel,1905

= *Phascolosoma anceps* Théel,1905

Kattegatt - Bohuslän - Skagerrak - all Norwegian coast, North Sea, Shetlands, Britain, Bretagne

N. rimicola (Gibbs,1973)

= *Golfingia (Phascoloides) rimicola* Gibbs,1973

Devon, Cornwall, N Spain

N. lilljeborgii (Danielssen & Koren,1881)

= *Phascolosoma lilljeborgii* Danielssen & Koren,1881

= *Golfingia (Phascoloides) lilljeborgii* (Danielssen & Koren,1881)

= *Oncimesoma glaciata* Danielssen & Koren,1880

= *Nephosoma mariniki* Pergament,1940 (TM - Arctic Seas)

Norwegian coast, Spitzbergen, Iceland in bathyal depths

N. eremita (M. Sars,1851)

= *Sipunculus eremita* M. Sars,1851

= *Golfingia (Phascoloides) eremita* (M. Sars,1851)

= *Sipunculus (Phascolosum) boreale* Keferstein,1865

= *Phascolosoma digitatum* Théel,1875

Bergen - NE Finnmark, Arctic Seas

N. abyssorum (Koren & Danielssen,1875)

= *Phascolosoma abyssorum* Koren & Danielssen,1875

- = *Golfingia (Phascoloides) abyssorum* (Koren & Danielssen,1875)
 = *Phascolosoma incompositum* Sluiter,1912
 W Norway (Bergensfj.), Helgoland
- N. diaphanes diaphanes** (Gerould,1913)
 = *Phascolosoma diaphanes* Gerould,1913
 = *Golfingia diaphanes* (Gerould,1913)
 = *Phascolosoma cinctum* Gerould,1913
 = *Phascolosoma improvius* Théel,1905
 Bohuslän, Oslofjorden - Trondheimsfjorden, Biscay in deep water, Cosmopolitan cold water species, chiefly from bathyal & abyssal depths
- N. diaphanes corrugatum** Cutler & Cutler,1986
 = *Golfingia schuttei* : Murina,1964, non (Augener,1903)
 Atlantic & Pacific Oceans, Mediterranean & Red Sea in 80-5900m
- N. delagei** (Hérubel,1903)
 = *Phascolosoma delagei* Hérubel,1903
 = *Golfingia (Phascoloides) delagei* (Hérubel,1903)
 * sensu Gibbs,1973 & Cutler & Cutler,1986 is this species indeterminable (incertae sedis)
 Ushant, Biscay
- N. bulbosum** (Southern,1913)
 = *Phascolosoma bulbosum* Southern,1913
 = *Golfingia (Phascoloides) bulbosa* (Southern,1913)
 Off SW Ireland in 957-1088m, off NE USA in bathyal depths
- N. constrictum** (Southern,1913)
 = *Phascolosoma constrictum* Southern,1913
 = *Golfingia (Phascoloides) constricta* (Southern,1913)
 Off SW Ireland in 1143-1317m, NE Atlantic in 1900-4000m
- N. wodjanizkii** (Murina,1973) *elisae* (Murina,1977)
 = *Golfingia (Phascoloides) elisae* Murina,1977
 Continental slope of NE Atlantic (43°-58°) in 1600-2300m
- N. constricticervix** (Cutler,1969)
 = *Golfingia constricticervix* Cutler,1969
 Biscay in more than 2000m
- N. flagriferum** (Selenka,1885)
 = *Phascolosoma flagriferum* Selenka,1885
 = *Golfingia flagrifera* (Selenka,1885)
 Biscay in depths greater than 1500m, N Atlantic in abyssal depths
- N. capilleiforme** (Murina,1973)
 = *Golfingia capilleiformis* Murina,1973
 Atlantic & Pacific ocean in 920-4900m
- Thysanocardia** Fisher,1950
- T. procerata** (Möbius,1875)
 = *Phascolosoma procerum* Möbius,1874 (TM)
 = *Golfingia procerata* : Auct.
 * May be a junior synonym of *T. catharinæ* (Grube,1868)
 Kattegatt - Skagerrak - Trondelag S of Trondheimsfjorden, N North Sea, W Scotland, W Ireland, Milford Haven
- Apionsoma** Sluiter,1902 (TM) *A. trichocephalus* Sluiter,1902 - Malaya
- A. capitata** (Gerould,1913)
 = *Physcosoma capitatum* Gerould,1913
 = *Physcosoma abyssorum* Southern,1913
 = *Phascolosoma abyssorum* (Southern,1913)
 SW Ireland in 1006-1096m, Biscay in 2800-4200m
- Phascolionidae** Cutler & Gibbs,1985
- Phascolion** Théel,1875
- P. strombus strombus** (Montagu,1804)
 = *Sipunculus strombi* Montagu,1804 (TM)
 = *Phascolosoma intermedium* Southern,1913
 Kattegatt - Bohuslän - all Norway, all around Britain, Mediterranean
- P. tuberculatum** Théel,1875
 = *P. pallidum* Koren & Danielssen,1877
 = *P. hirondellei* Sluiter,1900
 = *P. magadorensis* Sluiter,1912
 Bohuslän, Skagerrak, Bergen, Trondheimsfjord
- P. (Montuga) pacificum** Murina,1957
 Off W European coast in depths greater than ca 1000m
- P. (Montuga) lutense** Selenka,1885
 = *P. (Montuga) canum* Cutler & Cutler,1980
 Biscay in depths of more than 2000m
- Onchnesoma** Koren & Danielssen,1875
- O. squamatum squamatum** (Koren & Danielssen,1875)
 = *Phascolosoma squamatum* Koren & Danielssen,1875
 N Kattegatt, Skagerrak, North Sea, Oslofjord - Lofoten, Britain, Portugal, Mediterranean
- O. steenstrupii steenstrupii** Koren & Danielssen,1875 (TM)
 Kattegatt - Skagerrak - W Finnmark, North Sea, Britain, Portugal, Mediterranean, Ivory Coast
- O. magnibathra** Cutler,1969
 Off NW European coast in depths greater than 2000m
- POGONOPHORA** K.E. Johansson, 1937
 * More and more evidence seems to indicate an Annelid relationship.
- AFRENULATA** M. Webb,1969
 * Not represented in the area.
- FRENULATA** M. Webb,1969
- ATHECANEPHRIDA** Ivanov,1955
- Oligobrachiidae** Ivanov,1957
- Oligobrachia** Ivanov,1957
- O. webbi** Brattegaard,1966
 NW Tromsö in 270m, Varangerfjord area
- O. ivanovi** Southward,1959
 SW Engl. Channel in 1343-1435m
- O. gracilis** Southward,1978
 N Biscay in 720-2215m
- Siboglinidae** Caullery,1914
- Nereilinum** Ivanov,1961
- N. murmanicum** Ivanov,1961
 N & NE Finnmark, Barents Sea in 170-325m, Norw. Sea in 1280m
- Siboglinum** Caullery,1914
- S. brevicephalum** Flügel,1990
 Nor. Sea (1280 m)
- S. ekmani** Jägersten,1956
 Skagerrak and Norwegian fjords - N Finnmark in 396-1288m, SW Britain, Biscay
- S. fiordicum** Webb,1963
 Skagerrak and Norwegian fjords - N Finnmark in 30-600m
- S. poseidonii** Flügel & Langhof,1983
 Skagerrak (off Arendal - off Kristiansand) in 230-294m
- S. norvegicum** Ivanov,1960
 Off W Shetlands & off Norwegian coast in 120 & 1165m
- S. holmei** Southward,1963
 W Scotland, SW Ireland in 100-1300m
- S. atlanticum** Southward & Southward,1958
 NE Atlantic between SW Britain & N Spain in 956-1324m
- S. inerme** Southward & Southward,1958
 NE Atlantic off SW Britain in 522-1288m
- S. lacteum** E.C. Southward,1963
 NE Atlantic in 1105-1250m
- S. leucopleurum** Flügel & Callsen-Cencic,1993
 Biscay in 1138 m.
- THECANEPHRIDA** Ivanov,1955
- Polybrachiidae** Ivanov,1952
- Diplobrachia** Ivanov,1960
- D. capillaris** (Southward,1959)
 NE Atlantic off SW Britain in 550-1700m
- Selerolinidae** Webb,1964
- Selerolinum** Southward,1961
- S. brattstroemi** Webb,1964
 W Norwegian fjords in 100-870m in rotting wood, ropes, etc.
- HEMICORDATA** Bateson,1885
- ENTEROPNEUSTA** Gegenbaur,1870
 = **HELMINTHOMORPHA**
- Ptychoderidae** Spengel,1893
- Glossobalanus** Spengel,1901
- G. sarniensis** (Köhler,1886)
 ?Skagerrak off Arendal, S North Sea
- G. marginatus** Meek,1922
 Skagerrak off Arendal - Egersund, North Sea
- G. minutus** (Kowalevsky,1866)
 Portuguese coast, Mediterranean
- Balanoglossus** Delle Chiaje,1829
- B. clavigerus** Delle Chiaje,1829
 England, Portugal
- Spengeliidae** Willey,19??
 * Not represented in the area.
- Harrimanidae** Spengel,1901
- Harrimania** Ritter,1900
- H. kupfferi** (von Willemoes-Suhm,1871)
 Öresund - Bohuslän - Skagerrak, Trondheimsfjorden
- Saccoglossus** Schimkewitsch,1892
 = *Dolichoglossus* Spengel,1893
- * Along NW Atlantic shores two species, *S. kowalevskii* Agassiz,1873 & *S. bromophenolosus* King, Giray & Kornfield,1994 may be found.
- S. pygmaeus** Hinrichs & Jacobi,1938
 Helgoland
- S. ruber** Tattersall,1905
 = *S. serpentinus* Assheton,1908
- = *S. cambrensis* Brambell & Cole,1939
 Galway, Wales, Sound of Mull
- S. horsii** Brambell & Goodhart,1941
 = *S. kowalevskii* Auct. non Agassiz,1873
 Öresund?, Solent (Engl. Channel)
- S. mereschkowskii** Wagner,1???
- off N Russia
- Stereobalanus** Spengel,1901

S. canadensis (Spengel,1893)
Northern Scotland, Voring Plateau (deep water) in the Norwegian Sea

Protoglossidae Dawyodoff,1966
* This family is often included in Harrimaniidae.

Protoglossus van der Horst,1935
= *Protobalanus* Caullery & Mesnil,1904, *non* Hall & Clarke,1888 , *nec* Whitfield,1889 (Cirripedia)

P. koehleri (Caullery & Mesnil,1900)
Netherlands, Isle of Man

P. n. sp. Silén,1950
* Possibly identical with *P. koehleri* (Caullery & Mesnil)
Gulfmarnfjord area

PTEROBRANCHIA R. Lankester,1878

Cephalodiscidae

Cephalodiscus McIntosh,1882

C. levinseni Harmer,1905
?NE Atlantic?, off Japan & Corea

C. gracilis McIntosh,1882
Bermudas

Atubariidae

* The only species known was found by Sato from Japan.

Rhabdopleuridae

Rhabdopleura Allman,1869
= *Haliotopha* M. Sars,1869 (nom. nud.)

R. normani Allman,1869
= *Rhabdopleura mirabilis* M. Sars, *in* G.O. Sars,1872
@ often on branches of dead *Lophelia*
Bohuslän - Loftalen, ?Spitzbergen, Shetlands, Faeroes, Iceland, Barents Sea, W Greenland, Biscay, Mediterranean, Azores

R. compacta Hincks,1880
@ often on concave side of dead bivalve shells in 23-100 m
Plymouth, NE Ireland, N Bretagne

PLANCTOSPHAEROIDEA Van der Horst,1936

Planctosphaera Spengel,1932

P. pelagica Spengel,1932
Biscay in 270 m, Bermudas

CHAETOGNATHA Leuckart,1854

SAGITTOIDEA Claus & Grobben,1905

PHRAGMOPHORA Tokioka,1965

Spadellidae Tokioka,1965

Spadella Langerhans,1880

* World-wide 10 species are known, of which the 3 species in this list described by Casanova and the antarctic *S. antarctica* Casanova,1991 always have posterior teeth. One species *S. japonica* Casanova, 1993 do not have posterior teeth. The other five species, including *S. moretonensis* Johnston & Taylor, 1919 (Australia), *S. angulata* Tokioka,1951 (Japan), *S. gaetanoi* Alvarino,1978 (Hawaii) & *S. bradshawi* Bieri,1974 (California) may or may not have posterior teeth.

S. cephaloptera (Busch,1851)
Öresund - Bohuslän - Tröndelag

S. birostrata Casanova,1987
Gibraltar in 150-555m

S. equidentata Casanova,1987
Portugal in 452m

S. ledoyerii Casanova,1986
Marseille (in submarine cave)

Paraspadella Salvini-Plawen,1987
= *Gephyropadella* Salvini-Plawen,1987

* Having adhesive organs in contrast to *Spadella* which is restricted to species lacking adhesive organs. Not known from Scandinavia? The species: *P. schizophora* (Conant,1895) (Bahamas & Florida), *P. anops* Bowman & Bieri,1989 (Bahamas), *P. caecifera* (Salvini-Plawen,1987), *P. gotoi* (Casanova,1990) (Japan), *P. hummelincki* (Alvarino,1970) (which possibly is a synonym of *P. pulchella* Owre) (Puerto Rico), *P. johnstoni* (Mawson,1944) (Australia), *P. legazpicensis* (Alvarino,1981) (Marshall Islands), *P. nana* (Owre,1963) (Florida), *P. pinumukatharos* (Alvarino,1987) (California), *P. pulchella* (Owre,1963) (Puerto Rico) & *P. sheardi* (Mawson,1944) (Australia).

Bathypadella Tokioka,1939 (TM *Bathypadella edentata* Tokioka,1939 - Japan)

* Only one specimen of one species known from this genus.

Eukrohnidae Tokioka,1965

Eukrohnia Ritter-Záhony,1909

E. hamata (Möbius,1875)
Bohuslän, Bergen, Iceland, Scotland, Biscay

E. bathypelagica Alvarino,1962
* Aurich, 1970 is considering this species to be a breeding stage of *E. hamata* (Möbius)
Bohuslän, Bergen

E. fowleri Ritter-Záhony,1909
Norwegian Sea, off Ireland, Biscay

Heterokrohnia Ritter-Záhony,1911
= *Archeterokrohnia* Casanova,1986

H. palpifera (Casanova,1986)
= *Archeterokronia palpifera* Casanova,1986
W Mediterranean

APHRAGMOPHORA Tokioka,1965

CTENODONTINA Tokioka,1965

Pterosagittidae Tokioka,1965

Pterosagitta Costa,1869

P. draco (Krohn,1853)

Biscay

Sagittidae Claus & Grobben,1905

Sagitta Quoy & Gaimard,1827

S. bipunctata Quoy & Gaimard,1827

Atlantic west of Ireland, Biscay

Subgenus: *Parasagitta* Tokioka,1965

* Heterogeneous taxon.

S. (P.) elegans elegans Verrill,1873

Bohuslän - Baltic?, The Oslo Fjord, Britain

S. (P.) elegans baltica Ritter-Záhony,1911

Kattegatt - Baltic, Scotland

S. (P.) elegans arctica Aurivillius,1896

Norwegian Sea - off Ireland - Arctic

S. (P.) setosa J. Müller,1847

* Neritic

Bohuslän - Baltic, Netherlands, Britain, Mediterranean

S. (P.) friderici Ritter-Záhony,1911

W Scotland - Biscay

Subgenus: *Flaccisagitta* Tokioka,1965

S. (F.) hexaptera d'Orbigny,1836

= *Sagitta exaptera* d'Orbigny,1836

Faeroes, Shetlands, Atlantic west of Ireland, Biscay

S. (F.) enflata Grassi,1881

W Scotland, Biscay

Subgenus: *Pseudosagitta* Germain & Joubin,1912

S. (P.) maxima (Conant,1896)

Skagerrak - N North Sea, Iceland, Norwegian Sea - Arctic

S. (P.) lyra lyra Krohn,1853

N North Sea, off Ireland, Biscay

S. (P.) lyra gazellae : Auct. *non* Ritter-Záhony,1909

N North Sea?, Biscay

Subgenus: *Serratosagitta* Tokioka & Pathansali,1963

S. (S.) serratodentata serratodentata Krohn,1853

W Britain, Ireland, Biscay, Mediterranean

S. (S.) tasmanica Thomson,1947

Skagerrak, N North Sea, Norwegian Sea - N Norway, W Britain - NW Africa

Subgenus: *Caecosagitta* Tokioka,1965

S. (C.) macrocephala Fowler,1905 (TM of *Caecosagitta* Tokioka,1965))

Norwegian Sea, off Scotland & Ireland, Biscay

Subgenus: *Mesosagitta* Tokioka,1965

S. (M.) decipiens Fowler,1905

Norwegian Sea, Scotland, Biscay

S. (M.) minima Grassi,1881

Biscay

S. (M.) batava Biersteker & van der Spoel,1966

* Neritic

Netherlands

Subgenus: *Solidosagitta* Tokioka,1965

S. (S.) zetesios Fowler,1905

* Possibly a synonym of *S. planctonis* Steinhaus,1896

Norwegian Sea, Biscay

S. (S.) planctonis Steinhaus,1896

Atlantic west of Ireland, Biscay

FLABELLODONTINA Tokioka,1965

Krohnittidae Tokioka,1965

Krohnitta Ritter-Záhony,1910

K. subtilis (Grassi,1881)

Atlantic west of Ireland, Biscay

XENOTURBELLIDA

* This "phylum" will hopefully be reduced to a lower taxon within a near future, when DNA sequences from the only species have been compared to other sequences from other phyla.
According to Jondelius (pers. comm), DNA sequence data speaks in favour of a relationship with the "pseudocoelomate" taxa.

Xenoturbellidae

Xenoturbella Westblad,1949

X. bocki Westblad,1949 (TM)

Gulmarsfjord, Koster Channel, Bergen, Scotland
